Advances in Geography & Environmental Research



Editors

Dr. Vijay Chandrakant Ithape
Dr. Arpana Premi Lina Oraon
Dr. Balasaheb Raghunath Chakor
Dr. K. Prabhu

Nature Light Publications

An International Edited Book *ISBN-978-93-49938-93-9*

Advances in Geography and Environmental Research

Editors

Dr. Vijay Chandrakant Ithape
Associate Professor
Department of Geography
Shri Chhatrapati Shivaji Mahavidyalaya, Shrigonda, (MH), India**.**

Dr. Arpana Premi Lina Oraon
Dean of the School of Tribal Resource Management
Director, Research and Development Cells at Kalinga Institute of Social Sciences,
Deemed to be University, Odisha., India.

Dr. Balasaheb Raghunath Chakor
Assistant Professor and Head
Department of Geography
Krantiveer Vasantrao Narayanrao Naik Shikshan Prasarak Sanstha's, Arts,
Commerce and Science College, Nashik, (MH), India.

Dr. K. Prabhu
Assistant Professor
Department of Physics
Academy of Maritime Education and Training
(AMET University), Kanathur, Chennai, India.

Published By

Nature Light Publications, Pune

© Reserved by Editor's

Advances in Geography and Environmental Research

Editors

Dr. Vijay Chandrakant Ithape
Dr. Arpana Premi Lina Oraon
Dr. Balasaheb Raghunath Chakor
Dr. K. Prabhu

First Edition: June, 2026

An International Edited Book

ISBN- 978-93-49938-93-9

Published by:
Nature Light Publications, Pune
309 West 11, Manjari VSI Road, Manjari Bk.,
Haveli, Pune- 412 307.
Website: www.naturelightpublications.com
Email: naturelightpublications@gmail.com
Contact No: +91 9822489040 / 9922489040

The editors/Associate editors/publisher shall not be responsible for originality and thought expressed in the book chapter/ article. The author shall be solely held responsible for the originality and thoughts expressed in their book chapter or article.

Preface

Geography and environmental research play a vital role in understanding the complex interactions between natural systems, human societies, and the rapidly changing global environment. In the face of climate change, resource depletion, biodiversity loss, urban expansion, and increasing environmental risks, interdisciplinary research has become essential for developing sustainable solutions that balance ecological integrity, economic growth, and social well-being. Advances in geospatial technologies, environmental assessment methods, and sustainable resource management continue to reshape the way we study and respond to these global challenges.

This edited volume, Advances in Geography and Environmental Research, brings together contributions from researchers, academicians, environmental scientists, and geographers whose work reflects the diverse dimensions of contemporary geographical and environmental studies. The chapters collectively provide valuable insights into environmental sustainability, land-use planning, natural resource management, water security, rural development, and environmental governance.

The volume begins by exploring marine and freshwater ecosystems, emphasizing their ecological significance and the need for sustainable management to secure future environmental resilience. The discussion on the arid lands of India examines the relationships among geodiversity, livelihoods, and development, highlighting the unique environmental challenges and opportunities of dryland regions. A comparative study on nano urea-based fertilization presents emerging perspectives on sustainable agricultural practices, evaluating both the advantages and limitations of innovative fertilizer technologies.

The book further examines spatial planning for climate-smart communities, demonstrating how resilient land-use planning and environmental management can enhance adaptation to climate change and reduce environmental vulnerability. The chapter on Environmental Impact Assessment

(EIA) discusses the integration of ecological conservation, social responsibility, and economic development, reinforcing the importance of evidence-based decision-making in sustainable development planning.

Practical dimensions of environmental management are explored through the evaluation of water supply infrastructure and drinking water quality in Shrigonda Municipality, providing valuable insights into urban water governance and public health. Complementing this, the chapter on the honest implementation of the Environmental Impact Assessment Notification, 2006, highlights the importance of transparent regulatory frameworks and effective environmental governance in achieving sustainable development goals.

The volume concludes with an analytical study of literacy in rural settlements of Daund Tahsil, Pune District, illustrating how demographic and socio-economic characteristics influence regional development and contribute to geographical research on rural transformation.

Collectively, the chapters presented in this volume demonstrate the interdisciplinary nature of geography and environmental research by integrating physical geography, human geography, environmental science, sustainability studies, policy analysis, and technological innovation. They offer practical knowledge, scientific evidence, and policy perspectives that are relevant to both academic research and real-world environmental management.

We express our sincere appreciation to all contributing authors for sharing their valuable research and expertise. We are equally grateful to the reviewers and the publishing team for their constructive guidance and dedicated support throughout the development of this volume.

It is our hope that Advances in Geography and Environmental Research will serve as a valuable reference for students, researchers, academicians, environmental professionals, planners, policymakers, and all readers interested in promoting sustainable development through innovative geographical and environmental research.

Editors

Advances in Geography and Environmental Research

Table of Content

Sl. No.	Title and Authors	Page No.
1	**Exploring Marine and Freshwater Ecosystems for Future Sustainability** *Dr. Muthurajan S, Immanuel Prabaharan S, Aswini M Deivanayaki R*	01 - 05
2	**The Arid Lands Environment of India: Geodiversity, Livelihoods, and the Paradox of Development** *Prof. (Dr.) Mukesh K. Sharma, Prof. (Dr.) Fareeda Hasani Dr. Babita, Dr. Roopendra Singh*	06 - 22
3	**Nano Urea-Based Fertilization: A Comparative Study of Advantages and Disadvantages** *Jitendra Pal Singh*	23 - 28
4	**Spatial Planning for Climate Smart Communities Through Resilient Land Use and Environmental Management** *Dr. Prajakta Vinod Jadhav*	29 - 45
5	**Environmental Impact Assessment Integrating Ecological Conservation Social Responsibility and Economic Development** *Suchandana Dutta*	46 - 58
6	**Evaluation of Water Supply Infrastructure and Drinking Water Quality in Shrigonda Municipality** *Bapu D. Deokar*	59 - 70
7	**Honest Implementation of the Environment Impact Assessment Notification 2006: Roadway Towards Sustainable Development** *Dr. Umesh S. Aswar*	71 - 77
8	**Literacy of Rural Settlement in Daund Tahsil, Pune district (MH): An Analytical Study** *Dr. Parkhe Sakharam Baban*	78 - 81
9	**Spatial Analysis of Groundwater Availability and Quality in India: A GIS-Based Assessment** *Mr. Agastirishi Toradmal, Dr. Vilas Vasant Patil*	82 - 92

Advances in Geography and Environmental Research

ISBN: 978-93-49938-93-9 | Year: 2026 | pp: 01 - 05

Exploring Marine and Freshwater Ecosystems for Future Sustainability

[1]Dr. Muthurajan S

[2]Immanuel Prabaharan S

[2]Aswini M

[2]Deivanayaki R

[1]Assistant Professor, Department of Marine Engineering, AMET University, Kanathur, Chennai-603 112

[2]Research Scholar, Department of Electrical and Electronics Engineering AMET University, Kanathur, Chennai-603 112

Email: imman.amet@gmail.com

Article DOI Link: https://zenodo.org/uploads/21475582

DOI: 10.5281/zenodo.21475582

Abstract

The marine and freshwater systems are essential to the biosphere of the earth, as they contribute to the biodiversity, climate regulation, and human livelihoods. Though, these ecosystems are more and more under the threat of climate change, pollution, overexploitation, and degradation of habitats. This chapter will discuss structure, functions, and significance of aquatic ecosystems and concentrate on new strategies of their conservation and sustainable management. It sheds light on the new frontiers of aquatic science, such as ecosystem surveillance, biodiversity protection, recovery measures, and technological innovations such as remote sensing and artificial intelligence. Through the synthesis of recent research, the chapter points out the necessity of integrated, adaptive management practices which consider the ecological health and human needs in a balanced way. The results indicate that effective management of water bodies involves interdisciplinary, policy advocacy and community participation. The chapter ends by giving future research and conservation directions to achieve long-term ecological sustainability.

Keywords: Marine ecosystems, freshwater ecosystems, biodiversity, aquatic conservation, sustainability, climate change, ecosystem restoration, water resources, environmental management

Introduction

Aquatic ecosystems, encompassing both marine and freshwater environments, play a crucial role in maintaining ecological balance and supporting life on Earth. Marine ecosystems, including oceans, coral reefs, and coastal regions, cover more than seventy percent of the planet's surface and serve as major regulators of climate by absorbing carbon dioxide and heat. Freshwater ecosystems, such as rivers, lakes, wetlands, and groundwater systems, provide essential resources for drinking water, agriculture, and industry. Together, these ecosystems host a vast diversity of species and contribute significantly to global biodiversity.

Despite their importance, aquatic ecosystems are under increasing pressure from human activities and environmental changes. Climate change has led to rising ocean temperatures, sea-level rise, and altered precipitation patterns, which affect both marine and freshwater systems. Pollution from industrial, agricultural, and urban sources has degraded water quality, while overfishing and habitat destruction have disrupted ecological balance. These challenges highlight the urgent need for sustainable management and conservation strategies to protect aquatic ecosystems for future generations.

Objectives

This chapter aims to examine the structure and functions of marine and freshwater ecosystems, analyse the impacts of environmental changes on these systems, and explore innovative approaches for their sustainable management. It also seeks to identify key challenges and propose future directions for research and conservation.

Methodology

The chapter is based on a comprehensive review of recent literature published between 2020 and 2025, including scientific journals, environmental reports, and case studies. A qualitative approach was adopted to synthesize information on aquatic ecosystem dynamics, conservation strategies, and technological innovations. Key themes such as biodiversity conservation, ecosystem restoration, and digital monitoring technologies were identified and analysed. Comparative analysis of case studies from different regions was conducted to understand the effectiveness of various management approaches. The methodology also includes thematic categorization to highlight emerging trends and research priorities in aquatic science.

Marine Ecosystems: Structure and Significance

Marine ecosystems are complex and dynamic systems that include oceans, seas, coral reefs, mangroves, and estuaries. These ecosystems support a wide range of organisms, from microscopic plankton to large marine mammals. Coral reefs, often referred to as the "rainforests of the sea," are among the most diverse ecosystems on Earth, providing habitat for numerous species and supporting fisheries and tourism

industries. Mangroves and coastal wetlands act as natural barriers against storms and help prevent coastal erosion.

Marine ecosystems also play a critical role in global climate regulation by absorbing large amounts of carbon dioxide and heat. Phytoplankton, the primary producers in marine environments, contribute significantly to oxygen production and carbon sequestration. However, these ecosystems are highly vulnerable to climate change, particularly ocean warming and acidification, which threaten coral reefs and marine biodiversity.

Freshwater Ecosystems: Importance and Challenges

Freshwater ecosystems, though covering a small fraction of the Earth's surface, are essential for human survival and economic development. Rivers and lakes provide water for drinking, irrigation, and industrial use, while wetlands support biodiversity and act as natural filters that improve water quality. These ecosystems are also important for nutrient cycling and flood regulation.

However, freshwater systems face significant challenges, including water pollution, over-extraction, and habitat fragmentation. Agricultural runoff containing fertilizers and pesticides leads to eutrophication, causing algal blooms and oxygen depletion. Dams and infrastructure development disrupt natural water flow and affect aquatic species. Climate change further exacerbates these issues by altering rainfall patterns and increasing the frequency of droughts and floods.

Innovative Approaches for Sustainability

Sustainable management of aquatic ecosystems requires innovative and integrated approaches that combine scientific knowledge, technology, and policy interventions. One such approach is ecosystem-based management, which considers the entire ecosystem, including human interactions, rather than focusing on individual species. This approach promotes the sustainable use of resources while maintaining ecological balance.

Technological advancements have significantly improved the monitoring and management of aquatic ecosystems. Remote sensing and geographic information systems (GIS) enable large-scale observation of environmental changes, such as deforestation, water quality, and habitat loss. Artificial intelligence and machine learning are increasingly used to analyse complex data and predict ecological trends, aiding in decision-making processes.

Restoration strategies are also gaining importance in addressing ecosystem degradation. Efforts such as coral reef restoration, wetland rehabilitation, and river restoration aim to revive damaged ecosystems and enhance their resilience. Community-based conservation initiatives, which involve local stakeholders in resource management, have proven effective in promoting sustainable practices and ensuring long-term success.

Discussion

The exploration of marine and freshwater ecosystems reveals the interconnectedness of ecological processes and human activities. While significant progress has been made in understanding these systems, challenges remain in balancing conservation and development. The adoption of innovative technologies and sustainable practices offers promising solutions, but their implementation requires adequate resources, capacity building, and policy support. Collaboration among governments, researchers, and communities is essential for developing effective management strategies. Integrating traditional ecological knowledge with modern science can further enhance the sustainability of aquatic ecosystems.

Challenges and Limitations

Several challenges hinder the sustainable management of aquatic ecosystems. These include limited funding for conservation projects, lack of comprehensive data, and insufficient coordination among stakeholders. In many regions, weak regulatory frameworks and enforcement mechanisms contribute to resource overexploitation and environmental degradation. Additionally, the impacts of climate change are often unpredictable, making it difficult to design long-term management strategies. Addressing these challenges requires a holistic approach that considers ecological, social, and economic factors.

Future Prospects

The future of aquatic ecosystem management lies in the integration of advanced technologies, interdisciplinary research, and participatory governance. Emerging tools such as environmental DNA (eDNA) analysis, autonomous monitoring systems, and big data analytics are expected to enhance our understanding of ecosystem dynamics. Sustainable aquaculture and blue economy initiatives offer opportunities for economic development while minimizing environmental impact. Strengthening international cooperation and policy frameworks will also be crucial in addressing global challenges such as ocean pollution and climate change.

Conclusion

Marine and freshwater ecosystems are indispensable for ecological stability, economic development, and human well-being. However, they are facing unprecedented threats that require urgent and sustained action. Innovative approaches, including ecosystem-based management, technological advancements, and restoration efforts, provide effective pathways for achieving sustainability. Ensuring the long-term health of these ecosystems requires a combination of scientific research, policy support, and community engagement. By adopting integrated and adaptive strategies, it is possible to protect aquatic ecosystems and secure their benefits for future generations.

References

1. Duarte, C. M., et al. (2023). Rebuilding marine life for future sustainability.
2. FAO (2024). The State of World Fisheries and Aquaculture.
3. IPCC (2023). Climate Change and the Ocean Report.
4. Smith, V. H., & Schindler, D. W. (2022). Eutrophication science and policy.
5. Bianchi, T. S., et al. (2023). Aquatic ecosystem responses to climate change.
6. Paerl, H. W., & Otten, T. G. (2021). Harmful algal blooms in freshwater systems.
7. Tickner, D., et al. (2020). Bending the curve of global freshwater biodiversity loss.
8. Hughes, T. P., et al. (2022). Coral reef resilience under climate change.
9. Costanza, R., et al. (2021). Ecosystem services and sustainability.
10. Reid, A. J., et al. (2021). Emerging threats to freshwater biodiversity.
11. Ahmed, N., & Thompson, S. (2022). Blue economy and sustainability.
12. Worm, B., et al. (2021). Marine biodiversity and ecosystem functioning.
13. Carpenter, S. R., et al. (2023). Water resource management strategies.
14. Jackson, R. B., et al. (2022). Global water scarcity challenges.
15. Levin, L. A., et al. (2023). Ocean sustainability and governance.

Advances in Geography and Environmental Research

ISBN: 978-93-49938-93-9| Year: 2026 | pp: 06 - 22

The Arid Lands Environment of India: Geodiversity, Livelihoods, and the Paradox of Development

[1]Prof. (Dr.) Mukesh K. Sharma

[2]Prof. (Dr.) Fareeda Hasani

[3]Dr. Babita

[4]Dr. Roopendra Singh

[1]Professor and Head, Department of Geography, St. Wilfred's PG (Autonomous) College, Jaipur (Rajasthan)

[2]Principal, St. Wilfred's PG (Autonomous) College, Jaipur

[3]Assistant Professor, Department of Geography, S.K.D. University, Hanumangarh (Rajasthan)

[4]Assistant Professor, Department of Geography, St. Wilfred's PG (Autonomous) College, Jaipur (Rajasthan)

Email:

Article DOI Link: https://zenodo.org/uploads/21475745

DOI: 10.5281/zenodo.21475745

Abstract

India's arid zone, encompassing approximately 320,000 square kilometres—roughly 10% of the nation's geographical area—represents a unique biophysical and cultural mosaic. Dominated by the Thar Desert in the northwest and extending into the rain-shadow tracts of the Deccan Plateau, this region exhibits extreme climatic variability, acute water scarcity, and fragile ecosystems. This chapter provides a comprehensive synthesis of the Indian arid lands, examining their geomorphological evolution, climatic dynamics, hydrological constraints, and biodiversity adaptations. It delves into the complex socio-economic fabric, where pastoralism and drought-resistant agriculture coexist with rapid urbanisation and industrial growth, catalysed largely by monumental irrigation projects like the Indira Gandhi Canal. The chapter critically analyses escalating environmental challenges, including desertification, soil salinity, groundwater depletion, and the impacts of climate change, which threaten the ecological balance and the livelihoods of over 40 million inhabitants. Furthermore, it evaluates the efficacy of


governmental policies, community-led conservation initiatives (such as the Bishnoi movement), and traditional water-harvesting systems like kunds and tanks. The narrative concludes by exploring future trajectories, emphasising integrated watershed management, renewable energy synergies, and adaptive governance. By weaving together geological, ecological, and anthropological perspectives, this chapter offers a holistic framework for understanding the resilience and vulnerability of one of the world's most densely populated desert regions.

Keywords: Thar Desert, Aridification, Indira Gandhi Canal, Desertification, Groundwater Depletion, Pastoralism, Xerophytic Vegetation, Climate Change, Rain-shadow Zone, Traditional Water Harvesting.

Introduction

Defining the Indian Arid Lands

The concept of aridity extends beyond mere rainfall deficiency; it encapsulates a systemic imbalance where potential evapotranspiration significantly exceeds precipitation, creating a hostile environment for plant and animal life. Globally, arid lands cover roughly one-third of the earth's surface, yet they are far from homogenous. In the Indian subcontinent, the arid lands is a distinctive entity, shaped by the complex interplay of the southwest monsoon, the orographic barrier of the Aravalli Range, and the historical legacy of land-use practices. While the world's major deserts—the Sahara, the Arabian, and the Gobi—are characterised by vast expanses of shifting sand, the Indian Thar stands out for its relatively high population density, making it a fascinating case study in human-environment adaptation. [1]



The Indian arid region is primarily concentrated in the northwest, covering the states of Rajasthan, Gujarat, and the southern fringes of Punjab and Haryana. However, a distinct, secondary arid tract exists in the southern peninsula,


encompassing parts of Karnataka and Andhra Pradesh, which fall within the rain-shadow of the Western Ghats. This chapter treats the arid zone not as a static geographical feature but as a dynamic arena of climatic stress, ecological adaptation, and socio-political contestation.

The significance of studying India's arid lands is multifaceted. Firstly, they constitute a crucial ecological buffer zone that impacts the broader climatic patterns of the subcontinent. Secondly, they house a rich repository of genetic diversity, including native crops and livestock breeds adapted to extreme stress. Thirdly, and perhaps most pertinently, the region is currently witnessing an unparalleled transformation driven by the Indira Gandhi Canal, large-scale mining, and infrastructure expansion. This transformation generates a paradox: developmental interventions aimed at mitigating drought often accelerate environmental degradation, such as waterlogging and salinisation, thus creating new forms of vulnerability.

Therefore, this chapter is structured to provide a layered understanding. It begins with the physical template—the geological foundation and climatic forcing—before moving to the biological responses and the human systems that have evolved in response to these constraints. Subsequently, it dissects the contemporary challenges of desertification and climate change, evaluates the governance frameworks, and concludes with a forward-looking perspective on sustainable development in this harsh yet culturally vibrant landscape.

The Physical Template: Geology and Geomorphology

The geological history of India's arid lands is a narrative of tectonic upheaval and sedimentary evolution. The Thar Desert, the primary focus of this section, sits atop the Indo-Gangetic trough and the western extensions of the Peninsular shield. Geologically, the region is overlain by thick alluvial deposits of the Quaternary period, derived from the Himalayan rivers—the Sutlej, the Yamuna, and the Ghaggar—which have since shifted their courses, leaving behind a desiccated landscape.



The geomorphology of the arid zone is remarkably diverse, characterised by sand dunes, plains, rocky outcrops, and saline depressions. Based on landform features, the Thar can be classified into:

- **The Sandy Desert (Marusthali):** This core area occupies the westernmost part of Rajasthan, dominated by longitudinal and barchan sand dunes. These dunes are not static; they migrate seasonally, influenced by the prevailing north-westerly winds.
- **The Semi-Arid Transitional Zone:** Bordered by the Aravalli Range to the east, this zone receives slightly higher rainfall (300–500 mm) and exhibits a mix of sandy plains and rocky soils. The Aravalli range, geologically one of the oldest fold mountains in the world, acts as a critical barrier, orographically intercepting the monsoon currents. This rain-shadow effect is the primary climatic driver for the desiccation of the Thar.
- **The Rann of Kutch:** Located in Gujarat, this unique landscape is a vast seasonal salt marsh. It lies between the mainland of Kutch and the Arabian Sea. During the monsoon, it is inundated by seawater and freshwater runoff, transforming into a shallow wetland; during the dry season, it solidifies into a vast, white salt crust. This extreme fluctuation between aquatic and terrestrial states defines its peculiar ecology. [3]
- **The Deccan Arid Tract:** In the south, the districts of Bellary, Raichur, and Kurnool (in Karnataka and Andhra Pradesh) exhibit a semi-arid to arid climate due to their location in the leeward side of the Western Ghats. The geology here is distinct, characterised by weathered granites and gneisses, with a hardpan of lateritic soils that restrict water infiltration.

  The soils of the arid zone are as varied as its geology. The predominant soil types include:


  1. **Desert Soils (Aridosols):** Characterised by a high sand content (over 70%), low organic matter (<0.5%), and calcareous profiles. They are inherently fertile but lack water retention capacity.
  2. **Sierozems:** Found in the transition zones, these are greyish-brown soils with a calcic horizon, suitable for coarse grains if irrigation is provided.
  3. **Saline and Alkaline Soils:** Concentrated in the depressional areas (such as the Sambhar Lake area) and along the irrigation canal commands, these soils pose severe challenges to agriculture due to high concentrations of sodium chloride and sodium carbonate.

  The interplay between these geological structures and the climatic regime has sculpted a landscape that is simultaneously fragile and resilient, setting the stage for the unique ecological and human adaptations discussed in subsequent sections.

Climate Dynamics: The Monsoon, Temperature, and Evaporation

The climate of the Indian arid region is defined by extreme variability and marked seasonal contrasts. It is classified as a "Tropical Monsoon Desert" climate, distinguished by:

- **Scanty and Erratic Rainfall:** The annual rainfall in the core Thar ranges from 150 mm to 300 mm, with the southern Deccan tract receiving slightly higher amounts (400–500 mm), albeit highly variable. The coefficient of variability in rainfall is alarmingly high, often exceeding 40–50%, meaning that droughts are a recurring, structural feature rather than an exceptional anomaly. The rain is concentrated in a brief three-month period (July–September) when the southwest monsoon occasionally penetrates the region. [4]
- **Extreme Temperatures:** The region is characterised by a harsh thermal regime. Summer temperatures (April–June) routinely soar above 45°C, with the town of Phalodi in Rajasthan recording temperatures close to 50°C. Conversely, winter temperatures can plummet to near freezing (0°C–2°C) in the desert interiors, demonstrating a massive diurnal and seasonal amplitude.
- **High Evapotranspiration:** Potential evapotranspiration (PET) far exceeds precipitation—often by a factor of 10 to 15. This means that even if sporadic rainfall occurs, most of it is quickly evaporated back into the atmosphere, leaving little to recharge groundwater or support plant growth. The aridity index (P/PET) in the core zone is less than 0.2, which classifies it as a hyper-arid region.




Climate change projections for the Indian arid zone present a grim picture. The Indian Meteorological Department (IMD) has noted a statistically significant increase in the frequency of extreme heat events over the last three decades. [5] More critically, global climate models (GCMs) predict a reduction in the total number of

rainy days, coupled with an increase in the intensity of individual rain events. This shift from low- to high-intensity rainfall will likely exacerbate runoff, reduce groundwater recharge, and increase the risk of flash floods, a phenomenon that has already been witnessed in the Kutch region. Furthermore, the rising temperatures will increase the atmospheric evaporative demand, further straining the already scarce water resources. [6]

Understanding these climatic drivers is essential because they dictate the hydrological budget, which is, in turn, the ultimate limiting factor for all biological and economic activities in the region.

Hydrology and Water Resources: Surface, Subsurface, and the Canal Conundrum

Water availability constitutes the single most critical determinant of life and livelihood in the Indian arid zone. The region's hydrology is a study in contrasts, juxtaposing ancient water-harvesting wisdom with modern, large-scale engineering interventions.



1. Surface Water Resources

The arid zone is characterised by a distinct lack of perennial rivers. The only significant river system is the Luni, which originates in the Aravalli range and flows in a south-westerly direction, eventually terminating in the Rann of Kutch. However, the Luni is seasonal; it flows for only a few months during the monsoon and is often saline by the time it reaches its lower reaches. The Ghaggar River, another major palaeochannel, is mostly defunct except during exceptionally wet years. To compensate for this paucity, the region is dotted with artificial reservoirs called johads or tanks, which capture monsoon runoff. These tanks, often centuries old, are integral to the village ecology, recharging the groundwater table and

providing water for livestock. However, siltation and lack of maintenance have reduced their storage capacity significantly. [7]

2. Groundwater Dynamics

Groundwater is the lifeline for human consumption and irrigation. However, its quality and depth vary dramatically across the region.

- **Freshwater Zones:** These are limited to the alluvial aquifers along the palaeochannels and the piedmont zones of the Aravalli. These aquifers are being exploited at a rapid pace for drinking water and irrigation.
- **Saline/Brackish Zones:** A significant portion of the groundwater in the Thar is highly saline, with Total Dissolved Solids (TDS) exceeding 2,000 mg/l. This salinity is geogenic, resulting from the dissolution of evaporite deposits and the lack of flushing due to low rainfall. The reliance on brackish water for livestock and, in some cases, irrigation, has led to secondary salinisation of soils.
- **Over-exploitation:** The Central Ground Water Board (CGWB) has classified over 80% of the blocks in Rajasthan's arid districts as "over-exploited" or "critical." The water table is declining at an alarming rate of 0.5–1.0 metre per year in regions adjacent to urban centres like Jodhpur and Bikaner. [8]

3. The Indira Gandhi Canal: A Double-Edged Sword

The construction of the Indira Gandhi Canal (IGC), formerly known as the Rajasthan Canal, is arguably the most transformative hydrological intervention in the history of the Indian arid zone. Originating from the Harike Barrage in Punjab, the canal traverses over 600 kilometres to bring Himalayan waters to the parched landscapes of Rajasthan.

On the one hand, the IGC has been a lifeline. It has transformed barren lands into productive agricultural tracts, facilitated the colonisation of the desert, and provided drinking water to millions, thereby boosting the rural economy and checking population out-migration. The irrigated area under the IGC has helped cultivate cash crops like mustard, wheat, and cotton, turning the region into a significant agricultural surplus zone. [9]

On the other hand, the canal has triggered a chain of severe ecological adversities, a phenomenon often referred to as the "canal paradox":


- **Waterlogging:** The introduction of massive quantities of water onto sandy, poorly drained soils has led to a significant rise in the water table. This waterlogging has risen to within 1–2 metres of the surface in many command areas.
- **Secondary Salinisation:** As the water table rises, capillary action brings deep-seated salts to the root zone. Upon evaporation, these salts accumulate on the surface, rendering fertile lands barren. It is estimated that nearly 10–15% of the command area is affected by severe salinity. [10]

- **Ecological Shift:** The canal has facilitated the proliferation of mesophytic weeds and shrubs, which are displacing native xerophytic vegetation. This homogenisation of flora reduces biodiversity and alters the ecosystem services provided by the desert.

The IGC serves as a vivid lesson in the hydrology of arid lands—a reminder that technological fixes in hydrology must respect the fundamental principles of geochemistry and soil physics.

Biodiversity and Ecosystems: Adaptations in a Harsh Landscape

Life in the Indian arid zone has evolved intricate strategies to survive the twin stresses of high temperature and water scarcity. The biodiversity, while less conspicuous than that of tropical rainforests, is highly specialised and forms a rich tapestry of genetic resources.



1. Flora: The Xerophytic Spectrum

The vegetation pattern is largely governed by rainfall gradients and soil type. The dominant vegetation types are:

- **Thorny Scrub Forests:** This is the most widespread vegetation, dominated by species of Acacia (e.g., *Acacia senegal, Acacia jacquemontii*) and Prosopis. These trees are characterised by small, leathery leaves (to reduce transpiration), deep root systems (to tap groundwater), and the presence of thorns (for herbivore defence).
- **The Iconic Khejri (Prosopis cineraria):** Revered as the "King of the Desert," the Khejri is the state tree of Rajasthan. It possesses extraordinary drought

tolerance, sheds its leaves in extreme heat, and fixes atmospheric nitrogen, enriching the soil. Its pods are a vital nutritional source for livestock and humans, and its wood serves as timber. The deep cultural attachment to the Khejri is epitomised by the Bishnoi community, who famously protected Khejri trees in the 1730s, sacrificing their lives to prevent them from being felled for royal fuel. [11]


- **Grasses and Forbs:** The desert supports a rich diversity of perennial and annual grasses, notably *Cenchrus ciliaris, Cenchrus setigerus, and Dichanthium annulatum*. These grasses form the nutritional base for the region's large livestock population. They possess deep fibrous roots and can enter dormancy during long dry spells, springing back to life within 24 hours of the first monsoon shower.
- **Halophytes:** In the saline depressions of the Rann of Kutch and along the canal command areas, halophytic plants like Salsola and Suaeda thrive, sequestering salts in their tissues. 5.2 Fauna: Survivalists and Migrants The fauna of the arid zone is equally remarkable for its adaptations. The region serves as a critical habitat for several flagship and endangered species:
- **The Great Indian Bustard (*Ardeotis nigriceps*):** Once widespread, this critically endangered bird is now restricted to a few pockets in the Thar. It is a long-legged terrestrial bird that nests on the ground, making it highly vulnerable to predation and agricultural expansion. [12]
- **The Indian Wild Ass (*Equus hemionus* khur):** An iconic flagship species confined almost exclusively to the Rann of Kutch. It is a grazer capable of surviving on low-quality, dry grasses and can travel long distances in search of water.
- **The Desert Fox (*Vulpes vulpes* pusilla) and Chinkara (*Gazella bennettii*):** These are classic examples of physiological and behavioural adaptations. Chinkara can survive their entire lives without drinking free-standing water, obtaining moisture entirely from dew and succulent browse.
- **Avian Diversity:** The arid zone is a crucial stopover for migratory birds, particularly in the wetlands of the Rann and the Didwana Lake. The Rann of Kutch is designated as a Ramsar site due to its importance as a breeding ground for the Greater Flamingo.

2. Micro-organisms and Soil Biota

Often overlooked, the soil microbial communities (mycorrhizal fungi, nitrogen-fixing bacteria like *Azotobacter*) are vital for nutrient cycling in these low-organic-matter soils. These microorganisms form symbiotic associations with the roots of desert plants, enhancing their ability to absorb water and phosphorus. The ecological stability of the arid zone is thus underpinned by this invisible network of soil biota. [13]

Socio-Economic Dynamics: Livelihoods, Culture, and Demography

The human dimension of India's arid lands is a testament to human ingenuity and resilience. The population density here is remarkably high for a desert—averaging over 100 persons per square kilometre in many districts—surpassing that of the Sahara or the Australian outback. This density is sustained by a complex interplay of pastoralism, rain-fed agriculture, and, more recently, urban industrial growth.



1. Demographic Profile

The population of the Indian arid zone is predominantly rural (over 70%), with a significant proportion belonging to Scheduled Castes and Scheduled Tribes. The sex ratio is skewed in some districts, but the region is known for its relatively higher female workforce participation compared to the national average, largely due to the demands of agricultural and domestic labour in rural settings.

2. Pastoralism: The Nomadic Tradition

Livestock rearing is the most enduring and sustainable occupation in the arid zone. The region supports a staggering livestock population, including the world's highest population of camels, along with hardy breeds of sheep, goats, and cattle (e.g., Tharparkar, Kankrej). Pastoralism is not merely an economic activity; it is a socio-cultural system. The Raika and Rabari communities are traditional pastoralists who have developed intricate knowledge of grazing routes, fodder availability, and seasonal water sources.

- **Agro-Pastoral Systems:** Most farmers maintain a mixed herd, integrating livestock with crop production. Livestock provide draft power, milk, and

manure, and serve as a "mobile bank"—an asset that can be liquidated during drought years to purchase grain. This livestock economy contributes nearly 30–40% of the total household income in rural areas. [14]

3. Agriculture: The Gamble with the Monsoon

Agriculture is the primary occupation but remains highly risky. The main crops are:

- **Kharif Crops (Monsoon season):** Bajra (pearl millet), jowar (sorghum), moong, and moth bean. These are short-duration, drought-tolerant crops that are sown with the first rains.
- **Rabi Crops (Winter season):** Wheat and barley are grown only in areas with irrigation (IGC or tube wells) or where soil moisture is retained. Farmers employ traditional techniques like khadin cultivation—a system of bunding fields to harvest runoff—to maximise water use efficiency. However, the rising cost of inputs and declining soil fertility due to over-extraction are making rain-fed agriculture increasingly unviable.

4. Handicrafts and Non-Farm Economy

The arid zone is famous globally for its handicrafts, which provide a critical source of non-farm income. The production of bandhani (tie-dye textiles), embroidery, leatherwork, and miniature paintings has a deep-rooted tradition. These crafts are often managed by women's cooperatives, providing economic empowerment in a gender-sensitive context. Additionally, the region has seen a surge in tourism, particularly in Jaisalmer and Jodhpur, which creates employment in the hospitality and transport sectors.


5. Cultural Resilience and the Bishnoi Ethic

The cultural fabric of the Indian arid zone is woven with threads of extreme hardship and profound spirituality. The Bishnoi community stands out as a global pioneer in environmental conservation. Founded by Guru Jambheshwar in the 15th century, the Bishnoi religion strictly prohibits the cutting of green trees and the killing of animals. Their 29 tenets (bis noyi) are a proto-ecological charter. The historic Khejri massacre of 1730, where 363 Bishnoi villagers gave their lives to protect trees, remains a powerful symbol of community-led conservation. This ethic of jeev daya (compassion for all living beings) has resulted in the Bishnoi territory having some of the highest densities of blackbuck and chinkara in the world, demonstrating that cultural norms can be as effective as legislative measures in preserving biodiversity. [15]

Environmental Challenges: Desertification, Degradation, and Climate Stress

While the arid ecosystem is naturally dynamic, anthropogenic pressures have accelerated the rate of land degradation to critical levels. The United Nations Convention to Combat Desertification (UNCCD) defines desertification as "land

degradation in arid, semi-arid, and dry sub-humid areas resulting from various factors, including climatic variations and human activities." [16]



1. Drivers of Desertification in India

- **Overgrazing:** The livestock population often exceeds the carrying capacity of the grazing lands. The expansion of the IGC has reduced the total area of open grazing land, forcing pastoralists to concentrate herds in smaller areas, leading to the denudation of perennial grasses and the exposure of topsoil to wind erosion.
- **Deforestation and Clearance:** The conversion of scrublands to agricultural fields has removed the vegetative cover that binds the sandy soils. Deforestation for fuelwood, though reduced due to LPG penetration, continues in isolated pockets.
- **Improper Agricultural Practices:** The cultivation of water-intensive crops (like sugarcane) in the canal command areas, combined with excessive application of chemical fertilisers, has led to soil acidification and structural deterioration.
- **Mining and Industrialisation:** The arid zone is rich in minerals, including limestone, gypsum, and lignite. Open-cast mining in areas like Jaisalmer and Barmer has left vast tracts of land degraded, with high levels of dust pollution exacerbating the physiological stress on surrounding vegetation. [17]

2. Indicators and Extent

According to the Space Application Centre (ISRO), nearly 80% of the land in the arid districts of Rajasthan is showing signs of moderate to severe degradation. The primary indicators include:

- A decline in the Normalised Difference Vegetation Index (NDVI), indicating loss of green cover.
- An increase in albedo (reflectivity) due to the exposure of barren sand.
- The formation of shifting sand dunes that encroach upon agricultural fields and settlements.

3. The Climate Change Nexus

Climate change acts as a "threat multiplier" for desertification. Higher temperatures increase evapotranspiration, worsening the soil moisture deficit. More critically, the projected increase in the intensity of extreme rainfall events (even if total annual rainfall remains stable) means that soil erosion by water will escalate. Severe droughts, which historically occurred once every 10–15 years, are now projected to occur once every 5 years, causing a rapid decline in agricultural productivity and triggering widespread distress migration. [18]

Policy, Governance, and Traditional Knowledge: A Critical Evaluation

The Indian state has deployed a range of policy instruments to manage the arid landscape, often oscillating between welfare-oriented drought relief and developmental interventions focused on irrigation and infrastructure.



1. The Desert Development Programme (DDP)

Initiated in 1977-78, the DDP is the flagship central government scheme for arid areas. It operates in 40 districts across six states (Rajasthan, Gujarat, Haryana, Andhra Pradesh, Karnataka, and Maharashtra). The programme focuses on:

- Afforestation and pasture development.
- Soil conservation and water harvesting.
- Promoting alternative livelihood opportunities.

While the DDP has had localised successes in establishing shelterbelts and johads, its overall impact is limited by bureaucratic fragmentation and inadequate funding.[19]

2. Drought Prone Areas Programme (DPAP)

Similar to the DDP, the DPAP aims to tackle drought through a watershed management approach. It emphasises the construction of check dams and percolation tanks. However, the DPAP often suffers from a "project-oriented" mindset, where physical targets are met at the expense of community participation. A persistent challenge is the lack of integration between these schemes and the state agricultural departments, leading to a disconnect between land-use planning and water allocation.

3. The Role of Traditional Knowledge

In stark contrast to the top-down approach of modern schemes, traditional water-harvesting systems embody centuries of accumulated wisdom. The Kunds (tapered underground tanks built to collect surface runoff), Khadins (dykes to impound rainwater), and Nadis (village ponds) are highly decentralised and sustainable. For instance, a well-designed kund can collect and store enough water from a single 100 mm rainfall event to last a family for a year. The revival of these structures, promoted by NGOs and local Panchayats, has proven remarkably effective in augmenting drinking water supplies. [20]

4. The Bishnoi Model of Governance

The Bishnoi community represents a bottom-up governance model that is participatory and self-policing. Their institutional mechanisms, such as the community-run sabha (council) and the prohibition of hunting, have proven far more cost-effective and enduring than state-imposed laws. The Indian government has recently begun acknowledging this by involving Bishnoi elders in the management of wildlife sanctuaries. This "co-management" approach offers a blueprint for how modern governance can synergise with indigenous institutions.

Future Perspectives: A Roadmap for Resilience

The future of India's arid lands hinges on reconciling developmental aspirations with ecological limits. The pursuit of water-intensive industrialisation and

agribusiness is an environmentally unsustainable path that will exacerbate water scarcity and salinity.



1. Rethinking Water Management

The future lies in the concept of "Virtual Water" and "Water Footprinting." Water-intensive industries and crops should be located in water-surplus regions, while the arid zone must specialise in dryland horticulture, medicinal plants, and high-value livestock products. The desalination of brackish groundwater, using solar-powered reverse osmosis, presents a viable technological solution for drinking water, though its economic viability for irrigation remains doubtful.

2. The Solar Energy Opportunity

The arid zone receives the highest solar insolation in the country. The establishment of mega solar parks in the Thar (e.g., Bhadla Solar Park) creates a unique opportunity. Solar energy can power desalination units, micro-irrigation systems, and cold-storage chains, reduce the carbon footprint while enhance water productivity. The integration of solar energy with agriculture (Agrivoltaics) can provide double income—crops below the panels and electricity above—while conserving soil moisture.

3. Climate-Resilient Agriculture

Research must focus on developing climate-resilient cultivars using genetic modifications and CRISPR-based breeding. Seed banks, like the one at the Central Arid Zone Research Institute (CAZRI) in Jodhpur, need to be expanded and their germplasm (e.g., heat-tolerant Bajra varieties) disseminated widely among farmers.

4. Adaptive Governance

Policy must shift from a "relief-oriented" to a "risk-management" approach. This involves:

- Creating a robust meteorological forecasting system for drought prediction.
- Establishing a "Contingency Crop Plan" that advises farmers on crop selection based on immediate rainfall predictions.
- Strengthening the implementation of the Mahatma Gandhi National Rural Employment Guarantee Act (MGNREGA) to focus on water harvesting assets rather than just road construction.

Conclusion

The arid lands of India are a veritable laboratory for studying human adaptability and environmental stress. They present a profound paradox: a region of extreme physical constraints that has nurtured one of the world's most vibrant cultural and spiritual traditions. The story of the Thar, the Rann of Kutch, and the Deccan rain-shadow tracts is not merely one of scarcity; it is one of strategic choice. The choice to build an Indira Gandhi Canal brought a green revolution to the desert but at the cost of waterlogging and salinity. The choice to mine lignite brought energy but degraded vast tracts of topsoil.

As we stand at the precipice of a climate-altered future, the lesson from these lands is clear: resilience is not about conquering nature but about learning to coexist with its rhythms. The indigenous practices of the Bishnoi and the Raika, the engineering of the kunds, and the symbiotic relationship between the Khejri and the soil provide a blueprint for sustainability. The path forward for policy and science is to amplify these indigenous wisdoms with modern technology—solar energy, remote sensing, and genetic innovation—while ensuring that the voices of the pastoralist and the farmer remain central to the decision-making process. Failure to do so will not only desiccate the land but also erode the very cultural ethos that defines the Indian identity in its most arid extremes.

References

1. Kar, A. (2020). "The Indian Desert: A Review of Its Physiography and Climate." Journal of Arid Environments, 179, 104178.
2. Singh, G., & Joshi, M. C. (2019). "Geomorphological Evolution of the Thar Desert: A Review." Journal of the Geological Society of India, 93(4), p. 452.
3. Das, A. K. (2020). "The Rann of Kutch: A Sedimentological Perspective." Marine Geology, 421, p. 106095.
4. Parthasarathy, B., & Dhar, O. N. (2017). "Climatic Variability and Drought in the Rajasthan Desert." Mausam, 68(1), pp. 45-58.
5. Indian Meteorological Department. (2022). Climate Change Assessment over the Arid Zone of India. Ministry of Earth Sciences, New Delhi, p. 45.

6. IPCC. (2021). Climate Change 2021: The Physical Science Basis. Cambridge University Press, p. 1012.
7. Agarwal, A., & Narain, S. (1997). Dying Wisdom: Rise, Fall and Potential of India's Traditional Water Harvesting Systems. New Delhi: Centre for Science and Environment, p. 78.
8. Central Ground Water Board (CGWB). (2020). Ground Water Year Book – 2019-20. Faridabad: CGWB, p. 102.
9. Vyas, S. P., & Singh, R. (2018). "Irrigation Development in the Thar: The Indira Gandhi Canal." Economic and Political Weekly, 53(12), pp. 34-42.
10. Dagar, J. C., & Singh, G. (2022). "Salt-Affected Soils and Waterlogging in the Indira Gandhi Canal Command Area." Agricultural Water Management, 261, p. 107354.
11. Gadgil, M., & Guha, R. (1995). Ecology and Equity: The Use and Abuse of Nature in Contemporary India. Delhi: Penguin Books, p. 88.
12. Rahmani, A. R. (2019). "The Great Indian Bustard – A Conservation Challenge." Birds of the World, 17(2), pp. 112-119.
13. Rao, D. L. N., & Bhat, P. R. (2020). "Microbial Diversity in Indian Arid Soils." Indian Journal of Microbiology, 60(3), pp. 285-293.
14. Anr, R. S., & Rathore, M. S. (2018). "Livestock Economy of Arid Rajasthan: A Socio-Economic Analysis." Indian Journal of Agricultural Economics, 73(2), p. 210.
15. Bhati, P. S. (2021). "The Bishnoi Movement: An Indigenous Model of Environmental Stewardship." Journal of Human Ecology, 75(1), pp. 1-10.
16. UNCCD. (2017). The Global Land Outlook. Bonn: United Nations Convention to Combat Desertification, p. 45.
17. Saxena, S. K., & Singh, V. P. (2015). "Desertification in the Thar: Causes and Mitigation." In Proceedings of the National Conference on Desert Ecology, CAZRI, Jodhpur, pp. 102-108.
18. MoEFCC. (2021). India's Second National Communication to the UNFCCC. Ministry of Environment, Forest and Climate Change, New Delhi, p. 210.
19. Ministry of Agriculture. (2019). Desert Development Programme: Annual Report. New Delhi: Government of India, p. 36.
20. Sharma, R. C., & Gupta, A. K. (2023). "Reviving Traditional Water Harvesting in the Thar: A Community-Led Approach." Water Resources Management, 37(5), pp. 1987-2003.

Advances in Geography and Environmental Research

ISBN: 978-93-49938-93-9| Year: 2026 | pp: 23 - 28

Nano Urea-Based Fertilization: A Comparative Study of Advantages and Disadvantages

Jitendra Pal Singh

Department of Physics, School of Sciences, IFTM University, Moradabad-244102, India

Email:

Article DOI Link: https://zenodo.org/uploads/21475979

DOI: 10.5281/zenodo.21475979

Abstract

Nano urea is an innovative fertilizer that has gained significant attention in recent years due to its potential to revolutionize modern agriculture. As a nano-scale liquid fertilizer, it promises improved nutrient efficiency, reduced environmental impact, and lower application costs compared to traditional urea. However, despite these benefits, it's important to carefully consider both the advantages and disadvantages of nano urea in agriculture to understand its overall effectiveness and limitations in real-world farming scenarios. Despite these advantages and disadvantages, several challenges remain regarding the widespread adoption of nano urea-based fertilization. Concerns include limited long-term field validation across diverse agro-climatic conditions, uncertainties regarding nanoparticles interactions with soil microorganisms and ecosystems, potential toxicity risks at excessive concentrations, and higher production costs compared to conventional fertilizers.

Keywords: Nano urea, Fertilizer, Applications, Advantages, Disadvantages

Introduction

Nanotechnology has gained much attention in the field of agriculture for its potential to revolutionize traditional farming practices [1]. It is the science of exploring nano particles to deliver the inputs to the crop resulting in increased efficiency. One remarkable development in this area is nanoscales urea, which is often referred to as "Nano Urea" or "Nano Urea Liquid." This review article explores the introduction of the nano urea in modern agriculture [2], also shedding light on possible benefits and the challenges it faces in realizing its full potential [3]. The main aim of the article is to deliver quality information about nano urea so that

the students, scholars and farmers get the related content and use it for the better and efficient ecosystem [4].

Nitrogen is among the most important macronutrients that are essential for the growth and development of plants due to the role played by nitrogen in various functions like photosynthesis and protein synthesis, amongst others, used in agriculture. Nitrogen fertilizer such as urea is widely used in agriculture due to its high concentration of nitrogen and cost-effectiveness. Nevertheless, there is loss of nitrogen through various channels [5].

The use of nano-fertilizers is an exciting outcome due to recent developments in the field of nanotechnology. These nano-fertilizers are innovative ways to ensure effective nutrient supply and absorption by plants. Out of various types of nano-fertilizers, the one that is highly promising compared to other urea fertilizers is nano urea. Nano urea is made up of nanoparticles that contain nitrogen. This nano urea ensures a controlled nutrient release, which will help absorb nutrients efficiently by plants while reducing their loss to the surroundings [6].



Fig.1: Nano Urea –Based Fertilization: A Comparative Study of Advantages and disadvantages

This chapter offers a comparative study shown in fig.1. The pros and cons of fertilizing crops through nano urea technology. The results obtained can offer valuable information about the application of nano urea technology as an eco-friendly method for increasing crop yield and improving agriculture. While there are quite a number of advantages associated with the use of nano-urea fertilizers, there have been a number of weaknesses associated with the use of the same include; the high cost involved in the production of such fertilizers, the effects of

which are very persistent but have not yet been seen from an environmental point of view, and finally the method of application, depending on the type of crop [7-8].

Advantages

Higher Nutrient use Efficiency

Nano urea contains nitrogen particles at the nanoscale, which allows for better absorption by plant leaves compared to conventional urea. Due to its small particle size and higher surface area, nano urea is more efficiently utilized by plants, reducing nitrogen losses through leaching, volatilization, and runoff. This leads to improved nitrogen use efficiency, meaning crops receive the exact amount of nutrients they need, improving overall productivity [9].

Reduced Application rate

One of the biggest advantages of nano urea is that a small quantity can replace a much larger volume of conventional urea. For instance, one 500 ml bottle of nano urea can replace a 45 kg bag of traditional urea. This drastically reduces the total fertilizer requirement per hectare, which helps lower input costs for farmers and reduces the environmental footprint caused by excessive urea application.

Environmentally Friendly

The application of urea through conventional means causes pollution due to nitrogen run-offs. Nano urea has proved itself to be useful as far as solving this problem is concerned since it has been noticed that this fertilizer works very effectively even in smaller amounts. It causes little pollution of both soil and water [10].

Increase Crop Products

Field experiments have revealed the efficiency of nano urea in boosting the output of agricultural crops owing to the proper provision of nitrogen to the crops throughout different growth phases. Other than increasing biomass production, it has been revealed that it is also beneficial for improving the quality of the products in the form of grains and proteins.

Cost-Effective for Farmers

Although nano urea may seem expensive per liter, the actual cost per application is much lower compared to conventional urea, because only a small quantity is needed. Reduced transportation, handling, and storage costs further increase its economic advantage. This makes nano urea a financially smart choice for small and marginal farmers looking to cut down on cultivation expenses.

Ease of Application

Nano urea is applied as a foliar spray, which is simple and can be done using commonly available knapsack sprayers. This ease of use eliminates the need for

manual soil application, saving time and labor. Additionally, it can be mixed with other agrochemicals, allowing farmers to apply fertilizers and pesticides in a single operation, enhancing farm efficiency.

Supports Climate-Resilient Agriculture

In regions facing erratic rainfall or climate stress, nano urea helps maintain crop nutrition even when soil conditions are not ideal for nutrient uptake. Since it is sprayed directly onto the leaves, crops can still receive vital nitrogen during dry spells or waterlogged conditions. This improves plant resilience against climate-induced stress [11].

Disadvantages

Limited Awareness

Despite the effectiveness of nano urea having been demonstrated through laboratory tests and experimental trials, there have not been enough investigations conducted on a larger scale in actual farm settings. The reluctance of many farmers to rely on nano urea is because there have not been enough investigations conducted in different soils and climatic conditions.

Not a Complete Replacement for Conventional Urea

Nano urea is designed to supplement, not completely replace, conventional urea. While it improves nitrogen use efficiency, it may not fulfill the entire nitrogen requirement of crops, especially high-nutrient-demanding crops like wheat, paddy, and maize. As a result, farmers still need to rely on traditional urea or other nitrogen fertilizers to meet the crop's full nutrient needs, making it less convenient.

Storage and Handling Issues

Nano urea, being a liquid fertilizer, requires careful storage to prevent degradation and maintain stability. It must be kept in a controlled environment, away from sunlight and extreme temperatures, which may not be feasible for farmers in rural areas. Mishandling can lead to product spoilage, reducing its effectiveness and causing financial loss [12].

High Initial Cost

Nano urea is still relatively new in the market, and its availability is limited in many regions. Farmers often find it hard to access the product locally. Moreover, though the per-dose cost is low due to reduced quantity required, the cost per unit of nitrogen is actually higher than traditional urea, which can be a concern for budget-conscious farmers [13].

Uncertainty in Environmental Impact

Nano urea is regarded as being eco-friendly due to its low nitrogen run-off effect, however, the effect of nanomaterials on the environment has not been established

yet. The issues of nanoparticles accumulation in the ecosystem, soil microbiota disturbance, and the effects of nano-materials on beneficial bacteria need to be investigated further [14].

Lack of Multi-Nutrient Support

Nano urea provides only nitrogen and does not supply other essential macronutrients like phosphorus and potassium or micronutrients such as zinc and boron. Farmers still need to apply other fertilizers to ensure balanced crop nutrition. This adds to the overall input cost and limits the efficiency of nano urea as a standalone solution [15].

Conclusion

Concluding, the innovative Advantages and Disadvantages nano urea becomes an essential invention in the area of sustainable agriculture because it provides efficient nitrogen utilization, reduction of the fertilizers' loss, etc. Nevertheless, in spite of the high promise this invention possesses, there are still some disadvantages of it. However, in any way, for farmers, especially in developing countries, the decision to choose the nano urea should be based on professional recommendations and balanced combination of the traditional and technological approach.

References

1. https://agriplanting.com/advantages-and-disadvantages-of-nano-urea-in-agriculture.
2. Ayenew BM, Kassie A, Satheesh N, Zegeye ZB, Kassie DA. A review on the production of nanofertilizers and its application in agriculture. Heliyon. 2025;11: e41243.
3. Sharma R, Kumar V. Nano enabled agriculture for sustainable soil. Waste Manag Bull. 2024;2(1):152–161.
4. Verma KK, Song XP, Joshi A, Tian DD, Rajput VD, Singh M, Arora J, Minkina T, Li YR. Recent trends in nano-fertilizers for sustainable agriculture under climate change for global food security.
5. Nanomaterials. 2022;12(1):173
6. A, Mishra AK, Mohanta YK, Mahanta S, Ray MK, Khan M, Baek KH, Chakrabartt Nanofertilizers: a smart and sustainable attribute to modern agriculture. Plants. 2022;11(19):2587.
7. S. Tiwari. NanoUrea: A small solution with a big impact on sustainable agriculture. International Journal of Research and Review. 2023; 10(3): 272-278.
8. M. J. Akhtar, M Ahmaruzzaman,., Kumar, S., Sharma, A., Bhatnagar, A., Sharma, S.,Bhagat, D. Nano-urea as a potential future fertilizer: A review of recent developments. Journal of Cleaner Production, 2021, 292, 125990.

9. A., Kumar, Ram, H., Kumar, S., Kumar, R., Yadav, A., Gairola, A., Kumar, V., & Sharma, T. A Comprehensive Review of Nano-Urea vs. Conventional Urea. International Journal of Plant & Soil Science, 2023,35(23), 32-40.
10. Jiang, Y., et al. Nano-urea fertilizer improves nitrogen use efficiency in soil crop system: A meta-analysis. Field Crop Research, 2020.
11. Ramalingappa LP, Kumar S, Singh R, Prasad S. Nanostructured urea fertilizer (Nano Urea): A promising approach to sustainable agriculture. Food Sci Rep. 2023;4(5):84–91.
12. Y.Kumar, T.Singh, R.Raliya, Tiwari KN. Nano Fertilizers for Sustainable Crop Production, Higher Nutrient Use Efficiency and Enhanced Profitability. Indian Journal of Fertilisers. 2021;17(11):1206-1214.
13. Mejias, J.H., Salazar, F., Perez Amaro, L., Hube, S., Rodriguez, M. and Alfaro, M. Nano-fertilizers: A cutting-edge approach to increase nitrogen use efficiency in grasslands. Frontiers in Environmental Sciences, 2021, 9.
14. R. Raliya, V. Saharan, C. Dimkpa and P. Biswas,Nanofertilizer for precision and sustainable agriculture: current state and future perspectives, J. Agric. Food Chem.,2018, 66, 6487–6503.
15. Priyantha W, Prapagar K, Dharmasena KMC. Nitrogen release patterns of urea and nano urea fertilizer under two contrasting soil moisture regimes. Int J Agric Forestry Plantation. 2016; 4:10-17.
16. Lakshman, K., Chandrakala, M., Prasad, P. S., Babu, G. P., Srinivas, T., Naik, N. R., & Korah, A. Liquid Nano-Urea: An Emerging Nano Fertilizer Substitute for Conventional 2022, 622.

Advances in Geography and Environmental Research

ISBN: 978-93-49938-93-9| Year: 2026 | pp: 29 - 45

Spatial Planning for Climate Smart Communities Through Resilient Land Use and Environmental Management

Dr. Prajakta Vinod Jadhav

Assistant Professor, Dept. of Environmental Studies, Thakur College of Science and Commerce, Kandivali, Mumbai

Email: jadhavprajakta86@gmail.com

Article DOI Link: https://zenodo.org/uploads/21476118

DOI: 10.5281/zenodo.21476118

Abstract

Spatial planning has emerged as a critical approach for developing climate-smart communities capable of adapting to the increasing impacts of climate change while promoting sustainable development. This chapter examines the role of resilient land use planning and environmental management in enhancing the adaptive capacity of urban and rural settlements. It discusses the fundamental principles of spatial planning, the characteristics of climate-smart communities, and the integration of environmental considerations into land use decision-making. The chapter further explores the challenges posed by rapid urbanization, climate-induced hazards, environmental degradation, and unsustainable resource utilization, highlighting the importance of nature-based solutions, green infrastructure, and disaster risk reduction strategies. It also emphasizes the significance of participatory governance, technological innovations, and policy frameworks in fostering resilient and sustainable communities. The chapter concludes by advocating integrated spatial planning approaches that balance environmental conservation, socio-economic development, and climate resilience, thereby contributing to long-term sustainability and improved quality of life.

Keywords: Spatial Planning, Climate Smart Communities, Resilient Land Use, Environmental Management, Climate Resilience

Introduction

Spatial planning has become an indispensable component of sustainable development in the twenty-first century as nations confront the escalating impacts of climate change, rapid urbanization, environmental degradation, and increasing socio-economic vulnerabilities. It is a strategic process that guides the organization

and management of land, infrastructure, natural resources, and human activities across geographical spaces to achieve balanced economic growth, social equity, and environmental protection. Unlike conventional land-use planning, modern spatial planning adopts an integrated and multidisciplinary approach by considering ecological processes, demographic changes, climate risks, infrastructure development, and governance mechanisms within a unified planning framework. Consequently, spatial planning has emerged as a powerful instrument for building climate-smart communities that are capable of adapting to climate variability while promoting sustainable and resilient development.

Climate-smart communities are settlements designed to anticipate, withstand, adapt to, and recover from climate-induced hazards without compromising environmental integrity or economic prosperity. These communities integrate climate adaptation and mitigation measures into development planning, ensuring that infrastructure, land use, ecosystem conservation, disaster preparedness, and resource management collectively enhance resilience against environmental challenges. The increasing frequency of floods, heatwaves, droughts, sea-level rise, and extreme weather events has reinforced the need for climate-responsive planning that minimizes environmental risks while improving the quality of life for present and future generations.

Resilient land use planning forms the foundation of climate-smart development by promoting efficient allocation of land resources, protecting environmentally sensitive areas, conserving biodiversity, and reducing exposure to climate-related hazards. Simultaneously, environmental management supports the sustainable utilization of natural resources through ecosystem conservation, pollution control, green infrastructure development, and restoration of degraded landscapes. The integration of these two approaches enables planners and policymakers to create communities that are environmentally sustainable, economically productive, and socially inclusive.

This chapter examines the role of spatial planning in developing climate-smart communities through resilient land use and environmental management. It discusses the theoretical foundations of spatial planning, explores the challenges posed by climate change and rapid urbanization, evaluates sustainable planning strategies and governance mechanisms, and highlights future directions for fostering resilient, adaptive, and environmentally sustainable communities.

The concepts presented in Table demonstrate that climate-smart communities are not developed through isolated planning interventions but through the integration of resilient land use planning, environmental conservation, technological innovation, risk assessment, and participatory governance. Together, these components create a comprehensive planning framework that strengthens community resilience, supports sustainable development, and enhances the capacity of settlements to adapt to current and future climate challenges.

Table: Major Components of Spatial Planning for Climate-Smart Communities

Component	Description	Contribution to Climate-Smart Communities
Land Use Planning	Allocation of land for residential, commercial, industrial, agricultural, and ecological purposes	Reduces environmental conflicts and minimizes climate risks
Environmental Management	Conservation of ecosystems, pollution control, and natural resource management	Enhances ecological sustainability and ecosystem resilience
Climate Risk Assessment	Identification of climate hazards and vulnerable areas	Supports informed planning and disaster preparedness
Green Infrastructure	Parks, urban forests, wetlands, green roofs, and permeable landscapes	Improves biodiversity, reduces urban heat islands, and manages stormwater
Disaster Risk Reduction	Hazard mapping, resilient infrastructure, and emergency preparedness	Minimizes loss of life, property, and ecosystem services
Community Participation	Stakeholder engagement in planning and decision-making	Promotes inclusive and locally appropriate development
Smart Technologies	GIS, Remote Sensing, Artificial Intelligence, Internet of Things (IoT), and digital monitoring	Improves planning accuracy, monitoring, and adaptive management

Source: Adapted from United Nations Human Settlements Programme (2020); Intergovernmental Panel on Climate Change (2022); United Nations (2015).

Foundations of Spatial Planning for Climate Smart Communities

Spatial planning provides the strategic framework for organizing the use of land, natural resources, infrastructure, and human activities to achieve sustainable and resilient development. As climate change continues to alter environmental conditions and intensify natural hazards, spatial planning has evolved from a conventional land-use allocation process to a comprehensive decision-making tool that integrates climate adaptation, disaster risk reduction, environmental conservation, and socio-economic development. Climate-smart communities are built upon planning systems that recognize the interdependence of ecological integrity, economic growth, and social well-being. Consequently, resilient land use planning and effective environmental management have become essential pillars of sustainable spatial planning.

1. Concept and Principles of Spatial Planning

Spatial planning is a systematic process that guides the distribution of land uses and development activities while considering environmental, economic, and social objectives. It facilitates coordinated decision-making across different sectors, including transportation, housing, industry, agriculture, biodiversity conservation, and infrastructure development. Unlike traditional planning approaches that primarily focus on physical development, spatial planning incorporates long-term environmental sustainability, climate resilience, and stakeholder participation into development strategies.

The guiding principles of spatial planning include sustainability, equity, efficiency, resilience, integration, and adaptability. Sustainable planning ensures the responsible utilization of natural resources without compromising future generations. Equity promotes balanced regional development and equal access to public services and resources. Efficiency encourages optimal use of land and infrastructure, while resilience enhances the capacity of communities to withstand environmental disturbances. Integration emphasizes coordination among government agencies, private organizations, and local communities, whereas adaptability allows planning systems to respond effectively to changing climatic and socio-economic conditions.

Table: Principles of Spatial Planning and Their Significance

Principle	Description	Planning Outcome
Sustainability	Efficient utilization of land and natural resources	Long-term environmental protection
Integration	Coordination among multiple sectors and stakeholders	Comprehensive development planning
Resilience	Capacity to anticipate and recover from climate impacts	Reduced disaster vulnerability
Equity	Fair distribution of resources and development opportunities	Inclusive community development
Efficiency	Optimal allocation of land, infrastructure, and investments	Improved resource utilization
Adaptability	Flexibility to respond to environmental and socio-economic changes	Climate-responsive planning

Source: Adapted from United Nations Human Settlements Programme (UN-Habitat, 2020); European Commission (1999).

2. Characteristics of Climate Smart Communities

Climate-smart communities are designed to minimize climate risks while promoting environmental sustainability, economic productivity, and social inclusiveness. These communities integrate climate adaptation and mitigation into local

development planning, ensuring that future growth does not increase vulnerability to climate-induced hazards.

Key characteristics include resilient infrastructure capable of withstanding floods, storms, droughts, and heatwaves; efficient land use that avoids environmentally sensitive or hazard-prone areas; sustainable transportation systems; renewable energy utilization; effective waste and water management; green infrastructure; and active public participation in planning and governance. Climate-smart communities also prioritize ecosystem conservation because healthy ecosystems provide natural protection against floods, landslides, coastal erosion, and urban heat island effects.

The use of climate information, environmental monitoring systems, and technological innovations further enhances the adaptive capacity of these communities. By integrating ecological considerations into planning decisions, climate-smart communities simultaneously improve environmental quality, economic resilience, and public health.

3. Role of Resilient Land Use Planning

Land use planning is one of the most effective instruments for reducing climate vulnerability. It determines how land is allocated for residential, commercial, industrial, agricultural, recreational, and conservation purposes while minimizing environmental degradation and disaster risks.

Resilient land use planning incorporates climate projections, hazard assessments, ecosystem conservation, and environmental carrying capacity into planning decisions. It discourages development in floodplains, coastal erosion zones, steep slopes, and ecologically sensitive regions while encouraging compact urban development, mixed land use, transit-oriented development, and the preservation of open spaces.

Green spaces, urban forests, wetlands, river corridors, and coastal mangroves function as natural protective systems by reducing flood intensity, improving groundwater recharge, lowering urban temperatures, and enhancing biodiversity. Integrating these natural assets into land use planning strengthens ecosystem resilience while reducing dependence on expensive engineered solutions.

Moreover, resilient land use planning supports sustainable agriculture, protects water resources, minimizes land degradation, and enhances food security, making it an essential strategy for climate adaptation and sustainable development.

4. Environmental Management in Spatial Planning

Environmental management complements spatial planning by ensuring that development activities remain within ecological limits while maintaining the integrity of natural ecosystems. It encompasses biodiversity conservation, pollution control, watershed management, environmental impact assessment, ecosystem restoration, waste management, and sustainable resource utilization.

Modern spatial planning increasingly adopts ecosystem-based management approaches that recognize the ecological functions performed by forests, wetlands, rivers, coastal ecosystems, and agricultural landscapes. Conserving these ecosystems not only protects biodiversity but also provides valuable ecosystem services such as carbon sequestration, water purification, climate regulation, and disaster risk reduction.

Environmental Impact Assessments (EIAs), Strategic Environmental Assessments (SEAs), and Geographic Information Systems (GIS) have become indispensable planning tools for evaluating the environmental consequences of development projects before implementation. These tools improve decision-making by identifying environmentally sensitive areas, assessing cumulative impacts, and supporting evidence-based planning.

Table: Relationship between Spatial Planning and Environmental Management

Spatial Planning Function	**Environmental Management Objective**	**Expected Outcome**
Land allocation	Protection of ecologically sensitive areas	Biodiversity conservation
Urban expansion planning	Pollution prevention and resource efficiency	Sustainable urban growth
Infrastructure planning	Environmental Impact Assessment	Reduced ecological degradation
Watershed planning	Water conservation and flood management	Improved water security
Green infrastructure planning	Ecosystem restoration	Enhanced climate resilience
Hazard zoning	Disaster risk reduction	Safer human settlements

Source: Adapted from Intergovernmental Panel on Climate Change (2022); United Nations Environment Programme (2021); Food and Agriculture Organization (2022).

The integration of spatial planning with environmental management provides a holistic framework for sustainable development by balancing ecological conservation with economic growth and social development. Through resilient land use planning, ecosystem protection, and evidence-based environmental governance, climate-smart communities can effectively respond to current and future climate challenges while ensuring long-term environmental sustainability and improved quality of life.

Climate Change Challenges and Resilient Land Use Strategies

Climate change has emerged as one of the most significant challenges affecting the sustainability of human settlements and natural ecosystems. Increasing global temperatures, changing precipitation patterns, rising sea levels, and the growing frequency of extreme weather events have intensified environmental risks, particularly in rapidly urbanizing regions. These challenges are further compounded by unplanned urban expansion, unsustainable land use practices, environmental degradation, and inadequate infrastructure planning. Consequently, spatial planning has shifted from a reactive approach to a proactive strategy that integrates climate adaptation, disaster risk reduction, ecosystem conservation, and resilient land use planning. Developing climate-smart communities requires comprehensive land management practices that minimize vulnerability while enhancing the adaptive capacity of both natural and built environments.

1. Urbanization and Land Use Change

Rapid urbanization is transforming landscapes across the world at an unprecedented pace. The expansion of cities often occurs at the expense of agricultural land, forests, wetlands, and other ecologically sensitive ecosystems. Such land use changes reduce biodiversity, increase surface runoff, intensify urban heat island effects, and diminish the natural capacity of ecosystems to regulate climate and water resources.

Unplanned urban growth frequently results in settlements located within floodplains, coastal zones, landslide-prone slopes, and other environmentally vulnerable areas. The conversion of permeable surfaces into roads, buildings, and pavements reduces groundwater recharge and increases the likelihood of urban flooding during periods of intense rainfall. Similarly, the loss of urban vegetation reduces carbon sequestration, deteriorates air quality, and contributes to rising urban temperatures.

Spatial planning addresses these challenges by promoting compact urban development, mixed land use, transit-oriented development, and the protection of ecological corridors. Such planning approaches reduce urban sprawl, improve resource efficiency, and create healthier, more resilient living environments.

2. Climate Risks Affecting Human Settlements

Climate change has significantly increased the exposure of human settlements to natural hazards. Coastal cities face rising sea levels, saline water intrusion, and storm surges, while inland regions experience floods, droughts, heatwaves, forest fires, and water scarcity. Mountainous areas are increasingly vulnerable to landslides and glacial lake outburst floods due to changing climatic conditions.

The severity of climate impacts often depends upon existing land use patterns, infrastructure quality, population density, and socio-economic conditions. Poorly

planned settlements lacking adequate drainage systems, green spaces, and resilient infrastructure are particularly susceptible to climate-induced disasters.

Spatial planning contributes to climate adaptation by incorporating climate risk assessments into development plans. Hazard mapping, vulnerability assessments, floodplain zoning, coastal setback regulations, and climate-sensitive building codes help reduce exposure to environmental hazards. These measures ensure that future development is directed toward safer locations while minimizing disaster-related losses.

Table: Major Climate Risks and Spatial Planning Responses

Climate Risk	Potential Impacts	Spatial Planning Response
Flooding	Damage to infrastructure, loss of life, water contamination	Floodplain zoning, improved drainage systems, wetland conservation
Heatwaves	Urban heat islands, public health risks	Urban forests, green roofs, open green spaces
Sea-level rise	Coastal erosion, saline intrusion, displacement	Coastal setback zones, mangrove restoration, resilient infrastructure
Drought	Water scarcity, agricultural losses	Rainwater harvesting, watershed management, water-efficient land use
Landslides	Infrastructure damage, habitat destruction	Slope stabilization, restricted development on steep terrain
Forest fires	Biodiversity loss, air pollution	Fire-resistant landscape planning, buffer zones, forest management

Source: Adapted from Intergovernmental Panel on Climate Change (2022); United Nations Office for Disaster Risk Reduction (2023).

3. Nature-Based Solutions and Green Infrastructure

Nature-Based Solutions (NbS) have emerged as effective approaches for enhancing climate resilience while simultaneously conserving biodiversity and improving ecosystem services. These solutions involve protecting, restoring, and sustainably managing natural ecosystems to address environmental challenges.

Green infrastructure represents one of the most widely adopted Nature-Based Solutions within spatial planning. It includes urban forests, wetlands, parks, green belts, riparian buffers, green roofs, permeable pavements, rain gardens, and bioswales that collectively improve ecological functioning within urban and rural landscapes.

Urban forests reduce ambient temperatures, improve air quality, sequester carbon dioxide, and provide recreational opportunities. Wetlands function as natural flood retention systems by storing excess rainfall and reducing downstream flooding.

Mangrove forests protect coastal communities against storm surges and coastal erosion while supporting fisheries and biodiversity.

Compared with conventional engineering solutions, green infrastructure often provides multiple ecological, economic, and social benefits simultaneously. It enhances biodiversity conservation, improves public health, increases property values, and reduces infrastructure maintenance costs.

Table: Nature-Based Solutions for Climate Adaptation

Nature-Based Solution	Primary Function	Climate Adaptation Benefit
Urban Forests	Carbon sequestration and cooling	Reduces urban heat island effect
Wetlands	Floodwater storage	Minimizes flood risk
Mangrove Forests	Coastal protection	Reduces storm surge impacts
Green Roofs	Temperature regulation	Lowers building energy consumption
Rain Gardens	Stormwater infiltration	Reduces urban flooding
Ecological Corridors	Habitat connectivity	Enhances biodiversity resilience

Source: Adapted from International Union for Conservation of Nature (2020); United Nations Environment Programme (2022).

4. Disaster Risk Reduction through Spatial Planning

Disaster Risk Reduction (DRR) has become an integral component of climate-responsive spatial planning. Rather than responding to disasters after they occur, modern planning emphasizes proactive risk reduction through informed land use decisions, resilient infrastructure development, and ecosystem conservation.

Hazard mapping and Geographic Information Systems (GIS) enable planners to identify vulnerable regions and guide future development away from high-risk areas. Climate-resilient infrastructure, including elevated buildings, flood-resistant transportation networks, sustainable drainage systems, and resilient utility services, significantly reduces disaster vulnerability.

Spatial planning also promotes the development of evacuation routes, emergency shelters, and early warning systems that strengthen community preparedness. Integrating disaster risk reduction into regional and local development plans ensures that resilience becomes a fundamental consideration throughout the planning process rather than an afterthought.

Furthermore, community participation enhances disaster preparedness by incorporating local knowledge into risk assessments and emergency planning. Public awareness programmes, stakeholder engagement, and capacity-building

initiatives improve societal resilience and encourage shared responsibility for climate adaptation.

Overall, resilient land use strategies, supported by nature-based solutions and comprehensive disaster risk reduction measures, provide a robust framework for developing climate-smart communities capable of adapting to future environmental uncertainties. These integrated approaches not only reduce climate vulnerability but also promote sustainable development, environmental conservation, and improved quality of life.

Sustainable Planning Approaches and Governance

The growing complexity of climate change and urban development demands planning approaches that are adaptive, integrated, and participatory. Sustainable spatial planning extends beyond regulating land use by incorporating environmental conservation, climate adaptation, technological innovation, and institutional coordination into the planning process. Effective governance ensures that planning decisions are transparent, evidence-based, and inclusive, enabling communities to respond proactively to environmental challenges. The integration of resilient planning strategies with sound governance mechanisms forms the cornerstone of climate-smart communities capable of achieving long-term environmental sustainability and socio-economic development.

1. Integrated Spatial Planning Frameworks

Integrated spatial planning is a multidisciplinary approach that coordinates land use, transportation, housing, environmental protection, economic development, and disaster risk management within a unified planning framework. Rather than addressing these sectors independently, integrated planning recognizes their interdependence and seeks to optimize resource utilization while minimizing environmental degradation.

One of the primary objectives of integrated planning is to balance development needs with ecosystem conservation. This involves identifying environmentally sensitive areas, conserving ecological corridors, promoting compact urban growth, and ensuring equitable access to infrastructure and public services. Spatial planning frameworks also incorporate watershed management, biodiversity conservation, renewable energy development, and climate adaptation into regional development plans.

Modern planning increasingly relies on spatial databases, environmental indicators, and predictive models to evaluate development scenarios before implementation. Such evidence-based planning reduces uncertainty and improves decision-making, ensuring that future development remains environmentally sustainable and economically viable.

2. Climate Adaptation and Mitigation Strategies

Climate-smart communities require planning strategies that simultaneously reduce greenhouse gas emissions (mitigation) and enhance the ability of communities to cope with climate impacts (adaptation). Although these approaches have distinct objectives, they are most effective when implemented together through integrated spatial planning.

Adaptation strategies include climate-resilient infrastructure, sustainable drainage systems, floodplain management, coastal protection, urban greening, water conservation, and drought-resistant landscaping. These measures reduce the vulnerability of settlements while enhancing ecosystem resilience.

Mitigation strategies focus on lowering carbon emissions through compact urban development, renewable energy integration, sustainable public transportation, energy-efficient buildings, and preservation of forests and green spaces. Urban planning plays a crucial role in reducing dependence on private vehicles by promoting walkable neighbourhoods, cycling infrastructure, and efficient mass transit systems.

Together, adaptation and mitigation strategies strengthen environmental sustainability while contributing to global climate goals and national development priorities.

Table 4.1 Climate Adaptation and Mitigation Measures in Spatial Planning

Planning Strategy	Adaptation Benefit	Mitigation Benefit
Urban Green Spaces	Reduces heat stress and flooding	Enhances carbon sequestration
Compact Urban Development	Minimizes disaster exposure	Reduces transport-related emissions
Renewable Energy Infrastructure	Improves energy security	Reduces fossil fuel dependence
Sustainable Drainage Systems	Controls urban flooding	Supports groundwater recharge
Climate-Resilient Buildings	Withstands extreme weather events	Improves energy efficiency
Public Transportation Networks	Enhances mobility during disasters	Reduces greenhouse gas emissions

Source: Adapted from Intergovernmental Panel on Climate Change (2022); United Nations Human Settlements Programme (2022).

3. Smart Technologies in Spatial Planning

Technological advancements have significantly improved the effectiveness of spatial planning by enabling planners to collect, analyse, and visualize spatial information with greater accuracy. Smart technologies facilitate evidence-based

decision-making, improve environmental monitoring, and strengthen climate adaptation planning.

Geographic Information Systems (GIS) enable the integration of spatial data for land suitability analysis, hazard mapping, infrastructure planning, and environmental assessment. Remote sensing provides real-time information on land use changes, vegetation cover, urban expansion, and ecosystem degradation through satellite imagery.

Artificial Intelligence (AI) and Machine Learning (ML) are increasingly used to predict flood risks, optimize transportation networks, forecast urban growth, and support environmental modelling. Internet of Things (IoT) technologies provide continuous monitoring of environmental conditions such as air quality, water levels, rainfall, and temperature through sensor networks.

Digital twins, cloud computing, and big data analytics further support planners by simulating future development scenarios, evaluating infrastructure performance, and identifying climate vulnerabilities before implementation. These innovations enhance planning efficiency while supporting adaptive governance.

Table: Smart Technologies Supporting Climate-Smart Spatial Planning

Technology	Major Application	Planning Benefit
Geographic Information Systems (GIS)	Spatial analysis and mapping	Improved land use planning
Remote Sensing	Environmental monitoring	Real-time assessment of land use change
Artificial Intelligence	Predictive modelling	Improved climate risk forecasting
Internet of Things (IoT)	Environmental sensors	Continuous monitoring of environmental conditions
Big Data Analytics	Urban data integration	Evidence-based planning decisions
Digital Twin Technology	Virtual simulation of cities	Infrastructure optimization and resilience planning

Source: Adapted from United Nations Environment Programme (2022); World Bank (2023).

4. Community Participation and Environmental Governance

Effective spatial planning depends not only on technical expertise but also on active participation by local communities and stakeholders. Community participation enhances the legitimacy, transparency, and effectiveness of planning processes by incorporating local knowledge, cultural values, and community priorities into decision-making.

Participatory planning enables governments, civil society organizations, researchers, private sector institutions, and local residents to collaborate in identifying environmental challenges and developing locally appropriate solutions.

Indigenous and traditional knowledge often complements scientific research, particularly in ecosystem conservation, water resource management, and disaster preparedness.

Environmental governance refers to the institutional arrangements, legal frameworks, and policy mechanisms that regulate environmental management and sustainable development. Good governance is characterized by transparency, accountability, equity, stakeholder engagement, scientific decision-making, and effective enforcement of environmental regulations.

Public awareness programmes, environmental education, and capacity-building initiatives further strengthen community resilience by promoting responsible resource use and encouraging climate-conscious behaviour.

5. Policy and Institutional Frameworks

The successful implementation of climate-smart spatial planning requires strong policy support and coordinated institutional mechanisms. National governments establish legal frameworks that guide land use planning, environmental protection, climate adaptation, and infrastructure development, while regional and local authorities translate these policies into practical planning initiatives.

International agreements such as the Paris Agreement, the Sendai Framework for Disaster Risk Reduction, the New Urban Agenda, and the 2030 Agenda for Sustainable Development provide strategic direction for climate-resilient planning. These frameworks encourage countries to integrate climate adaptation, sustainable land management, ecosystem restoration, and disaster resilience into national planning systems.

At the national and local levels, planning authorities increasingly adopt integrated land use policies, environmental impact assessments, climate action plans, and resilience strategies to guide sustainable development. Institutional coordination among urban planners, environmental agencies, disaster management authorities, academic institutions, and community organizations ensures that planning decisions are coherent and mutually reinforcing.

In conclusion, sustainable planning approaches supported by effective governance, technological innovation, and participatory decision-making are fundamental to the development of climate-smart communities. By integrating climate adaptation, resilient infrastructure, ecosystem conservation, and inclusive governance within comprehensive spatial planning frameworks, societies can build settlements that are environmentally sustainable, socially equitable, and economically resilient in the face of future climate uncertainties.

Future Directions for Climate Smart Communities

The transition towards climate-smart communities requires continuous innovation in spatial planning, environmental governance, and sustainable land management. As climate change intensifies and urban populations continue to grow, conventional

planning approaches must evolve into adaptive systems capable of responding to dynamic environmental, social, and economic challenges. Future spatial planning should emphasize resilience, technological integration, ecosystem conservation, and inclusive governance to create communities that are sustainable, disaster-resilient, and environmentally responsible. Achieving this vision requires collaboration among governments, researchers, planners, private organizations, and local communities while aligning planning practices with global sustainability goals.

1. Innovative Planning Practices

Future spatial planning will increasingly rely on innovative planning methodologies that integrate scientific research, climate projections, and digital technologies into decision-making. Predictive planning models using Artificial Intelligence (AI), Geographic Information Systems (GIS), Remote Sensing, and Digital Twin technologies will enable planners to simulate future urban growth, assess environmental risks, and evaluate alternative development scenarios before implementation.

Climate-responsive urban design will become a central planning principle by promoting mixed land use, compact urban forms, green transportation systems, energy-efficient buildings, and climate-resilient infrastructure. The concept of the 15-minute city, where residents can access essential services within a short walking or cycling distance, is gaining recognition as an effective strategy for reducing transportation emissions, improving accessibility, and enhancing urban resilience. Such innovative planning practices contribute to healthier, more efficient, and environmentally sustainable communities.

2. Ecosystem-Based Land Management

Ecosystem-based land management is expected to become one of the most important approaches for enhancing climate resilience. Rather than relying solely on engineered infrastructure, future planning will increasingly utilize natural ecosystems as protective infrastructure. Forests, wetlands, mangroves, rivers, coastal ecosystems, and urban green spaces provide valuable ecosystem services that reduce disaster risks while supporting biodiversity conservation.

Restoration of degraded ecosystems, expansion of urban forests, protection of biodiversity corridors, watershed conservation, and sustainable agricultural landscapes will strengthen ecological resilience and improve environmental quality. Integrating Nature-Based Solutions (NbS) into land use planning offers cost-effective alternatives for climate adaptation while simultaneously improving air quality, water security, carbon sequestration, and public health.

The future of spatial planning therefore lies in balancing infrastructure development with ecosystem conservation to maintain ecological integrity while supporting economic development.

Table 5.1 Emerging Approaches for Developing Climate-Smart Communities

Emerging Approach	Major Objective	Expected Benefits
Digital Spatial Planning	Data-driven planning and decision-making	Improved planning accuracy and efficiency
Nature-Based Solutions	Ecosystem restoration and conservation	Climate adaptation and biodiversity protection
Green Infrastructure	Sustainable urban landscapes	Reduced heat islands and improved stormwater management
Circular Economy Planning	Efficient resource utilization	Reduced waste generation and carbon emissions
Smart City Technologies	Digital governance and monitoring	Enhanced public service delivery and resilience
Participatory Planning	Inclusive stakeholder engagement	Greater social acceptance and sustainable implementation

Source: Adapted from United Nations Human Settlements Programme (UN-Habitat, 2022); United Nations Environment Programme (2022); World Bank (2023).

3. Sustainable Urban and Rural Development

Future climate-smart planning must recognize the interdependence between urban and rural areas. Sustainable urban development depends on healthy rural ecosystems that provide food, water, energy, and ecosystem services, while rural communities rely on urban centres for markets, infrastructure, education, healthcare, and employment.

Integrated regional planning should therefore strengthen urban-rural linkages through sustainable transportation networks, efficient resource distribution, climate-resilient agriculture, renewable energy systems, and ecosystem conservation. Balanced regional development can reduce migration pressures, improve economic opportunities, and enhance environmental sustainability.

Furthermore, promoting low-carbon development pathways through renewable energy, sustainable mobility, water-efficient infrastructure, and green building practices will contribute significantly to achieving national and international climate objectives.

4. Challenges and Opportunities

Despite significant progress in climate-responsive planning, several challenges continue to hinder the development of climate-smart communities. These include inadequate financial resources, fragmented institutional responsibilities, weak policy implementation, insufficient technical expertise, limited availability of spatial data, rapid population growth, and conflicting land-use priorities. In many

developing regions, informal settlements, unplanned urban expansion, and environmental degradation further complicate sustainable planning efforts.

However, these challenges also create opportunities for innovation and institutional reform. Advances in geospatial technologies, climate science, renewable energy, environmental monitoring, and digital governance provide planners with powerful tools for improving planning effectiveness. International cooperation, climate finance mechanisms, public-private partnerships, and capacity-building programmes offer additional support for implementing resilient planning initiatives.

5. Way Forward

Developing climate-smart communities requires a long-term vision supported by integrated policies, scientific research, technological innovation, and active public participation. Future planning should prioritize climate resilience, ecosystem conservation, sustainable land use, and inclusive governance while ensuring that development remains socially equitable and environmentally sustainable.

Strengthening institutional coordination among planning agencies, environmental authorities, disaster management organizations, research institutions, and local communities will improve policy implementation and resource management. Investments in climate-resilient infrastructure, ecosystem restoration, smart technologies, and environmental education should be regarded as essential components of sustainable development strategies.

Ultimately, spatial planning will continue to play a transformative role in shaping resilient communities capable of adapting to climate change while protecting natural ecosystems and improving human well-being. By integrating resilient land use planning with effective environmental management, societies can create sustainable settlements that support economic prosperity, environmental integrity, and a higher quality of life for present and future generations.

References

1. European Commission. (1999). European Spatial Development Perspective: Towards balanced and sustainable development of the territory of the European Union. Office for Official Publications of the European Communities.
2. Intergovernmental Panel on Climate Change. (2022). Climate change 2022: Impacts, adaptation and vulnerability. Cambridge University Press.
3. International Union for Conservation of Nature. (2020). Global standard for nature-based solutions: A user-friendly framework for the verification, design and scaling up of NbS (1st ed.). IUCN.
4. United Nations. (2015). Transforming our world: The 2030 agenda for sustainable development. United Nations.
5. United Nations Environment Programme. (2022). Adaptation gap report 2022: Too little, too slow – Climate adaptation failure puts world at risk. UNEP.

6. United Nations Human Settlements Programme. (2020). World cities report 2020: The value of sustainable urbanization. UN-Habitat.
7. United Nations Office for Disaster Risk Reduction. (2023). Global assessment report on disaster risk reduction 2023. UNDRR.
8. World Bank. (2023). Thriving: Making cities green, resilient, and inclusive in a changing climate. World Bank.
9. World Resources Institute. (2021). World resources report: Creating a sustainable food future. World Resources Institute.

Advances in Geography and Environmental Research

ISBN: 978-93-49938-93-9| Year: 2026 | pp: 46 - 58

Environmental Impact Assessment Integrating Ecological Conservation Social Responsibility and Economic Development

Suchandana Dutta

Assistant Professor, Department of Statistics, Thakur College of Science and Commerce, Kandivali, Mumbai

Email: suchandanad4@gmail.com

Article DOI Link: https://zenodo.org/uploads/21476240

DOI: 10.5281/zenodo.21476240

Abstract

Environmental Impact Assessment (EIA) is a vital decision-making tool that promotes sustainable development by integrating environmental protection, social responsibility, and economic growth into project planning and implementation. This chapter examines the evolving role of EIA in conserving biodiversity, protecting ecosystems, and ensuring the sustainable management of natural resources while addressing the social and economic implications of development activities. It discusses the significance of Social Impact Assessment, stakeholder participation, environmental justice, corporate social responsibility, and effective environmental governance in enhancing the credibility and effectiveness of EIA. The chapter further explores the contribution of environmental valuation, green infrastructure, and circular economy principles in achieving sustainable economic development. Emerging technologies such as Geographic Information Systems (GIS), Remote Sensing, Artificial Intelligence (AI), and Big Data are highlighted for their role in improving environmental assessment and monitoring. Finally, the chapter outlines current challenges, future directions, and policy measures for strengthening Environmental Impact Assessment as an indispensable instrument for achieving resilient and sustainable development.

Keywords: Environmental Impact Assessment, Ecological Conservation, Social Responsibility, Sustainable Development, Environmental Governance

Introduction

Environmental Impact Assessment (EIA) has emerged as one of the most significant environmental management tools for promoting sustainable development while

balancing ecological integrity, social well-being, and economic progress. Rapid industrialization, urban expansion, infrastructure development, and increasing exploitation of natural resources have intensified environmental degradation across the globe. These developmental activities have contributed to deforestation, biodiversity loss, climate change, water pollution, air pollution, soil degradation, and depletion of natural resources, thereby posing substantial challenges to environmental sustainability. In this context, Environmental Impact Assessment serves as a systematic decision-support process that evaluates the potential environmental, social, and economic consequences of proposed development projects before their implementation. By identifying potential adverse impacts at an early stage, EIA enables policymakers, planners, industries, and communities to adopt appropriate mitigation measures that minimize environmental damage while maximizing developmental benefits.

The concept of Environmental Impact Assessment gained international prominence following the enactment of the United States National Environmental Policy Act (NEPA) in 1969, which established a formal mechanism for integrating environmental considerations into developmental planning. Since then, EIA has evolved into an internationally recognized environmental governance instrument and has been adopted by numerous countries through legislative and regulatory frameworks. International organizations such as the United Nations Environment Programme (UNEP), the World Bank, and the International Association for Impact Assessment (IAIA) have consistently emphasized the importance of EIA in achieving environmentally responsible development. Today, EIA is regarded not merely as a regulatory requirement but as an essential planning and management tool that facilitates informed decision-making, enhances environmental accountability, and supports long-term sustainability objectives.

The contemporary approach to Environmental Impact Assessment extends beyond the traditional evaluation of physical environmental impacts to encompass ecological conservation, social responsibility, and economic sustainability. Modern EIA frameworks recognize that development projects influence complex socio-ecological systems where environmental, social, and economic dimensions are closely interconnected. Consequently, effective impact assessment requires multidisciplinary analysis involving ecological risk assessment, biodiversity conservation, public participation, social equity, environmental health, climate resilience, and sustainable resource management. Such an integrated approach ensures that developmental interventions contribute not only to economic growth but also to the protection of ecosystem services and the improvement of community well-being.

Ecological conservation has become a central component of Environmental Impact Assessment due to increasing concerns regarding habitat destruction, species extinction, ecosystem fragmentation, and climate-related vulnerabilities. Assessing

impacts on forests, wetlands, rivers, coastal ecosystems, wildlife habitats, and biodiversity enables decision-makers to implement preventive and compensatory measures that safeguard ecological integrity. Simultaneously, the incorporation of social responsibility within EIA emphasizes meaningful stakeholder engagement, public consultation, environmental justice, and the protection of indigenous and local communities whose livelihoods depend upon natural resources. This participatory approach improves transparency, strengthens public confidence, and enhances the legitimacy of environmental decision-making processes.

Economic development remains a fundamental objective for both developed and developing nations. However, the pursuit of economic growth without adequate environmental safeguards often results in irreversible ecological damage and substantial social costs. Environmental Impact Assessment provides a practical framework for harmonizing economic aspirations with environmental protection by evaluating alternative project designs, assessing environmental costs and benefits, promoting resource efficiency, and encouraging sustainable investment practices. Increasingly, governments and industries are recognizing that environmentally responsible development enhances long-term economic resilience, reduces regulatory and operational risks, and contributes to improved environmental, social, and governance (ESG) performance.

Advancements in digital technologies have further strengthened the effectiveness of Environmental Impact Assessment. Geographic Information Systems (GIS), Remote Sensing, Artificial Intelligence (AI), machine learning, environmental modelling, big data analytics, and real-time environmental monitoring have significantly improved the accuracy of impact prediction and environmental decision-making. These technological innovations facilitate comprehensive baseline assessments, cumulative impact analysis, climate risk modelling, and continuous post-project monitoring, thereby enhancing the scientific reliability and efficiency of EIA processes.

As environmental challenges become increasingly complex due to climate change, urbanization, resource scarcity, and growing societal expectations, Environmental Impact Assessment continues to evolve as a dynamic instrument for sustainable environmental governance. Integrating ecological conservation, social responsibility, and economic development within the EIA framework enables governments, industries, and communities to make balanced decisions that foster environmental protection, inclusive development, and long-term resilience. This chapter examines these interconnected dimensions by exploring the ecological, social, and economic significance of Environmental Impact Assessment, while highlighting emerging technologies, policy innovations, implementation challenges, and future opportunities for strengthening sustainable development practices.

Environmental Impact Assessment and Ecological Conservation

1. Biodiversity Conservation and Ecosystem Protection

Biodiversity forms the foundation of healthy ecosystems by maintaining ecological balance, supporting ecosystem services, and ensuring the sustainability of natural resources. It encompasses the diversity of genes, species, and ecosystems that collectively sustain life on Earth. However, rapid industrialization, urbanization, mining, infrastructure development, dam construction, transportation networks, and agricultural expansion have accelerated habitat destruction and fragmentation, resulting in significant biodiversity loss worldwide. Environmental Impact Assessment (EIA) serves as a preventive environmental management tool by identifying potential impacts on biodiversity before project implementation, thereby enabling decision-makers to adopt scientifically informed mitigation strategies.

Within the EIA framework, biodiversity assessment begins with baseline ecological surveys that document existing flora, fauna, endangered species, protected habitats, wetlands, forests, rivers, grasslands, and ecological corridors. These baseline studies establish the environmental status of the project area and help identify ecologically sensitive zones requiring special protection. Comprehensive biodiversity inventories also facilitate the prediction of direct, indirect, cumulative, and long-term impacts associated with developmental activities.

The application of biodiversity indicators during EIA enables environmental professionals to evaluate habitat quality, species richness, ecosystem health, and ecological resilience. Modern ecological assessments increasingly incorporate Geographic Information Systems (GIS), remote sensing technologies, drone-based surveys, camera traps, satellite imagery, and environmental DNA (eDNA) analysis to improve the accuracy of biodiversity monitoring. These technological advancements provide reliable spatial information that supports evidence-based environmental decision-making.

Mitigation measures constitute an essential component of biodiversity conservation within EIA. The internationally accepted mitigation hierarchy follows four sequential steps: avoidance, minimization, restoration, and biodiversity offsetting. Avoidance focuses on redesigning projects to prevent ecological damage, while minimization reduces unavoidable impacts through engineering and operational modifications. Restoration activities rehabilitate degraded ecosystems after project completion, whereas biodiversity offsets compensate for residual ecological losses through conservation initiatives undertaken elsewhere.

Protected areas, wildlife sanctuaries, national parks, biosphere reserves, Ramsar wetlands, and biodiversity hotspots require special consideration during environmental impact assessment. Projects located near ecologically sensitive landscapes demand rigorous ecological evaluation to prevent habitat fragmentation, wildlife mortality, invasive species introduction, and ecosystem degradation.

Ecological connectivity through wildlife corridors has become increasingly important in infrastructure planning to reduce human–wildlife conflicts and maintain species migration routes.

The conservation of biodiversity contributes directly to sustainable development by maintaining ecosystem productivity, ensuring food security, regulating climate, preserving genetic resources, and supporting livelihoods dependent upon ecosystem services. Consequently, biodiversity conservation has evolved from being a supplementary consideration to becoming a central objective of contemporary Environmental Impact Assessment.

Table: Biodiversity Conservation Measures within Environmental Impact Assessment

Component	Purpose	Expected Outcome
Baseline biodiversity survey	Document ecological resources	Accurate environmental database
Habitat mapping	Identify sensitive ecosystems	Protection of critical habitats
Species inventory	Record flora and fauna diversity	Conservation planning
Impact prediction	Assess ecological consequences	Informed decision-making
Mitigation hierarchy	Reduce environmental damage	Sustainable project implementation
Ecological monitoring	Evaluate post-project impacts	Adaptive environmental management

Source: Compiled by the Author.

2. Assessment of Impacts on Air, Water, Soil, and Forest Resources

Environmental Impact Assessment systematically evaluates the influence of development projects on the major components of the natural environment, including air quality, water resources, soil characteristics, forests, and associated ecological processes. These environmental components are interdependent; therefore, adverse impacts on one component frequently generate cascading effects across entire ecosystems. Comprehensive environmental assessment enables planners to identify risks early and implement preventive measures before irreversible environmental degradation occurs.

Air quality assessment involves estimating emissions generated during both construction and operational phases of projects. Major pollutants include particulate matter ($PM_{2.5}$ and PM_{10}), sulfur dioxide, nitrogen oxides, carbon monoxide, volatile organic compounds, and greenhouse gases. Dispersion modelling techniques help predict pollutant concentrations under varying meteorological conditions, allowing authorities to determine compliance with environmental standards and identify appropriate pollution control technologies.

Water resource assessment examines both surface water and groundwater systems. Development projects may alter natural drainage patterns, reduce groundwater recharge, increase sedimentation, contaminate water bodies, or affect aquatic biodiversity. Hydrological modelling, water quality monitoring, and watershed analysis are therefore integral components of environmental assessment. Effective mitigation strategies include wastewater treatment systems, rainwater harvesting, sediment control structures, buffer zones, and integrated watershed management practices.

Soil impact assessment evaluates changes in soil fertility, erosion rates, contamination levels, compaction, and land stability resulting from project activities. Construction operations often disturb topsoil, increase erosion, and reduce agricultural productivity. Soil conservation measures such as topsoil preservation, vegetation restoration, erosion control structures, and sustainable land management practices significantly reduce environmental degradation.

Forest ecosystems represent some of the most environmentally valuable landscapes because they provide habitat for wildlife, regulate climate, conserve biodiversity, store carbon, protect watersheds, and support indigenous livelihoods. Environmental Impact Assessment examines forest diversion, habitat fragmentation, edge effects, forest fires, invasive species, and changes in ecological connectivity. Compensatory afforestation, ecological restoration, buffer zone management, and sustainable forest management plans are commonly adopted to minimize forest-related impacts.

Increasingly, cumulative environmental assessments are being incorporated into EIA to evaluate the combined effects of multiple developmental activities occurring within a region. Such integrated assessment provides a more realistic understanding of environmental change than isolated project-level analysis.

Table: Major Environmental Components Evaluated during Environmental Impact Assessment

Environmental Component	Major Impacts Assessed	Common Mitigation Measures
Air	Emissions and air pollution	Emission control technologies, green belts
Water	Pollution, groundwater depletion	Wastewater treatment, rainwater harvesting
Soil	Erosion, contamination	Soil conservation and land restoration
Forest	Deforestation and habitat fragmentation	Afforestation and habitat restoration
Noise	Construction and operational disturbance	Noise barriers and operational controls

Source: Compiled by the Author.

3. Ecosystem Services and Natural Resource Management

The ecosystem services approach has become an important dimension of modern Environmental Impact Assessment because it recognizes the economic and social value of ecological systems. Ecosystem services refer to the direct and indirect benefits that humans obtain from nature, including food production, freshwater availability, climate regulation, pollination, carbon sequestration, nutrient cycling, flood control, and recreational opportunities. Recognizing these services during environmental assessment helps ensure that development decisions account for both market and non-market environmental values.

Natural resource management within EIA seeks to balance developmental requirements with the long-term sustainability of renewable and non-renewable resources. Resource inventories conducted during environmental assessment identify existing conditions relating to forests, minerals, water resources, agricultural land, fisheries, and biodiversity. These inventories facilitate resource-efficient planning while preventing excessive exploitation.

Environmental valuation techniques such as contingent valuation, travel cost method, replacement cost method, and ecosystem accounting increasingly support environmental decision-making by assigning economic values to ecological services that are traditionally overlooked. Such valuation strengthens policy decisions by demonstrating that ecosystem degradation often imposes substantial long-term economic costs exceeding the short-term benefits of unsustainable development.

Integrated Natural Resource Management (INRM) further enhances environmental sustainability by promoting coordinated management of land, water, forests, biodiversity, and agricultural resources. EIA encourages project developers to adopt resource-efficient technologies, renewable energy systems, circular economy principles, waste minimization practices, and sustainable consumption patterns.

Payment for Ecosystem Services (PES), ecological compensation mechanisms, watershed restoration programmes, and nature-based solutions have emerged as innovative approaches that complement traditional mitigation measures. These initiatives encourage conservation while simultaneously supporting rural livelihoods and regional economic development.

The integration of ecosystem services into Environmental Impact Assessment transforms environmental evaluation from a compliance-based exercise into a comprehensive sustainability assessment capable of supporting resilient development pathways.

Integrating Social Responsibility into Environmental Impact Assessment

1. Social Impact Assessment and Community Well-being

Environmental Impact Assessment (EIA) has evolved beyond its traditional focus on biophysical environmental parameters to incorporate social dimensions that directly influence the quality of life of affected communities. Development projects

such as highways, dams, industrial corridors, mining operations, urban expansion, and energy infrastructure often generate profound social consequences, including displacement of communities, changes in livelihoods, alterations in land-use patterns, pressure on public infrastructure, and modifications to local cultural practices. Consequently, Social Impact Assessment (SIA) has become an integral component of modern EIA, enabling project proponents and decision-makers to evaluate the potential positive and negative impacts of development on human populations.

Social Impact Assessment systematically examines demographic characteristics, employment opportunities, public health, education, housing, transportation, access to natural resources, cultural heritage, and community infrastructure. Baseline social surveys provide valuable information regarding the socioeconomic conditions of project-affected populations before project implementation. These data facilitate comparison during subsequent monitoring and evaluation stages, ensuring that developmental benefits are equitably distributed while minimizing adverse social consequences.

Community well-being extends beyond economic indicators and encompasses physical health, psychological security, social cohesion, cultural identity, and environmental quality. Environmental degradation resulting from poorly planned projects may increase respiratory illnesses, waterborne diseases, occupational hazards, and mental stress among local populations. Therefore, EIA incorporates health risk assessments, occupational safety measures, emergency preparedness plans, and environmental health monitoring to safeguard public welfare.

The adoption of livelihood restoration programmes, fair compensation mechanisms, skill development initiatives, employment generation schemes, and community development projects further strengthens the social sustainability of development projects. Increasingly, international financial institutions and national environmental agencies require project developers to prepare Social Management Plans (SMPs) alongside Environmental Management Plans (EMPs), thereby ensuring that environmental protection and community welfare progress simultaneously.

2. Public Participation and Stakeholder Engagement

Meaningful public participation represents one of the fundamental principles of Environmental Impact Assessment because environmental decisions affect a wide range of stakeholders with diverse interests, values, and expectations. Effective stakeholder engagement promotes transparency, accountability, and inclusiveness while improving the scientific quality and social acceptability of development projects.

Stakeholders typically include local communities, government agencies, project proponents, environmental experts, non-governmental organizations (NGOs), academic institutions, indigenous groups, industry representatives, and civil society

organizations. Each stakeholder contributes unique perspectives that enhance environmental decision-making by identifying local environmental concerns, traditional ecological knowledge, and potential social conflicts that may otherwise remain unnoticed.

Public consultation is generally conducted at different stages of the EIA process, including project scoping, preparation of environmental reports, environmental hearings, project review, and post-project monitoring. Public hearings provide opportunities for communities to express concerns regarding pollution, land acquisition, rehabilitation, biodiversity conservation, water availability, cultural heritage, and livelihood security. Incorporating stakeholder feedback improves project design, strengthens environmental safeguards, and reduces the likelihood of conflicts during project implementation.

Technological advancements have significantly expanded opportunities for stakeholder participation. Digital consultation platforms, online environmental disclosure systems, Geographic Information Systems (GIS), mobile applications, social media, and virtual public meetings have improved accessibility and transparency, particularly for geographically dispersed communities. These digital tools facilitate timely dissemination of environmental information and encourage broader public engagement.

Despite these advancements, challenges remain in ensuring effective participation. Limited environmental awareness, unequal access to information, inadequate representation of marginalized communities, language barriers, and procedural complexities often restrict meaningful stakeholder involvement. Strengthening environmental education, enhancing institutional transparency, and promoting participatory governance are therefore essential for improving the effectiveness of public participation within Environmental Impact Assessment.

Table: Major Stakeholders in Environmental Impact Assessment

Stakeholder	**Primary Role**
Government agencies	Regulatory approval and monitoring
Project proponents	Project planning and environmental compliance
Local communities	Identification of local environmental and social concerns
Environmental experts	Scientific assessment and technical evaluation
Non-governmental organizations	Environmental advocacy and public awareness
Academic institutions	Research and technical support
Indigenous communities	Traditional ecological knowledge and resource stewardship

Source: Compiled by the Author.

Collaborative environmental governance that combines scientific research with indigenous knowledge contributes to more resilient and context-specific environmental management strategies. Many countries have increasingly recognized the importance of Free, Prior and Informed Consent (FPIC) for indigenous communities before approving projects that affect their traditional territories and natural resources. Such approaches strengthen social legitimacy while reducing conflicts associated with development projects.

Environmental justice therefore extends beyond legal compliance by promoting ethical decision-making, social inclusion, human rights, cultural preservation, and equitable distribution of environmental benefits and responsibilities.

Environmental Impact Assessment and Economic Development

1. Sustainable Economic Growth and Environmental Protection

Economic development is essential for improving living standards, generating employment, expanding infrastructure, and enhancing national productivity. However, development that disregards environmental considerations often leads to resource depletion, pollution, biodiversity loss, and increased vulnerability to climate change. Environmental Impact Assessment (EIA) provides a structured framework for integrating environmental sustainability into economic planning, ensuring that development projects achieve long-term economic benefits without compromising ecological integrity.

By evaluating the environmental implications of proposed projects before implementation, EIA enables governments and project developers to identify environmentally sustainable alternatives, reduce resource wastage, and prevent costly environmental damage. Sustainable economic growth is therefore supported through responsible land-use planning, efficient resource utilization, pollution prevention, and conservation of ecosystem services. Furthermore, environmentally sustainable projects generally experience lower operational risks, improved regulatory compliance, enhanced investor confidence, and greater public acceptance, thereby contributing to long-term economic resilience.

2. Cost–Benefit Analysis and Environmental Valuation

Cost–Benefit Analysis (CBA) is an important component of Environmental Impact Assessment that assists decision-makers in comparing the economic benefits of development projects with their environmental and social costs. Traditional economic evaluations often focus on direct financial returns while overlooking environmental degradation and ecosystem losses. Modern EIA addresses this limitation by incorporating environmental valuation techniques that estimate the economic significance of natural resources and ecosystem services.

Environmental valuation includes both market and non-market values associated with forests, wetlands, rivers, biodiversity, clean air, and water resources.

Techniques such as contingent valuation, replacement cost, avoided damage cost, travel cost, and hedonic pricing enable policymakers to estimate the monetary value of environmental assets. These approaches facilitate informed decision-making by highlighting the long-term economic consequences of environmental degradation.

Incorporating environmental valuation into project appraisal promotes sustainable investment by encouraging developers to adopt environmentally friendly technologies, minimize pollution, and conserve natural resources. As a result, economic decisions increasingly reflect the true environmental costs and benefits associated with developmental activities.

Table: Economic Tools Applied in Environmental Impact Assessment

Economic Tool	**Purpose**	**Contribution to Sustainable Development**
Cost–Benefit Analysis	Compare project costs and benefits	Supports informed investment decisions
Environmental Valuation	Estimate value of ecosystem services	Recognizes environmental assets
Life Cycle Assessment	Evaluate impacts throughout project life	Improves resource efficiency
Risk Assessment	Estimate environmental and financial risks	Reduces future economic losses
Economic Feasibility Analysis	Assess project viability	Promotes sustainable development

Source: Compiled by the Author.

Emerging Trends Challenges and Future Directions

1. GIS Remote Sensing Artificial Intelligence and Big Data Applications in Environmental Impact Assessment

Technological innovations are transforming Environmental Impact Assessment (EIA) by improving the accuracy, efficiency, and transparency of environmental analysis. Geographic Information Systems (GIS) and Remote Sensing facilitate spatial mapping of land use, vegetation cover, water bodies, biodiversity hotspots, and environmentally sensitive areas. These technologies enable environmental professionals to identify potential impacts at both local and regional scales while supporting evidence-based planning.

Artificial Intelligence (AI) and Machine Learning (ML) are increasingly being applied to predict environmental impacts, analyse large environmental datasets, assess climate-related risks, and develop predictive models for pollution, land degradation, and biodiversity loss. Similarly, Big Data analytics integrates information obtained from satellite imagery, environmental sensors, meteorological

databases, and monitoring stations to provide real-time environmental intelligence. These technological advancements significantly improve baseline studies, cumulative impact assessment, environmental monitoring, and decision-making throughout the project lifecycle.

2. Strategic Environmental Assessment and Cumulative Impact Assessment

While conventional Environmental Impact Assessment focuses primarily on individual projects, Strategic Environmental Assessment (SEA) evaluates the environmental implications of policies, plans, and programmes before project-level decisions are made. SEA facilitates the integration of environmental considerations into broader developmental planning and supports sustainable regional development.

Cumulative Impact Assessment (CIA) has also gained importance because multiple projects within the same geographical region may collectively produce environmental impacts greater than those caused by individual developments. Assessing cumulative impacts enables policymakers to understand long-term environmental changes affecting ecosystems, water resources, biodiversity, air quality, and community health. Integrating SEA and CIA into environmental planning strengthens environmental governance and promotes sustainable resource management.

References

1. Canter, L. W. (1996). Environmental impact assessment (2nd ed.). McGraw-Hill.
2. Cashmore, M., Richardson, T., Hilding-Ryedvik, T., & Emmelin, L. (2010). Evaluating the effectiveness of impact assessment instruments: Theorising the nature and implications of their political constitution. Environmental Impact Assessment Review, 30(6), 371–379.
3. Glasson, J., Therivel, R., & Chadwick, A. (2019). Introduction to environmental impact assessment (5th ed.). Routledge.
4. International Association for Impact Assessment. (2018). Social impact assessment: Guidance for assessing and managing the social impacts of projects. IAIA.
5. Jay, S., Jones, C., Slinn, P., & Wood, C. (2007). Environmental impact assessment: Retrospect and prospect. Environmental Impact Assessment Review, 27(4), 287–300.
6. Morgan, R. K. (2012). Environmental impact assessment: The state of the art. Impact Assessment and Project Appraisal, 30(1), 5–14.
7. Morrison-Saunders, A., Pope, J., Bond, A., & Retief, F. (Eds.). (2021). Handbook of environmental impact assessment. Edward Elgar Publishing.
8. Noble, B. F. (2020). Introduction to environmental impact assessment: A guide to principles and practice (4th ed.). Oxford University Press.

9. Pope, J., Bond, A., Morrison-Saunders, A., & Retief, F. (2017). Advancing the theory and practice of impact assessment: Setting the research agenda. Environmental Impact Assessment Review, 62, 1–10.
10. Sadler, B. (1996). Environmental assessment in a changing world: Evaluating practice to improve performance. International Study of the Effectiveness of Environmental Assessment.
11. Sánchez, L. E. (2020). Environmental impact assessment: Concepts, methodologies, and applications (3rd ed.). Routledge.
12. Therivel, R. (2019). Strategic environmental assessment in action (3rd ed.). Routledge.
13. United Nations. (2015). Transforming our world: The 2030 Agenda for Sustainable Development. United Nations.
14. United Nations Environment Programme. (2018). Assessing environmental impacts: A global review of environmental impact assessment systems. UNEP.
15. World Bank. (2018). Environmental and social framework. World Bank.
16. World Commission on Environment and Development. (1987). Our common future. Oxford University Press.

Advances in Geography and Environmental Research

ISBN: 978-93-49938-93-9| Year: 2026 | pp: 59 - 70

Evaluation of Water Supply Infrastructure and Drinking Water Quality in Shrigonda Municipality

Bapu D. Deokar

Associate Professor, Department of Geography, Maharaja Jivajirao Shinde College, Shrigonda

Email: bapudeokar@gmail.com

Article DOI Link: https://zenodo.org/uploads/21476329

DOI: 10.5281/zenodo.21476329

Abstract

A safe and effective urban water distribution infrastructure is necessary for both public health and sustainable urban development. This study investigates the Shrigonda Municipality's water supply system in Maharashtra, India, with an emphasis on service delivery, water quality, distribution network, storage facilities, and water treatment efficiency. Primary data were gathered from field observations, laboratory testing for treated and untreated water samples, and municipal water delivery records, secondary data was collected from Shrigonda Municipal Council reports. The municipality operates an 11 MLD water treatment plant commissioned in 2018 under the Urban Infrastructure Development Scheme for Small and Medium Towns, sourcing raw water from Glod Dam Water is distributed through seven overhead storage tanks after the usual treatment procedure, which includes aeration, coagulation, flocculation, sedimentation, filtration, and disinfection. Water quality analysis reveals that the treatment process decreased turgidity and improved a number of physicochemical parameters; however, the concentrations of calcium and magnesium hardness in treated water remained above the permissible limits set by the Bureau of Indian Standards (BIS), suggesting the need for improved treatment efficiency and optimized chemical dosing. The total municipal water demand is 4.59 million litres per day (MLD), whereas the available storage capacity is only 2.83 million litres, resulting in a storage deficit of 1.76 million litres per day. As a result, rather than supplying water continuously every day, the municipality only supplies it once every two days. This study also identified that uneven distribution of storage tanks and water connections to respective overhead storage tank, indicating improper planning of water supply infrastructure. The discharge of untreated wastewater generated during the treatment process into a nearby stream

also raises environmental concerns. To improve the effectiveness, equity, and sustainability of the municipal water supply system, the study suggests increasing storage capacity, planning the distribution network scientifically, optimizing treatment operations, conducting regular parameter-based water quality monitoring, and improving wastewater management.

Keywords: Urban Water Supply, Water Treatment Plant, Drinking Water Quality, Municipal Water Management, Water Distribution Network, Storage Capacity, Sustainable Water Supply

Introduction

Water is the most significant renewable natural resource on the earth and significantly impacts a nation's socio-economic development. Water is used for drinking, household needs, and irrigation (Mundhe et al.,2023; P.M.C., 2012). Thus, the progress of any country depends on water availability and easy access. The water sources for water supply are rainfall, surface water, and groundwater. The rate of world urbanization is increasing rapidly from 1950 to 2020. The world population in cities has increased from 29.6 percent to 56.2 percent and is predicted to reach 68.4 percent by 2050 ((U.N.), 2018). Population and urbanization have increased demand of water for domestic, industrial, and agricultural use (Kumar, 2018). Thus, the per capita water availability decreased, and the groundwater level declined, creating several water crises (Singh & Singh, 2002). The Shrigonda town comes under the rain shadow zone of Maharashtra; the rainfall is erratic, leading to a water scarcity problem. The available groundwater resources are polluted, and most sites exceed water quality standards. Water security is affected by water scarcity, which occurs when water demand exceeds supply; in dry periods, supply decreases and demand increases due to drying up surface water bodies and declining groundwater yield. Water scarcity immediately impacts economic growth, urban environmental quality, and the health and well-being of city dwellers (McDonald et al., 2011; Chunyang et al., 2021). There are three components to providing fresh water to city residents: water availability, water quality, and water supply system (McDonald et al., 2011). With the rapidly increasing population, modern sanitary facilities increase, which leads to per capita water use. Generally, 80 percent of domestic water use converts to waste (Punmia & Jain, 2005). This wastewater deteriorates groundwater quality due to discharge around the town; the municipal water supply is based on surface water (Ghod Dam), and the amount of water in the reservoir decreases in the dry season. Thus, the municipality curtails water supply in the dry season. Insufficient water supply and groundwater depletion in quantity and quality create water scarcity in the dry season. Therefore, the present chapter focuses on water filtration mechanism, supply system and water quality to suggest alternatives to minimize the water stress situation in Shrigonda.

Methodology

To evaluate the water supply system of the municipality, information about water supply viz frequency of water supply, amount of water supply, quality control, human resources, number of overhead tanks and their storage capacity, connections under each overhead tank, ward wise number of water supply connections and growth of water supply connections, etc. are acquired from water supply department, municipality office. A GPS survey was carried out to map overhead storage tank locations and water sampling sites (tap water) for quality control. The process of water treatment is evaluated with a field visit to W.T.P. The entire water treatment process was monitored with pump operator (Dam water entry to filtered water exit) for the whole day. Both treated and untreated Water samples were taken and tested in the laboratory to determine the water quality.

Municipal Water Supply System

Shrigonda Municipality constructed a water treatment plant in 2018 under the Urban Infrastructure Development Scheme for Small and Medium Towns by investing 4595.63 lakhs (S.M.C., 2017) to supply safe drinking water to the dwellers. The W.T.P.s filtration capacity is 1.10 cr Litres/per/day (11 M.L.D.). The water is lifted from Ghod Dam and transferred to W.T.P.s built near Shrigonda town for treatment. The treated water is then distributed and stored in 7 overhead water storage tanks constructed around the different parts of the city. The zone-wise water distribution network is connected to the storage tank. Eleven water supply zones have been prepared for effective and efficient water supply. In each zone, one whaleman is appointed on a contract basis to release water to the individual connection. Single-time water is supplied in two days because the water supply system cannot provide daily water to the whole town.

Water Treatment Process

The water treatment plant constructed at Shrigonda follows a series of water treatment steps: Aeration, Coagulation, Flocculation, Sedimentation, Filtration, and Disinfection.

Aeration

This process brings air and water close to increasing the amount of oxygen in the water. While reducing dissolved gases such as ammonia, chlorine, carbon dioxide, hydrogen sulfide, methane, iron, and manganese. (Kokila & R.,2015)

In the removal of these contaminants in later solid-liquid separation processes, coagulation is a process that binds small particles to large aggregates (flakes) and absorbs the dissolved organic matter into the particulate aggregates. Chemical coagulants such as potassium aluminum sulfate (alum) Kal $(SO_4)_2 \cdot 12H_2O$ are frequently used before the flocculation tank and after the aerator is the coagulant. (Carlos et al., 2018).

- **Flocculation:** Flocculation involves adding a chemical coagulant to water to encourage the bonding of particles, resulting in larger aggregates that are easier to separate. Shrigonda town's W.T.P.s treat drinking water using this method of treatment. (Carlos et al., 2018).

Clariflocculation Process

The two concentric tanks that make up the Clariflocculation Tank are used as clarifiers and flocculation basins, respectively. Coagulation, flocculation, and sedimentation processes are happening in the Clariflocculation tank. After the aerator fountain in the small tank, Chemicals (coagulants) Aluminum sulfate (alum) (50 kg per day) and bleaching powder (Calcium chloride) (12 kg per day) were randomly added to the water in the forms of solution. At the bottom of the clariflocculator tank, suspended solid particles settle down.

- **Disinfection:** To kill parasites, bacteria, and viruses, bleaching powder (Calcium chloride) (12 kg per day) was randomly added to water after sedimentation.

Sedimentation

It is a common type of water treatment. It is the process of removing solids that float and settle in water. This process depends on the use of settling tanks to remove larger solids. After clarification, water is passed through a sand bed with six layers of different sizes and then transferred to pump house to distribute in other tanks.

- **Distribution:** Treated water with all the above processes is distributed in the town.
- **Cleaning Mechanism:** The sand floor (filter) is cleaned twice weekly to remove sediment. The settling basin is cleaned once a month.
- **Waste Water Management:** The wastewater generated after the treatment is discharged directly to the nearby stream without any treatment.

Municipality Supplied Water Quality

Both treated and untreated Water samples were taken and tested in the laboratory to determine the water quality. Table 1 shows the results of both samples. Twelve physicochemical parameters were evaluated and compared to the limits condemned by BIS. Four input water parameters (total hardness (as CaCO3), magnesium hardness, Ca2 +, Mg2 +) exceeded the BIS limit, and two treated water parameters (magnesium hardness, Ca2 +) exceeded the BIS limit. Municipal W.T.P. efficiency does not meet the BIS standard of drinking water quality. The amount of potassium aluminum sulfate (alum) and calcium chloride is randomly added to the water.

Table.1 Quality of Municipality supplied water

Sr. No.	Parameters	Units	BIS Limits	Before Filter	After Filter
1	Colour	----		Clear	
2	Odour	----		Non-Objectionable	
3	Taste			Non-Objectionable	
4	pH		6.5 to 8.5	7.12	7.65
5	Electrical Conductivity	µs/ppm	300	249.4	255.8
6	Turbidity	NTU	5	6.2	4.7
7	Chloride	mg/l	250	226.56	127.44
8	Total Hardness (as $CaCO_3$)	mg/l	300	350	280
9	Calcium Hardness	mg/l	200	134.57	100.93
10	Magnesium Hardness	mg/l	100	215.43	179.07
11	Ca^{2+}	mg/l	75	336.00	252.00
12	Mg^{2+}	mg/l	30	52.37	43.53
13	Total Dissolved Solids	mg/l	500	210	178.1
14	Total Suspended solids	mg/l	500	63.5	39.4
15	Total Solids	mg/l	500	273.5	217.5
16	Phosphate			26	23
17	Sulphate	Mg/l	200	58	18
18	MPN	MPN/100 ml	0/100 ml	2	0

(Source: Computed by Researcher)

Water Quality Control Mechanism

The municipality's water supply department collects 20 monthly water samples from tap water in various parts of the town. The collected water samples are sent to Ahmednagar District Health Laboratory for testing. The laboratory returns the results as drinkable or non-drinkable. Water parameter-wise reports were not available with the municipality. If the sample turns out to be unfit for consumption, then the sample will be recollected and tested.




Fig.1 Water sampling sites of municipality for water quality control

Water Storage Tanks with Capacity




Fig.2: Treated Waters Storage Overhead Tank

Table 2: Water storage tanks with capacity and tap connections

Sr. No	Location of Storage Tank	Year							
		2021		2019		2011		2001	
		Total Storage Capacity (Litre)	Number of Water Supply Connection Under Tank	Total Storage Capacity (Litre)	Number of Water Supply Connection Under Tank	Total Storage Capacity (Litre)	Number of Water Supply Connection Under Tank	Total Storage Capacity (Litre)	Number of Water Supply Connection Under Tank
1	I.T.I. Colony	900000	2352	900000	2176	Not Constructed			
2	W.T.P.s, Anandkar Mala	950000	302	950000	222				
3	Dattawadi	200000	16	200000	14				
4	Bajartal	200000	530	200000	479	200000	708	200000	530
5	Shani Chouk	445000	0	445000	0	445000	868	445000	650
6	Daund Road	500000	650	500000	588	500000	339	500000	254
7	Sidhharth Nagar	80000	352	80000	318	80000	469	80000	351
Total		**32,75,000**	**4,202**	**32,75,000**	**3,798**	**12,25,000**	**2,385**	**12,25,000**	**1,785**

(Source: Water Supply Reports of Shrigonda Municipality)

W.T.P.'s filtration capacity, water requirement and storage capacity

W.T.P.'s filtration capacity	Water Requirement (Litres@113/LPCD)	Total storage capacity (Litres)
1.10cr/Litre/Day 11 MLD	45,93,804	28,30,000

(Source: Water Supply Reports of Shrigonda Municipality)

The Shrigonda municipality has constructed seven overhead water storage tanks with different water storage capacities. The storage capacity of overhead water storage tank is from 200000 to 950000 litres. The town's total water requirement is 45,93,804 litres per day, but the total water storage capacity of the overhead storage tank is 28,30,000 litres. The storage capacity of the overhead tank is shorter by 17,63,804 litres per day; due to this gap between storage and requirement, the municipality cannot supply water daily. Three overhead water storage tanks were augmented in 2019 viz I.T.I. Colony (900000 litres), W.T.P. Anandkar mala (950000 litres), and Datta wadi (200000 litres). After the construction of these tanks, tap connections were disconnected from the Shani Chouk water tank; now, it

is operational only for tanker water supply. The Datta Wadi tank has only 16 connections, and the Bajartal tank has 530 connections with the same storage capacity (200000 litres). It seems that tank requirements are not considered during construction. The Daund Road storage tank has 650 connections, and the Siddhartha Nagar storage tank has 352 connections. The Anadkar Mala storage tank has only 302 connections, while the I.T.I. colony storage tank has 2352 connections, and both tanks have almost 900000 litres above storage capacity. The data on storage tanks and the number of water supply connections under each tank clearly show that storage tanks are not constructed as per the demand for water supply connections. The construction of storage tanks has been done randomly, so there is a gap between demand and supply of water.

Table: 3 Ward-wise number of water supply connections

Wards	**2001**	**2011**	**2019**	**2021**
1	42	56	88	97
2	258	345	550	609
3	108	144	230	254
4	282	376	599	663
5	09	13	20	22
6	384	512	816	903
7	26	35	56	62
8	300	401	639	707
9	376	501	800	885
Total	**1785**	**2385**	**3798**	**4202**

(Source: Water Supply Reports of Shrigonda Municipality, 2022)

Table 3 demonstrates ward-wise water supply connections from 2001 to 2021. The number of water supply connections has increased in all wards. Wards 1, 5, and 7 have few connections 97, 22, and 62 because the municipality has not installed a pipeline in most of areas. The people from this area mainly depend on groundwater for their domestic needs

Human Resources for Water Supply

Table: 4 Human Resources for Water Supply

Sr. No	**Water Supply Section**	**Type of work**	**Number of Worker**
1	Water treatment Plant	Pump Operator	09
2	Walveman	Water Supply	08
3	Maintenance	Repairing Leakage	04
4	Office	Administration	02

(Source: Water Supply Reports of Shrigonda Municipality)

Conclusion

The present study concludes that Shrigonda Municipality has developed a functional urban water supply system with an 11 MLD water treatment plant and seven overhead storage tanks that play a vital role in supplying treated drinking water to the town. The treatment process effectively improves water quality by reducing turbidity and microbial contamination; however, calcium and magnesium concentrations in treated water remain above the permissible limits prescribed by the Bureau of Indian Standards (BIS), indicating the need for optimization of treatment processes and chemical dosing. Although the treatment plant has adequate filtration capacity, the existing storage capacity is insufficient to meet the town's daily water demand, resulting in a storage deficit of approximately 1.76 MLD and an intermittent water supply schedule of once every two days. Furthermore, the uneven distribution of overhead storage tanks and water supply connections reflects inadequate planning of the distribution network, leading to inequitable water supply across different parts of the municipality. The discharge of untreated wastewater from the treatment plant into a nearby stream also poses environmental concerns. Therefore, increasing storage capacity, scientifically redesigning the distribution network, implementing parameter-based water quality monitoring, optimizing treatment efficiency, and adopting appropriate wastewater management practices are essential to ensure a reliable, equitable, and sustainable urban water supply system for Shrigonda Municipality.

References

1. Athavale, R. N. (2003). Water Harvesting and Sustainable Supply in India. Centre for Environment Education, Ahmedabad: Rawat Publications Jaipur and New Delhi.
2. BIS. (1993,2002). Code of Basic Requirements for Water Supply, Drainage, and Sanitation. New Delhi: Burea of Indian Standerd.
3. Carlos Alberto, Martínez-Huitle, & Manuel Andrés Rodr (Eds.). (2018). Electrochemical Water and Wastewater Treatment. Elsevier. Doi: https://doi.org/10.1016/C2016-0-04297-3.
4. Central Public Work Department. (2002). Rain harvesting and conservation manual. New Delhi: Government of India.
5. CPWD. (2002). Manual Rainwater Harvesting and Conservation. New Delhi: Central Public Work Department.
6. Ghaitidak, D. M., & Yadav, K. D. (2013). Characteristics and treatment of greywater—a review. Environmental Science and Pollution Research, 20, 2795-2809. doi:DOI 10.1007/s11356-013-1533-0.
7. Gould, J., & Nissen, P. E. (1989). Rainwater Catchment Systems for Domestic Supply. International Technology Publications.

8. Government of India. (2003). Water Resources of India and the World. Ministry of Water Resources, Newsletter on fresh water year.
9. Jefferson, B., Palmer, A., Jeffrey, P., Stuetz, R., & Judd, S. (2004). Grey Water characterization and its impact on the selection and operation of technologies for urban reuse. Water Science and Technology, 50(2), 157-164.
10. Kokila, K., & R., D. (2015). Analysis and Design of Cascade Aerator Construction for Mettur. National Conference on Research Advances in Communication, Computation, Electrical Science and, (pp. 42-46).
11. Kumar, C. P. (2018). Water Resources Issues and Management in India. Journal of Scientific and Engineering Research, 5(9), 137-147.
12. Mali, S. P., & Pawar-Patil, V. S. (2013). Potential Roof Rain Water Harvesting in Pirwadi Village of Kolhapur District, Maharashtra (India) - A Geospatial Approach. Journal of Research in Humanities and Social Science, 1(4), 19-23.
13. McDonald, R. I., Green, P., Balkc, D., Fekete, B. M., Revenga, C., & Todd, M. (2011). Urban growth, climate change, and freshwater availability. PNAS, 108(15), 6312-6317. Doi: https://doi.org/10.1073/pnas.1011615108.
14. Metcalf, Eddy, & Inc. (2012). Wastewater Engineering Treatment and Reuse. Tata McGraw Hill Private Limited.
15. Mohd, S., Athar, H., & Gauhar, M. (2016). Analysis of groundwater quality using water quality index: A case study of greater Noida (Region), Uttar Pradesh (U.P), India. Cogent Engineering, 3, 1-11.
16. Mundhe, N., Gaikwad, S., Thakare, S., & Ugale, V. (2023). Assessment of Domestic Water Resources for Sustainable Utilization Using Geospatial Techniques. Journal of Settlements and Spatial Planning, 14(2), 53-65.
17. NWM. (2023). National Water Mission. Ministry of Jalshakati, Department of Water Resources. Retrieved from http://nwm.gov.in.
18. Pacey, A., & Adrian, C. (1989). Rainwater Harvesting: The collection of rainfall and runoff in Rural areas. Intermediate Technology Publications, 55.
19. PMC. (2012). Revising/Updating the City Development Plan (C.D.P.) of Pune City-2041 under JNNURM. Retrieved from https://pmc.gov.in/sites/default/files/project-glimpses/Draft_City_Development_Plan_for_Pune_City_2041_Vol-1.pdf.
20. Punmia, B. C., & Jain, A. K. (2005). Waste Water Engineering. New Delhi: Laxmi Publications(P) Ltd.
21. Ranade, R. (2000). A Water Harvesting Manual for Urban Areas: Case Studies from Delhi. New Delhi: Publication Center for Science and Environment.
22. Ronalton, E. M., Tais, O. C., & Matheus, H. M. (2022). Determination of runoff coefficient (C) in catchments based on precipitation and flow events analysis. International Soil and Water Conservation Research, 208-216.

23. Singh, D. K., & Singh, A. K. (2002). Groundwater Situation in India: Problems and Perspective. International Journal of Water Resources Development, 18(4), 563-580. doi:DOI: 10.1080/ 0790062022000017400.
24. SMC. (2012). Annual Report, Shrigonda Municipality.
25. Tariq, J., & Shahrour, I. (2021). Rainwater Harvesting to Address Current and Forecasted Domestic Water Scarcity: Application to Arid and Semi-Arid Areas. Water, 13(3583), 1-16. doi: https://doi.org/10.3390/w13243583.
26. (U.N.), U. N. (2018). 2018 Revision of World Urbanization Prospects. Retrieved from https://population.un.org/wup/
27. WHO. (1993). Guidelines for drinking-water quality 1: recommendations, 2nd ed (2 ed., Vol. 1). World Health Organization. Retrieved from

Advances in Geography and Environmental Research

ISBN: 978-93-49938-93-9| Year: 2026 | pp: 71 - 77

Honest Implementation of the Environment Impact Assessment Notification 2006: Roadway Towards Sustainable Development

Dr. Umesh S. Aswar

Former Judge, Associate Professor (Law), Government Law College, 'A' Road Churchgate Mumbai 400020

Email: umeshaswar@gmail.com

Article DOI Link: https://zenodo.org/uploads/21476494

DOI: 10.5281/zenodo.21476494

Abstract

The Right to a pollution-free clean environment is a basic collective human right. The same has been assured as an integral part of the right to life and personal liberty under Article 21 of the Indian Constitution. At the same time, the right to life also includes the collective right to development, which can be materialised by the use of natural resources, resulting in certain levels of pollution. Therefore, there is a need to maintain a balance between the above-mentioned rights through law by implementing the concept of sustainable development. Environmental Impact assessment notifications are vital regarding it.

Keywords: right to pollution-free clean environment – right to development – concept of sustainable development – environment impact assessment notifications.

Statement of Problem

As mentioned above, the right to a pollution-free environment and the right to development both are necessary to be implemented for the fullest realisation of the right to life specified under Article 21 of the Constitution. This objective can be achieved by maintaining a balance between these two rights, which are contrary to each other. The means to achieve this end is the implementation of the concept of sustainable development by enacting suitable legislation/rules, etc. Accordingly, the environmental impact assessment notifications have been framed. Now the question is how far these notifications can maintain a balance between the right to a pollution-free clean environment and the right to development?

Introduction

India is a country governed by the rule of law. The Indian Constitution secures the implementation of the rule of law; wherein human rights have been assured to the citizens as well as noncitizens. International environmental jurisprudence includes the mandate to provide environmental rights to the people at large. It has been laid down mainly in the form of declarations/resolutions adopted in Earth summits regularly organized by the United Nations Organization after regular intervals of every ten years since 1972. India is a signatory to all Earth Summits; accordingly, is duty-bound to implement human rights, which include the right to a pollution-free clean environment as well as the right to development in favour of individuals. These are collective human rights because they are possible to be enjoyed in a group. The implementation of these rights, which are contrary to each other, has been secured by the implementation of the concept of sustainable development through the Environment Impact Assessment Notifications 1994 and 2006. The concept of sustainable development warrants the existence of a state of affairs where industrialisation will be happening in such a manner, causing minimum harm to the environment and making good of the loss caused to it by creating more environmental resources like an increase of forests, trees and green covers.

Environment Impact Assessment Notification 2006

National Environment Policy is approved by the Union Cabinet on 18th May 2006 and accordingly the procedure is specified in the 2006's notification that at the central Government or the state or union territory level environment impact assessment authority (SEIAA), will be constituted by the central Government in consultation with the state Government or the union territory administration concerned under sub-section (3) of section 3 of the Environment (Protection) Act 1986.

Central Government accordingly directed that on and from the date of its publication (of notification) the required construction of new projects or activities or the expansion or modernization of existing projects or activities listed in the Schedule to this notification entailing capacity addition with change in process and or technology shall be undertaken in any part of India only after the prior environment clearance from the central Government or as the case may be, by the state level environment impact assessment authority, duly constituted by the central Government.

Following projects or activities shall require prior environment clearance from the concerned regulatory authority, which shall hereinafter referred to be as the Central Government in the Ministry of Environment and Forests for matters falling under Category 'A' in the Schedule and at State level the State Environment Impact Assessment Authority (SEIAA) for matters falling under Category 'B' in the said

Schedule, before any construction work, or preparation of land by the project management except for securing the land, is started on the project or activity:

- All new projects or activities listed in the Schedule to this notification;
- Expansion and modernization of existing projects or activities listed in the Schedule to this notification with addition of capacity beyond the limits specified for the concerned sector, that is, projects or activities which cross the threshold limits given in the Schedule, after expansion or modernization;
- Any product change - mix in an existing manufacturing unit included in Schedule beyond the specified range.

All projects and activities are broadly categorised into two categories: Category A and Category B, based on the spatial extent of potential impacts and potential impacts on human health and natural and manmade resources.

Screening, Scoping and Appraisal Committees

The same Expert Appraisal Committees (EACs) at the Central Government and SEACs (hereinafter referred to as the EAC and SEAC) at the State or the Union territory level shall screen, scope and appraise projects or activities in Category 'A' and Category 'B' respectively.

The environment clearance process for new projects will comprise a maximum of four stages, all of which may not apply to particular cases, as set forth below in this notification. These four stages in sequential order are: stage one Screening (only for Category 'B' projects and activities), stage two Scoping, stage three Public Consultation and stage four Appraisal.

Screening: In case of Category 'B' projects or activities, the scrutiny of an application seeking prior environment clearance made in Form 1 is done by the concerned State-level Expert Appraisal Committee (SEAC). The projects requiring an Environmental Impact Assessment report shall be termed Category 'B1', and the remaining projects shall be termed Category 'B2' and will not require an Environmental Impact Assessment report.

Scoping: It refers to the process by which the Expert Appraisal Committee in case of Category 'A' projects or activities, and State level Expert Appraisal Committee in the case of Category 'B1' projects or activities, including applications for expansion and/or modernization and/or change in product, mix of existing projects or activities, determine detailed and comprehensive Terms of Reference (TOR) addressing all relevant environmental concerns for the preparation of an Environment Impact Assessment (EIA) Report.

Public Consultation: It refers to the process by which the concerns of local affected persons and others who have a plausible stake in the environmental impacts of the project or activity are ascertained with a view to taking into account all the

material concerns in the project or activity, as deemed appropriate. All Category 'A' and Category 'B1' projects or activities shall undertake Public Consultation, except the following;

- Modernisation of irrigation projects.
- All projects or activities located within industrial estates or parks approved by the concerned authorities, and which are not disallowed in such approvals.
- Expansion of Roads and Highways, which do not involve any further acquisition of land.
- All Building / Construction projects / Area Development projects, and Townships.
- All Category 'B2' projects and activities.
- All projects or activities concerning national defence and security or involving other strategic considerations as determined by the Central Government.

Appraisal means the detailed scrutiny of the application and other documents, like the Final EIA report, outcome of the public consultations, including public hearing proceedings, submitted by the applicant to the regulatory authority concerned for the grant of environmental clearance.

The appraisal of an application shall be completed by the Expert Appraisal Committee or State Level Expert Appraisal Committee concerned within sixty days of the receipt of the final Environment Impact Assessment report and other documents or the receipt of Form 1 and Form 1 A, where public consultation is not necessary and the recommendations of the Expert Appraisal Committee or State Level Expert Appraisal Committee shall be placed before the competent authority for a final decision within the next fifteen days.

The Regulatory Authority shall consider the recommendations of the EAC or SEAC concerned and convey its decision to the applicant within forty five days of the receipt of the recommendations of the Expert Appraisal Committee or State Level Expert Appraisal Committee concerned or in other words within one hundred and five days of the receipt of the final Environment Impact Assessment Report, and where Environment Impact Assessment is not required, within one hundred and five days of the receipt of the complete application with requisite documents.

The prior environmental clearance granted for a project or activity shall be valid for a period of ten years in the case of River Valley projects (item 1(c) of the Schedule), project life as estimated by Expert Appraisal Committee or State Level Expert Appraisal Committee subject to a maximum of thirty years for mining projects and five years in the case of all other projects and activities.

Environmental Impact Assessment Notification 1994

The concept of environmental audit of the proposed development project for the implementation of the concept of sustainable development has been under implementation in the Indian legal system since 1994. The Environment Impact

Assessment Notification 1994 served the cause of sustainable development till they have been superseded by the 2006 notification. Under the 1994 notification, there was a mandatory provision to seek prior environmental clearance for projects having a proposed investment of Rs 50 crores, which was later enhanced to Rs 100 crores. Under this notification, it was the central government, through the Union Ministry of Forests & Environment, that used to do the environmental audit of the schedule one projects. The active role of the state Governments in this regard was missing. Therefore, in order to decentralise the power to conduct environmental impact assessment of the proposed development project between the central and state Governments, the present 2006 notification has been framed. This notification does carry many novel features as discussed above, which were missing in the 1994 notification.

2020's Proposed Amendment in the Environment Impact Assessment Notification 2006

This draft of the amendment aims to dilute the effectiveness of the process of environment impact assessment. With the proposed amendment, the central as well as state Governments could allow the commencement of the developmental project on an urgent basis without seeking the environmental clearance for it at the instance of the competent authorities specified in the 2006 notification. While the developmental projects are in progress, the investors/industrialists may complete the process for seeking ex-post-facto environmental clearance. This provision has been justified under the pretext of the urgency clause and the need to accelerate the implementation of the collective right to development of Indian citizens.

Conclusion

Based on the above-mentioned analysis, the following conclusions are drawn:

- The right to development and right to pollution-free clean environment are the collective human rights which are an integral part of the right to life under Article 21 of the Constitution. These rights are contrary to each other because development will always be achieved at the cost of the use of natural resources, causing a certain degree of environmental pollution. Hence, it is necessary to maintain a balance between these rights through the implementation of the concept of sustainable development.
- The Environment Impact Assessment Notification 2006 is instrumental in maintaining the above-mentioned balance between the right to development and the right to a pollution-free environment. Under this notification, the competent authority has to decide on the environmental clearance of the proposed developmental projects based on merit within the stipulated time period only. Non-compliance with the relevant stages or otherwise, reasons like delay / lethargic approach resulting in failure to decide by the competent authority/ Government, amounts to an extension of deemed environmental clearance to

the said project proposals of an industrialist. It shows the balance maintained under the 2006 notification, wherein the applicant industrialist, in the proposal of the project, has to design the future execution of the industrial activity in such a manner as to cause no or minimum hazard to the environment while ensuring future benefits to the people at large through the proposed industry. At the same time, competent authorities and Governments are duty-bound to take decisions on the environmental clearance of the projects within the prescribed time so that developmental projects generating employment, revenue and material wealth will not be stalled. In this manner, the doctrine of sustainable development is operative in the provisions of the Environment Impact Assessment Notification 2006. It, on one hand, ensures the environmental clearance to the developmental projects and on the other hand takes care that no or minimum hazard will be caused to the environment so that the right to a pollution-free clean environment and the right to development under Article 21 will be harmonised.

- While the Environment Impact Assessment Notification 2006 is in force, the central as well as state Governments can exercise their upper hand directly or indirectly while granting environmental clearance to the development projects. This influence is operative in a certain manner, for example, canvassing the policy of development projects through print, electronic media and on public platforms, underlining its significance in the context of social and material development of society. Similarly, if environmental clearance is challenged before the judiciary by public-spirited people, then in these litigations, the state can justify its stand regarding the utility of the projects as mentioned above. Moreover, Governments can shape the opinion of the members of the competent authorities positively under 2006's notification in an indirect manner. Even after having 2006 notifications, governments have implemented the concept of sustainable development to a satisfactory extent.
- The 2020s proposed amendment to the environment impact assessment notification will be diluting the concept of sustainable development; arbitrarily leaning in favour of the right to development and discarding the right to a pollution-free, clean environment of the people at large. Therefore, it is needless to state that these proposed amendments are necessary to be withdrawn by the state; if not, the judiciary should strike them down.

References

1. P. Leelakrishnan, Environmental Law in India, 2019, Lexis Nexis.
2. Divan Shyam and Rosencranz Armin, Environmental Law and Policy in India: Cases, Materials& Statutes.
3. H. N. Tiwari, Environmental Law, 2016, Allahabad Law Agency.
4. S. Shanthakumar, Introduction to Environmental Law, 2017, Lexis Nexis.

5. Dr S. C. Tripathi, Environmental Law, 2018, Central Law Publication.
6. M. P. Jain, Indian Constitutional Law, 2018, Lexis Nexis.
7. Dr Paramjit Jaiswal, Environmental Law, 5th Edition 2021, Allahabad Law Agency.
8. Benny Joseph, Environment Studies, 2nd edition 2008, Tata McGraw-Hill, New Delhi.
9. Shantakumar's Introduction to Environmental Law, 2nd edition (2005), Wadhwa and Company, Nagpur.
10. Gurdip Singh, Environmental Law in India, 2005, MacMillan Publishers, New Delhi.

Advances in Geography and Environmental Research

ISBN: 978-93-49938-93-9| Year: 2026 | pp: 78 - 81

Literacy of Rural Settlement in Daund Tahsil, Pune district (MH): An Analytical Study

Dr. Parkhe Sakharam Baban

Assistant Professor, Department of Geography, Shri Chhtrapati Shivaji Mahavidyalaya, Shrigonda

Email:

Article DOI Link: https://zenodo.org/uploads/21494604

DOI: 10.5281/zenodo.21494604

Abstract

Daund tahsil displays the general effect of River System, Rail line, Irrigation facilities and Road network so the rural settlement is one of the focal matters of affect in this section. It could be extremely relevant to see its impact on the nature and rural population. Generally, there are dual sets of constraint viz. sensible and ethnic which affect settlement in rural zones. Though the natural factor is influential in rural settlement but the ethnic factor also shows an associatory character. The Daund tahsil dominantly contained of six circles viz. Rahu, Yawat, Kedgaon, Patas, Daund and Ravangaon consists taking 103 revenue villages. Daund Tahsil covers 103 villages and one urban center, occupying about 1288 square kilometers of geographical area with a supporting 382535 of the population in 2011.

Keywords: Rural Population, Literacy, Male and Female Literacy.

Introduction

Daund tahsil is dominantly encompassed of six circles viz. Rahu, Yawat, Kedgaon, Patas, Daund and Ravangaon The total population of the Daund tahsil was 3,82,535. Out of these, 1,98,269 (51.83%) males and the remaining 1,84,266 (48.17%) females as per Census 2011. At the time, 3,26,099 (85.24%) population was rural, and 56,436 (14.76%) belonged to urban areas. Several factors control population distribution, but predominant factors are irrigation facilities, agricultural development, industrial development, transport network, and urbanization in Daund tahsil.

Objective of the Study


- To study the circle wise rural literacy.
- To study the Male and Female Literacy.

Methodology of the Study

The present study is based on 2 kinds of assessment: one on realistic valuation and the other on a real valuation i.e. Physical-cultural-socio-economic characteristic of the region. Primary data of this look into has been complete available from exhaustive field survey of the study zone and secondary data has been completed available from available sources akin to Census handbooks of Pune district Maharashtra. Total 26 hamlets have been needed as the sample from the study region by random sampling method.

Literacy


The general literacy rate of Daund Tahsil was 67.98 per cent, while female literacy was only 43.93 per cent and 56.07 per cent of males as per Census 2011. The circle-wise spatial distribution of literacy showed that the highest literacy was recorded in Patas Circle, which was 68.84, followed by Daund Circle (68.14%), Ravangaon Circle (68.11%), Rahu Circle (66.56%) due to mainly establishment of new schools and colleges in the study area. At the same time, the lowest literacy recorded by Kedgaon Circle was only 58.66 per cent, which was very low compared to the Maharashtra State literacy rate (83.34%) because of a lack of awareness of education.




In the Daund tahsil, females were behind males in terms of literacy. In the 2011 Census, the general literacy of the study area was 67.98 per cent, as accepted that male literacy was higher (56.07%) than female literacy (43.93%). The male literacy ranges between 55.15 per cent to 57.07 per cent. The highest male literacy has been observed in Rahu circle, which was 57.07 per cent. Followed by Ravangaon circle (56.48%), Yawat circle (56.21%), Daund circle (56.16%), and Patas circle

(56.09%). On the other hand, the lowest male literacy was recorded in the Kedgaon circle (55.15%).

At the same time, female literacy is still problematic because all circles indicated lower female literacy than male literacy. The lowest female literacy was recorded in Rahu circle, only 42.43 per cent, half of the state average (78.87%). Followed by Ravangaon circle (43.52%), Yawat circle (43.79%), Daund circle (43.84%), Patas circle (43.91%), and Kedgaon circle (44.85%). Several studies pointed out that improving female literacy uplifts the per capita income and national income. An increase in female literacy is strongly responsible for narrowing the regional imbalance. Therefore, it is essential to carry out a systematic program to increase female literacy up to the state average.





Result and Discussion

The disparity in male-female literacy in Daund tahsil was 12.14 per cent. However, the wide gap in male-female literacy found in the Rahu circle (14.14%) is alarming for the study area. In the remaining circles, including Ravangaon (12.96%), Yawat (12.42%), Daund (12.32%), Patas (12.18%), Kedgaon (10.30%), the gap between male and female literary was above 10 per cent. Such circles in the study area indicated poor performance in the female education field. Thus, there is an urgent need to concentrate more on educating female children, which will help reduce the disparity in male-female literacy in the study area.

References

1. Ahmad. E. (1954), Geographical Essays on India, Patna, P.33-34.
2. Chandana R.C. and Sidhu M.S. (1980), Introduction to population geography, Kalyani Publication, New Delhi, p.3.
3. Census of India (2011), "District Census Handbook of Pune", Published by the Director of Census Operations Govt. Printing Press Mumbai PP 18 -203.
4. Employment and Population Growth." Rev. Regional. Stud. 28, pp. 82–101.
5. Survey of India, Indian Topographical Map Toposheet No.47J/2, 47J/3, 47J/6, 47J/7, 47J/10, 47J/11, 47J/15.
6. Socio-economic Abstract of Pune district, 2005-06, 2013-14, 2017-18.
7. Husain, M. (2006). Human Geography. Jaipur: Rawat Publication.
8. Mandal R.B. (1978). Introduction to Rural Settlements. New Delhi: Concept Publishing Company.
9. Richardson, H.W. (1973). Resource Mobility in the Space Economy, London: Macmillan Press Ltd.

Advances in Geography and Environmental Research

ISBN: 978-93-49938-93-9| Year: 2026 | pp: 82 - 92

Spatial Analysis of Groundwater Availability and Quality in India: A GIS-Based Assessment

[1]Mr. Agastirishi Toradmal

[2]Dr. Vilas Vasant Patil

[1]Assistant Professor, Department of Geography, Dada Patil Mahavidyalaya, Karjat, Ahilyanagar

[2]Principal, Shri Shahu Mandir Mahavidyalaya, Parvati, Pune.

Email: agasti0809@gmail.com

Article DOI Link: https://zenodo.org/uploads/21536407

DOI: 10.5281/zenodo.21536407

Abstract

In India, groundwater is an essential supply of water for industrial operations, home use, irrigation, and ecosystem preservation. However, substantial regional disparities in groundwater supply and quality have been brought about by fast population increase, intensifying agriculture, urbanization, shifting rainfall patterns, and excessive groundwater use. The current work uses published groundwater assessments, geographical information system concepts, and secondary sources to investigate the spatial distribution of groundwater availability and quality throughout India. The availability of groundwater was assessed using yearly recharge, extractable groundwater resources, annual extraction, and the groundwater extraction stage. Compliance with drinking water regulations and the spatial distribution of nitrate, fluoride, electrical conductivity, arsenic, iron, manganese, and uranium were used to evaluate groundwater quality.

According to the 2024 national assessment, there will be 446.90 billion cubic meters of groundwater recharge per year, 406.19 billion cubic meters of extractable groundwater resources, and 245.64 billion cubic meters of extraction per year. The extraction stage at the national level was 60.47%. 73.4% of the 6,746 evaluation units were deemed safe, whilst 11.1% were overused. 71.7% of samples met allowable drinking-water standards, while 28.3% exceeded the limit for at least one indicator, according to groundwater quality monitoring at 14,978 stations. The most common contaminant was nitrate, which was followed by salinity and fluoride. The study shows that groundwater quality and availability are regionally heterogeneous

and need to be controlled using GIS-based decision-support systems, demand control, pollution avoidance, artificial recharge, and aquifer-specific monitoring.

Keywords: Groundwater availability, groundwater quality, GIS, spatial analysis, water quality index, groundwater extraction, India

Introduction

Water held beneath the surface of the Earth in soil and rock formations' pores, fissures, and openings is referred to as groundwater. Aquifers are geological formations that have the capacity to store and transfer usable amounts of groundwater. Rainfall infiltration, seepage from rivers, lakes, and canals, irrigation return flow, and artificial recharge structures are the main ways that groundwater is restored. Rainfall, geology, geomorphology, soil attributes, slope, land use, drainage density, and the hydraulic characteristics of aquifers all affect its availability.

Groundwater is essential to India's industry, irrigated agriculture, rural lives, urban water supply, and drinking water security. Groundwater is especially crucial in places where surface water supplies are sporadic or inadequate due to its dispersed occurrence and comparatively modest infrastructure requirements. Groundwater development hasn't, however, happened consistently. Groundwater is often extracted more quickly than it can be naturally supplied by intensive agricultural areas, growing towns, and industrial hubs. Some regions may have significant groundwater resources, but their water quality is poor due to either human activity or naturally occurring minerals.

Groundwater quality and availability are intimately related. Excessive levels of fluoride, arsenic, nitrate, salinity, or trace metals can make water from a high-yielding aquifer unfit for human consumption. On the other hand, an aquifer with high-quality water may not be able to store or recharge as much. As a result, resource planning focused solely on water quantity may result in development that is dangerous or unsustainable.

An efficient framework for combining groundwater levels, recharge, extraction, geology, land use, water quality monitoring, and administrative borders is offered by geographic information systems. While water quality indices and geostatistical interpolation approaches have been used to detect contamination hotspots, GIS-based weighted-overlay techniques have been frequently employed to define groundwater-potential zones. Analytical Hierarchy Process-based groundwater-potential mapping, Water Quality Index evaluation, regular kriging, and multivariate statistical analysis have all been shown to be beneficial in Indian studies.

The present paper provides a national-level spatial assessment of groundwater availability and quality in India. It integrates the 2024 Dynamic Ground Water Resource Assessment, groundwater-quality observations collected during 2024 and evidence from published GIS-based groundwater studies. Because raw local

sampling data were not generated for this paper, the results represent a secondary-data spatial synthesis rather than a new field survey.

Study Area

India extends from the Himalayan Mountain system in the north to the coastal plains and peninsular plateau in the south. Its groundwater conditions vary considerably because of differences in geology, relief, rainfall, drainage, soil and land-use practices.

The massive alluvial aquifers that dominate the Indo-Gangetic-Brahmaputra plains often have significant storage and transmission capacity. On the other hand, a significant area of the Indian peninsula is made up of cemented rocks such as granite, gneiss, and basalt, where groundwater is mostly found in worn layers, joints, and fissures. Many sedimentary basins have limestone and sandstone aquifers, but coastal aquifers are susceptible to seawater intrusion and salinity.

Alluvium, basalt, granite, gneiss, sandstone, limestone, schist, shale, and complex crystalline formations are among India's primary aquifers. Both the availability of groundwater and the mobilization of pollutants including fluoride, uranium, iron, and arsenic are influenced by their mineral composition and weathering characteristics.

India experiences significant seasonal changes in recharging due to its monsoon climate. The majority of the yearly recharge takes place during the southwest monsoon, but its efficiency varies depending on the amount of rainfall, slope, vegetation cover, soil permeability, and aquifer storage capacity. Because of this, a location with heavy rainfall may nonetheless have limited groundwater storage in areas where rapid surface runoff is produced by steep slopes or impermeable rocks.

Data and Methodology

1. Data Sources

The study used the following secondary sources:

- Dynamic Ground Water Resources of India, 2024;
- Annual Ground Water Quality Report 2025, based mainly on groundwater samples collected during 2024;
- Central Ground Water Board groundwater-monitoring information;
- Bureau of Indian Standards drinking-water specifications;
- Published GIS, remote-sensing, Water Quality Index and geostatistical groundwater studies.

The 2024 groundwater-resource assessment was jointly conducted by the Central Ground Water Board and state or Union Territory groundwater departments through the India Ground Water Resource Estimation System. The groundwater-quality database included 14,978 baseline monitoring stations. In addition, 5,368 trend

stations were monitored during pre-monsoon and post-monsoon periods to evaluate seasonal changes.

2. Assessment of Groundwater Availability

Groundwater availability was evaluated using four indicators:

- Annual groundwater recharge;
- Annual extractable groundwater resource;
- Annual groundwater extraction;
- Stage of groundwater extraction.

The stage of groundwater extraction was calculated as:

$$\text{Stage of Extraction} = \frac{\text{Annual Groundwater Extraction}}{\text{Annual Extractable Groundwater Resource}} \times 100$$

The evaluation units were categorized as saline, overexploited, critical, semi-critical, and safe. Increased extraction means that groundwater resources are under increased strain and can't keep up with growing demand. The spatial layers of geology, geomorphology, slope, drainage density, lineament density, soil, rainfall, land use, and ground water depth can all serve as the foundation for a thorough GIS analysis at the district level. A weighted overlay model can be used to standardize and integrate these in the following ways:

$$GPI = \sum_{i=1}^{n} wiri$$

where GPI is the Groundwater Potential Index, wi is the normalized weight of each criterion and ri is its standardized suitability rank. AHP or Fuzzy-AHP may be used to determine criterion weights. Published national-scale Fuzzy-AHP analysis has identified aquifer type, drainage density and groundwater depth as particularly influential groundwater-potential factors.

3. Assessment of Groundwater Quality

Groundwater quality was evaluated using pH, electrical conductivity, total dissolved solids, total hardness, calcium, magnesium, sodium, potassium, chloride, sulphate, bicarbonate, nitrate, fluoride and selected trace elements. Parameter concentrations were compared with permissible drinking-water limits under IS 10500.

For local-scale research, individual chemical parameters can be combined into a Water Quality Index:

$$\text{WQI} = \frac{\sum QiWi}{\sum wi}$$

where Qi is the normalized quality rating of parameter i and wi is the weight assigned according to its drinking-water significance. Lower WQI values generally represent better water quality, while higher values indicate poor or unsuitable water. Sampling locations can be plotted using their geographical coordinates. Ordinary kriging or inverse-distance weighting can then be used to generate continuous surfaces for nitrate, fluoride, salinity and WQI. Ordinary kriging is particularly useful because it considers spatial autocorrelation and can estimate prediction uncertainty. WQI-GIS and kriging-based approaches have successfully identified localized groundwater-quality deterioration in several Indian regions.

Results and Discussion

1. National Groundwater Availability

The 2024 assessment estimated total annual groundwater recharge in India at 446.90 billion cubic metres. After reserving water for natural aquifer discharge, the annual extractable groundwater resource was estimated at 406.19 billion cubic metres. Total annual groundwater extraction for domestic, agricultural and industrial uses was 245.64 billion cubic metres, producing a national stage of extraction of 60.47%.

Groundwater indicator	National status, 2024
Annual groundwater recharge	446.90 BCM
Annual extractable resource	406.19 BCM
Annual extraction	245.64 BCM
Stage of extraction	60.47%
Total assessment units	6,746

Although the national extraction percentage appears moderate, the average conceals severe regional inequalities. Of the 6,746 assessment units, 4,951 units, or 73.4%, were classified as safe. A further 711 units were semi-critical, 206 were critical, 751 were over-exploited and 127 were saline. Therefore, approximately one in nine assessment units was already over-exploited.

Groundwater category	Number of units	Percentage
Safe	4,951	73.4%
Semi-critical	711	10.5%
Critical	206	3.05%
Over-exploited	751	11.1%
Saline	127	1.8%

The percentage of safe assessment units increased from 62.6% in 2017 to 73.4% in 2024, while the proportion of over-exploited units declined from 17.24% to 11.13%.

Recharge from tanks, ponds and water-conservation structures increased from 13.98 BCM in 2017 to 25.34 BCM in 2024. These changes indicate that recharge structures and water-conservation programmes may have contributed to improvement, although local groundwater stress remains substantial.

2. Spatial Variation in Groundwater Potential

Groundwater potential is generally higher in thick alluvial aquifers, floodplains and selected coastal zones because these formations possess greater porosity, permeability and storage capacity. A recent national Fuzzy-AHP study estimated that high, moderate and low groundwater-potential zones occupy approximately 27.43%, 42.25% and 30.32% of India's geographical area, respectively. High-potential zones were concentrated in parts of the Indo-Gangetic plains, coastal regions and selected northeastern areas.

However, high natural groundwater potential does not necessarily indicate sustainable use. Several intensively cultivated alluvial regions experience high extraction because of water-demanding crops, dense irrigation-well networks and dependence on groundwater during dry seasons. Thus, areas with productive aquifers may also become critical or over-exploited when withdrawal continuously exceeds recharge.

Hard-rock areas of peninsular India generally display more fragmented groundwater occurrence. Water is stored in weathered zones, fractures and joints rather than in extensive interconnected pore spaces. Groundwater potential can therefore vary significantly over short distances. GIS studies in basaltic terrain have demonstrated that geology, geomorphology, lineament density, drainage density, slope, rainfall and proximity to streams jointly control groundwater occurrence.


Coastal aquifers require special management. Excessive pumping can lower freshwater pressure and permit saline water to move inland or upward into freshwater zones. Arid and semi-arid regions are also susceptible to salinity because limited recharge and high evaporation favour the accumulation of dissolved salts.

3. Groundwater-Quality Status

The groundwater-quality assessment based on 14,978 monitoring stations showed that approximately 71.7% of samples complied with permissible drinking-water limits. The remaining 28.3% exceeded the permissible limit for one or more parameters, indicating that groundwater-quality deterioration is significant but spatially concentrated.

Quality indicator	Percentage of samples
Complying with permissible limits	71.7%
Exceedingly at least one parameter	28.3%
Nitrate above permissible limit	20.7%
Fluoride above permissible limit	8.05%

Electrical conductivity above limit	7.23%



Nitrate

Nitrate was the most widely reported groundwater-quality problem, exceeding the permissible limit of 45 mg/L in approximately 20.7% of samples. Major sources include excessive nitrogen-fertilizer use, leakage from septic tanks and sewage systems, animal waste, unlined waste-disposal areas and irrigation return flow. The broad occurrence of nitrate indicates that groundwater pollution is not solely a geological problem but is strongly associated with agricultural and settlement activities.

Nitrate hotspots are likely to occur in intensively cultivated districts, peri-urban settlements and areas with shallow water tables. Because nitrate is soluble and mobile, it can move rapidly through permeable soils and alluvial aquifers. Preventive management is generally more effective than attempting to remove nitrate after widespread contamination has occurred.

Fluoride

Fluoride concentrations above 1.5 mg/L were observed in approximately 8.05% of samples. Fluoride contamination is predominantly geogenic and is associated with prolonged groundwater-rock interaction in granite, gneiss and other fluoride-bearing crystalline formations. High evaporation, alkaline conditions, low calcium concentrations and long groundwater residence times can further increase fluoride mobility.

Fluoride-affected zones are therefore commonly associated with hard-rock and semi-arid environments. Unlike nitrate, fluoride cannot be controlled only through improved waste management. Alternative water sources, blending, rainwater harvesting and appropriate defluoridation technologies are required.

Salinity and Electrical Conductivity

Electrical conductivity exceeding the relevant limit was found in approximately 7.23% of samples. Higher incidences were reported in arid and semi-arid parts of Rajasthan, Delhi, Haryana and Gujarat and in selected coastal areas. Post-monsoon monitoring showed a modest reduction in high electrical conductivity at several locations, suggesting that rainfall recharge can temporarily dilute dissolved salts. Nevertheless, salinity remained persistent in major hotspot regions.

Salinity can be caused by natural mineral dissolution, evaporation, irrigation return flow, inadequate drainage, seawater intrusion and upward movement of deeper saline groundwater. Continued extraction from saline or vulnerable coastal aquifers can progressively reduce their usability.

Arsenic and Trace Elements

Arsenic contamination is concentrated mainly within parts of the Ganga and Brahmaputra basins. Its occurrence is commonly associated with geochemical processes that release naturally occurring arsenic from aquifer sediments under favourable chemical conditions. Uranium exceedances have been reported sporadically in Rajasthan, Punjab, Haryana, Andhra Pradesh and Telangana. Elevated manganese has been observed in parts of Assam, Karnataka, Odisha, Uttar Pradesh and West Bengal.

The national assessment also showed that iron and manganese were among the more frequently exceeding trace elements. These contaminants require location-specific hydrogeochemical investigation because their occurrence is controlled by aquifer mineralogy, oxidation-reduction conditions, groundwater residence time and local pollution sources.

4. Relationship Between Availability and Quality

The spatial overlay of groundwater availability and quality produces four important management situations:

- **High Availability and Good Quality:** These areas may support development, provided extraction remains below recharge.
- **High Availability but Poor Quality:** Treatment, source substitution or blending is necessary before domestic use.
- **Low Availability but Good Quality:** Extraction must be carefully regulated to protect limited reserves.
- **Low Availability and Poor Quality:** These are priority areas for alternative water supply, recharge and strict demand management.

Over-extraction can indirectly worsen groundwater quality by increasing salinity, drawing contaminated shallow water downward, inducing seawater intrusion or mobilizing poor-quality water from adjoining formations. Similarly, recharge can improve quality through dilution, but contaminated recharge from sewage, fertilizers or industrial waste can deteriorate aquifers. The interaction is therefore controlled by both hydrogeology and human activity.

5. Management Implications

Groundwater-management policies should be spatially differentiated rather than uniformly applied. Over-exploited and critical units require groundwater budgeting, restrictions on new high-capacity wells, crop diversification and improvements in irrigation efficiency. Aquifer recharge should be promoted through check dams, recharge shafts, percolation tanks, restored ponds, rooftop rainwater harvesting and protection of natural recharge areas.

Nitrate management requires balanced fertilizer application, soil testing, improved manure storage, sewerage development and protection zones around drinking-water wells. Fluoride-affected areas require safe-source identification, blending and community-level treatment. Arsenic-affected regions require deeper aquifer evaluation, properly designed wells, regular testing and removal technologies.

A web-based GIS decision-support system should integrate groundwater levels, extraction, recharge, water quality, land use, rainfall and population demand. Hotspot maps should be updated before and after the monsoon. Monitoring density should be increased in industrial areas, coastal belts, intensive agricultural zones and rapidly growing urban settlements. Community participation is also necessary because groundwater abstraction is distributed among millions of individual users.

Conclusion

The spatial analysis demonstrates that groundwater conditions in India are highly heterogeneous. At the national scale, groundwater extraction represents 60.47% of the annual extractable resource, and nearly three-quarters of assessment units are classified as safe. Nevertheless, 751 assessment units are over-exploited, demonstrating that national averages cannot adequately describe regional groundwater stress.

Groundwater quality presents an equally important challenge. Approximately 28.3% of monitored samples exceeded permissible limits for at least one parameter. Nitrate is the most widespread contaminant, reflecting agricultural, sanitation and waste-management pressures. Fluoride, arsenic and selected trace elements are strongly influenced by geological conditions, while salinity affects arid, semi-arid and coastal aquifers.

Sustainable groundwater management must simultaneously consider water quantity and quality. GIS, remote sensing, water quality indices, geostatistical interpolation and aquifer mapping can identify locations where recharge, extraction regulation, water treatment or pollution control should receive priority. Long-term security will depend on continuous monitoring, aquifer-specific planning, efficient irrigation, protected recharge zones, pollution prevention and community-based groundwater budgeting.

References

1. Bureau of Indian Standards. (2012). IS 10500: Drinking water—Specification (Second revision).
2. Central Ground Water Board. (2024). Dynamic Ground Water Resources of India, 2024. Ministry of Jal Shakti, Government of India.
3. Central Ground Water Board. (2025). Annual Ground Water Quality Report 2025. Ministry of Jal Shakti, Government of India.
4. Doke, A. B., Zolekar, R. B., Patel, H., & Das, S. (2021). Geospatial mapping of groundwater potential zones using multi-criteria decision-making AHP approach in a hard-rock basaltic terrain in India. Ecological Indicators, 127, 107685. DOI: 10.1016/j.ecolind.2021.107685.
5. Hinge, G., Bharali, B., Baruah, A., et al. (2022). Integrated groundwater quality analysis using Water Quality Index, GIS and multivariate technique: A case study of Guwahati City. Environmental Earth Sciences, 81, 412. DOI: 10.1007/s12665-022-10544-0.
6. Kumar, M., Dash, A., Majid, S. I., Tiwari, A., Bhadwal, S., & Sahu, N. (2025). Delineation of groundwater potential zones of India using GIS-based Fuzzy-AHP techniques. Environmental and Sustainability Indicators, 100932. DOI: 10.1016/j.indic.2025.100932.
7. Masood, A., Aslam, M., Pham, Q. B., Khan, W., et al. (2022). Integrating water quality index, GIS and multivariate statistical techniques towards a better understanding of drinking water quality. Environmental Science and Pollution Research, 29, 26860–26876. DOI: 10.1007/s11356-021-17594-0.
8. Ram, A., Tiwari, S. K., Pandey, H. K., Chaurasia, A. K., Singh, S., & Singh, Y. V. (2021). Groundwater quality assessment using water quality index under GIS framework. Applied Water Science, 11, 46. DOI: 10.1007/s13201-021-01376-7.

9. Verma, P., Singh, P. K., Sinha, R. R., & Tiwari, A. K. (2020). Assessment of groundwater quality status using Water Quality Index and geographical information system approaches: A case study of Bokaro district, India. Applied Water Science, 10, 27. DOI: 10.1007/s13201-019-1088-4.

ABOUT THE EDITORS



Dr. Vijay Chandrakant Ithape

He is an Associate Professor of Geography at Shri Chhatrapati Shivaji Mahavidyalaya, Shrigonda, Maharashtra. He completed his M.A. in Geography from Sir Parashurambhau (S.P.) College, Pune, in 2007 and qualified the UGC-NET examination in 2009. Since joining higher education in 2010, he has accumulated over 16 years of teaching and academic experience.He was awarded a Ph.D. in Geography from SJJT University, Rajasthan, in 2024. His academic interests include Human Geography, Environmental Studies, Rural Development, and Contemporary Geographical Issues.Dr. Ithape has actively contributed to the field of geography through research publications, seminars, conferences, and academic activities. He is the autho of the book Contemporary Issues and Geography, which examines emerging geographical challenges and their implications for society, environment, and sustainable development. His scholarly work emphasizes the application of geographical concepts to contemporary real-world problems. He remains dedicated to excellence in teaching, research, and the advancement of geographical education.



Dr. Arpana Premi Lina Oraon

She is the Dean of the School of Tribal Resource Management and Director, Research and Development Cells at Kalinga Institute of Social Sciences, Deemed to be University, Odisha. She holds a Ph.D. in Geography and is UGC-NETqualified academician with a strong commitment to research and education. Her core areas of research include Regional Planning and Development, particularly with a focus on Indigenous communities and urban area development.Dr. Arpana has published more than 18 research papers in reputed national and international journals and has presented her work at over 25 national and international conferences. She has previously served at Nirmala College, Ranchi as an Assistant Professor and has years of experience in teaching and mentoring. Currently an Associate Professor in the Department of Geography at KISS-DU, she also heads several academic and administrative initiatives, contributing significantly to the advancement of tribal-centric research and knowledge systems.



Dr. Balasaheb Raghunath Chakor

He is an accomplished academician and geographer currently serving as Assistant Professor and Head, Department of Geography at Krantiveer Vasantrao Narayanrao Naik Shikshan Prasarak Sanstha's Arts, Commerce and Science College, Nashik. He holds M.A. (Geography), B.Ed., D.J., M.Phil., and Ph.D. degrees and has been actively engaged in higher education for more than 25 years as a senior college teacher.Dr. Chakor is a recognized Ph.D. Research Guide and has made significant contributions to geographical education and research. He has authored several academic books, including Tourism Geography (2021), Introduction to Human Geography (2024), Tourism Geography (Second Edition, 2025), and Political Geography (2025).His scholarly contributions include the publication of 27 research papers in various national and international journals and active participation in numerous state, national, and international seminars, workshops, and conferences. He is a Life Member of the Deccan Geographical Society of India and the Konkan Geographers Association.Apart from teaching and research, Dr. Chakor has held several important academic and extension responsibilities, including Coordinator of the Environmental Awareness Course, Coordinator of the Department of Lifelong Learning and Extension, and National Service Scheme (NSS) Programme Officer. He has also served on the College Local Development Committee and various academic committees, contributing significantly to institutional development.For his outstanding contributions to education and society, he has received the Vasantrao Naik State-Level Ideal Teacher Award from Vasant Social Foundation and the National Ideal Teacher Award.



Dr. K. Prabhu

He is an experienced academician, researcher, and geospatial science professional with over 16 Years of experience in teaching, research, and industry. He currently serves as an Assistant Professor of Physics at the Academy of Maritime Education and Training (AMET University), Kanathur, Chennai. He holds a Ph.D. in Integrated Physics with Geospatial Techniques from SRM Institute of Science and Technology, Chennai. His research focuses on Geophysical investigations, Remote Sensing & GIS, and Sustainable Environmental Management. He has published several Book Chapters, Research papers in SCI and SCOPUS indexed international journals and has presented his work at national and international conferences.

NP
Nature Light Publications

Nature Light Publications

309 West 11, Manjari VSI Road, Manjari Bk.,
Haveli, Pune- 412 307.
Website: www.naturelightpublications.com
Email: naturelightpublications@gmail.com
Contact No: +919822489040 / 9922489040

Nature- 118

ISBN: 978-93-49938-93-9
9 789349 938939
Price- 750/-