FRONTIERS IN ANIMAL BIOLOGY AND BIODIVERSITY



Editors

Dr. Esha Yadav

Ms. Hemangi Jayram Gavit

Dr. Swapna Sachin Patil

Mr. Anup Kumar Verma









Nature Light Publications

An International Edited Book *ISBN-978-93-49938-98-4*

Frontiers in Animal Biology and Biodiversity

Editors

Dr. Esha Yadav
Professor
Department of Zoology
Brahmanand P.G. College, The Mall,
Kanpur, Uttar Pradesh, India.

Ms. Hemangi Jayram Gavit
Assistant Professor
Department of Zoology
Rayat Shikshan Sanstha's Annasaheb Awate Arts, Commerce & Hutatma Babu Genu Science College, Manchar, Pune, (MH), India.

Dr. Swapna Sachin Patil
Assistant Professor and Head
Department of Zoology
Rayat Shikshan Sanstha's Annasaheb Awate Arts, Commerce & Hutatma Babu Genu Science College, Manchar, Pune, (MH), India.

Mr. Anup Kumar Verma
Research Scholar
Department of Zoology
Dr. Ram Manohar Lohia Avadh University, Ayodhya, Uttar Pradesh, India

Published By

Nature Light Publications, Pune

© Reserved by Editor's

FRONTIERS IN ANIMAL BIOLOGY AND BIODIVERSITY

Editors

Dr. Esha Yadav
Ms. Hemangi Jayram Gavit
Dr. Swapna Sachin Patil
Mr. Anup Kumar Verma

First Edition: June, 2026

An International Edited Book

ISBN- 978-93-49938-98-4

Published by:
Nature Light Publications, Pune
309 West 11, Manjari VSI Road, Manjari Bk.,
Haveli, Pune- 412 307.
Website: www.naturelightpublications.com
Email: naturelightpublications@gmail.com
Contact No: +91 9822489040 / 9922489040

The editors/Associate editors/publisher shall not be responsible for originality and thought expressed in the book chapter/ article. The author shall be solely held responsible for the originality and thoughts expressed in their book chapter or article.

Preface

The remarkable diversity of animal life forms the foundation of healthy ecosystems and sustains ecological processes that are essential for life on Earth. Rapid environmental changes, habitat degradation, climate change, pollution, and emerging infectious diseases have intensified the need for scientific research and innovative conservation strategies. In this context, Frontiers in Animal Biology and Biodiversity brings together contemporary research and interdisciplinary perspectives that address key challenges and emerging opportunities in animal biology, biodiversity conservation, environmental health, and sustainable development.

This edited volume presents contributions from scholars and researchers working across diverse fields of biological sciences. The chapters explore a broad spectrum of topics, including the diversity and ecological significance of invertebrates, biodiversity hotspots and conservation priorities, climate change impacts on ecosystems, artificial intelligence applications in biodiversity monitoring, multi-omics approaches in animal science, environmental toxicology, heavy metal contamination, zoonotic diseases from a One Health perspective, and the therapeutic potential of natural bioactive compounds such as curcumin. Together, these contributions reflect the growing integration of traditional biological research with modern technologies and interdisciplinary methodologies.

The book emphasizes that effective biodiversity conservation requires scientific innovation, evidence-based policy, technological advancement, and collaborative efforts among researchers, conservationists, healthcare professionals, and policymakers. It also highlights the interconnectedness of environmental integrity, animal health, and human well-being, reinforcing the importance of sustainable practices for future generations.

We sincerely appreciate all the authors who have contributed their valuable research and insights to this volume. Their dedication and scholarly

efforts have made this publication possible. We also extend our gratitude to the reviewers for their constructive suggestions, which have significantly enhanced the quality of the chapters. Special thanks are due to the editorial team and the publisher for their continuous support and commitment throughout the publication process.

We hope that Frontiers in Animal Biology and Biodiversity will serve as a valuable reference for researchers, academicians, students, environmental professionals, and policymakers. It is our sincere expectation that this volume will stimulate further research, promote scientific collaboration, and contribute meaningfully to the understanding, conservation, and sustainable management of animal biodiversity in an ever-changing world.

Editors

Frontiers in Animal Biology and Biodiversity

Table of Content

Sl. No.	Title and Authors	Page No.
1	**Invertebrates: Diversity, Classification, Ecological Roles, and Conservation** *Dr. Vivek Kumar, Sorabh*	01 - 12
2	**Climate Change Impacts on Ecosystems Challenges and Sustainable Solutions** *Dr. Muthurajan S, Immanuel Prabaharan S, Aswini M Deivanayaki R*	13 - 17
3	**Biodiversity Hotspots: Distribution, Significance, and Conservation Priorities** *Dr. Anand Kumar, Prof. Veena P. Swami*	18 - 32
4	**Artificial Intelligence in Biodiversity Monitoring** *Pradnyesh Valekar, Khodade Hrishikesh, Arpita Khatal*	33 - 39
5	**Biodiversity Hotspots and Conservation Priorities** *Pradnyesh Valekar, Khodade Hrishikesh, Arpita Khatal*	40 - 48
6	**Multi-Omics Approaches in Animal Science: Genomics, Proteomics and Metabolomics** *Ranjana*	49 - 70
7	**A Study on Therapeutic Potential of Curcumin in Cardio Protection** *Anand Konkala, A. Jayasankar, R. Kamli Naik, M. Venkata Chandrudu*	71 - 82
8	**Environmental and Health Consequences of Toxic Heavy Metal Exposure: A Comprehensive Review** *A. Jayasankar, Anand Konkala, M. Venkata Chandrudu R. Kamli Naik*	83 - 95
9	**Emerging and Re-emerging Zoonotic Diseases: A One Health Perspective** *Isai M, Annadurai T, Nagaraj K, Subavathy P*	96 -108
10	**Heavy Metals and Emerging Contaminants in Drinking Water: Ecological Risks and Health Implications** *Anup Kumar Verma, Dr. Abhinav Singh*	109 -118

Frontiers in Animal Biology and Biodiversity

ISBN: 978-93-49938-98-4 | Year: 2026 | pp: 01 – 12 |

Invertebrates: Diversity, Classification, Ecological Roles, and Conservation

[1]Dr. Vivek Kumar

[2]Sorabh

[1]Assistant Professor, Department of Zoology, School of Sciences, IFTM University, Moradabad, Uttar Pradesh (India)

[2]Research Scholar, Department of Zoology, School of Sciences, IFTM University, Moradabad, Uttar Pradesh (India)

Email: drvivek15010@gmail.com

Article DOI Link: https://zenodo.org/uploads/21281279

DOI: 10.5281/zenodo.21281279

Abstract

Invertebrates make up the majority of the world's biodiversity, accounting for about 95% of all known animal species. They live in a variety of freshwater, marine, and terrestrial habitats and are essential to preserving ecological equilibrium and ecosystem function. An overview of the biology and diversity of invertebrates is given in this chapter, with particular attention to their categorization, evolutionary relevance, ecological roles, and economic value. The discussion focuses on the unique traits and adaptations of major invertebrate phyla, such as Porifera, Cnidaria, Ctenophora, Platyhelminthes, Nematoda, Annelida, Mollusca, Arthropoda, and Echinodermata. The chapter looks at how invertebrates contribute to soil formation, pollination, decomposition, nutrient cycling, and food web dynamics. Additionally, it looks at how they are used in biotechnology, medicine, sericulture, apiculture, fisheries, and agriculture. Many invertebrate populations are becoming threatened by habitat loss, pollution, invasive species, overexploitation, and climate change, despite their enormous ecological and economic significance. Therefore, maintaining their diversity and ecosystem services requires sustainable management techniques and effective conservation methods. Recent developments in DNA barcoding, molecular taxonomy, genomics, and biodiversity assessment have greatly advanced our knowledge of the evolution, classification, and conservation of invertebrates. This chapter emphasizes the vital role that invertebrates play in maintaining biodiversity, ecosystem stability, and human well-being by fusing traditional zoological knowledge with contemporary scientific methods.

Keywords: Invertebrate Diversity, Animal Biodiversity, Invertebrate Phyla, Ecosystem Services, Conservation Biology.

Introduction

Nearly 95% of all known animal species are invertebrates, making them the most abundant and diverse category of animals on Earth. These animals are remarkably diverse in terms of shape, physiology, behavior, and ecological adaptations, and they do not have a vertebral column. As evidence of their ecological adaptability and evolutionary success, invertebrates can be found in almost any habitat, including freshwater, marine, terrestrial, and even harsh settings [2]. Each of the major invertebrate phyla—Porifera, Cnidaria, Ctenophora, Platyhelminthes, Nematoda, Annelida, Mollusca, Arthropoda, and Echinodermata—has distinctive structural and functional characteristics that support their ecological roles and evolutionary significance [4]. According to the latest Catalogue of Life database, invertebrates account for the vast majority of described animal species and encompass numerous taxonomic groups with extensive morphological and ecological diversity [1]. Invertebrates are essential to the stability and proper operation of ecosystems. They provide biological control, pollination, soil aeration, nutrient cycling, organic matter breakdown, and food-web dynamics. While terrestrial invertebrates, especially insects and annelids, promote agricultural productivity and biodiversity conservation, marine invertebrates, such as corals, mollusks, and echinoderms, are vital parts of aquatic ecosystems [3]. With over a million known species, arthropods alone make up the largest animal phylum and are important markers of environmental health [10]. Our knowledge of invertebrate phylogeny, species diversity, and evolutionary relationships has been enhanced as a result of recent developments in molecular systematics, genomics, and DNA barcoding [4]. Many invertebrate populations are decreasing despite their ecological and economic significance as a result of pollution, invasive species, habitat destruction, climate change, and unsustainable exploitation [3]. Therefore, ecosystem management, biodiversity protection, and the sustainable use of biological resources all depend on a thorough understanding of invertebrate biology and variety. An overview of the major invertebrate groups' categorization, biology, ecological relevance, and conservation is given in this chapter.

General Characteristics of Invertebrates

Invertebrates are the most varied category of animals on Earth, accounting for almost 95% of all known animal species. They live in a variety of habitats, including seas, freshwater ecosystems, woodlands, deserts, arctic areas, and other harsh settings, and they do not have a vertebral column. Their remarkable diversity in physiological adaptations, ecological roles, and bodily organization has greatly aided in their evolutionary success [2].

- **Absence of a Vertebral Column:** The lack of a backbone or vertebral column is what distinguishes all invertebrates. They lack an internal skeletal structure made up of vertebrae, in contrast to vertebrates. Alternative support structures, such as exoskeletons, shells, or hydrostatic skeletons, have been developed by numerous groups.
- **Levels of Body Organization:** The structural complexity of invertebrates varies. Cnidarians have tissue-level organization, while sponges (Porifera) have cellular organization. Organ-system organization is shown in more complex groupings like Annelida, Mollusca, Arthropoda, and Echinodermata, where specialized organs carry out specific tasks [4].
- **Body Symmetry:** Different types of body symmetry are observed among invertebrates:
  - **Asymmetrical body plan:** Found in most sponges.
  - **Radial symmetry:** Present in cnidarians and adult echinoderms.
  - **Bilateral symmetry:** Characteristic of most advanced invertebrate groups.

    Body symmetry is closely associated with movement, feeding behavior, and habitat adaptation.
- **Body Cavity (Coelom):** The body cavity or coelom varies among different invertebrate phyla:
  - **Acoelomates:** Lack a body cavity (e.g., Platyhelminthes).
  - **Pseudocoelomates:** Possess a false body cavity (e.g., Nematoda).
  - **Coelomates:** Have a true coelom lined by mesoderm (e.g., Annelida and Mollusca) [7].

    The presence of a coelom facilitates organ development and efficient body functioning.
- **Adaptations in Physiology:** Invertebrates have a variety of physiological systems that are suited to various environmental circumstances.
  - **Breathing** through the trachea, gills, book lungs, or body surface.
  - **Circulation** via closed or open circulatory systems.
  - **Excretion** by specialized glands, nephridia, Malpighian tubules, or flame cells.


    Survival in harsh, terrestrial, and aquatic environments is made possible by these adaptations.
- **Reproductive Strategies and Development:** Both sexual and asexual reproduction are used by invertebrates. While some species use internal fertilization, many display exterior fertilization. In insects, metamorphosis is especially highly developed, and complex developmental stages are typical. Their vast distribution and ecological success are facilitated by this reproductive variety [8].

- **Ecological and Evolutionary Significance:** Invertebrates have a vast evolutionary history spanning hundreds of millions of years, which is reflected in their astonishing morphological and functional variety. They carry out vital ecological tasks such pollination, soil formation, decomposition, nitrogen cycling, and food web maintenance. As a result, invertebrates are essential to ecosystem functioning and global biodiversity [13; 2].

Major Invertebrate Phyla: Classification and Diversity

Based on body organization, symmetry, embryological development, and evolutionary links, invertebrates—which make up the majority of animal diversity—are conventionally divided into a number of main phyla [2]. Every phylum has unique physiological and physical adaptations that support its ecological success.

Figure 1. Major Invertebrate Phyla and Representative Organisms




(A) (B) (C)

Figure 2. Diversity of Marine and Terrestrial Invertebrates



(A) (B) (C)

Diagram 1. Evolutionary Relationships among Major Invertebrate Phyla



- **Phylum Porifera (Sponges):** The most basic multicellular animals are represented by Porifera. Globally, some 9,000 species have been identified [2]. Although there are a few freshwater species, sponges are mostly marine. They lack real tissues and organs, and their bodies are lopsided. Numerous pores allow water to enter, while the osculum allows it to depart. Water currents are produced and feeding is made easier by specialized cells known as choanocytes. Because they recycle nutrients and filter vast amounts of water, sponges play a vital ecological role. Additionally, they serve as habitat for a variety of marine species [7]. Euplectella, Sycon, and Spongilla are a few examples.
- **Phylum Cnidaria:** Corals, hydras, jellyfish, and sea anemones are examples of cnidarians. There are currently over 11,000 known species [13]. These creatures have tissue-level organization and radial symmetry. Mesoglea separates the two germ layers that make up their bodies. The phylum is characterized by specialized stinging cells known as cnidocytes.
- **Cnidarians have two body forms:** medusa and polyp. Throughout their life cycle, many animals rotate between these phases. Because they create reef ecosystems that sustain a high level of marine biodiversity, corals are very

significant. According to [4], coral reefs are among the planet's most prolific ecosystems. Hydra, Aurelia, Metridium, and corals that form reefs are a few examples.

- **Phylum Ctenophora:** Comb jellies, or ctenophores, are only found in marine environments. There are currently about 200 known species [2]. They have eight rows of ciliary comb plates for movement and biradial symmetry. Ctenophores typically lack real stinging cells, in contrast to cnidarians. Numerous species produce amazing light shows in ocean waters due to their bioluminescence. In marine food webs, they play a significant role as predators. Ctenoplana and Pleurobrachia are two examples.
- **Phylum Platyhelminthes (Flatworms):** There are over 30,000 recognized species of platyhelminthes [7]. Members are triploblastic acoelomates with bilateral symmetry. They have flattened dorsoventral bodies. There are organ systems, but there are no respiratory or circulatory systems. Flatworms are both significant parasites and free-living organisms. Humans, animals, and wildlife are all impacted by parasitic species. Planaria, Taenia (tapeworm), and Fasciola (liver fluke) are a few examples. This group is where cephalization and bilateral symmetry first appear, marking a significant development in animal evolution [4].
- **Phylum Nematoda (Roundworms):** Among the most prevalent creatures on Earth are nematodes. Although the true variety is thought to approach one million species, more than 25,000 species have been officially described [13]. They have long, cylindrical bodies that are covered in a hard cuticle, making them pseudo-coelomate organisms. The mouth and anus are distinct parts of the digestive tract. Numerous nematodes can live freely in soil and water environments. Others are significant parasites that affect humans, animals, and plants. Globally, plant-parasitic nematodes result in large agricultural losses. In certain areas, human parasites like Ascaris and Wuchereria continue to be serious public health issues [3].
- **Phylum Annelida:** There are over 17,000 species of segmented worms known as annelids [2]. They have metameric segmentation and a genuine coelom.
  More specialization and effective mobility are made possible by segmentation. The circulatory system of the majority of annelids is closed. By aerating the soil and breaking down organic debris, earthworms increase soil fertility. Many marine benthic ecosystems are dominated by polychaetes. Because of their anticoagulant secretions, leeches are used in microsurgical techniques and have medical relevance. Hirudo, Nereis, and Pheretima are examples.
- **Phylum Mollusca:** After Arthropoda, Mollusca is the second-largest phylum of animals. There are over 100,000 known species [2]. Mollusks usually have a mantle, visceral bulk, and a muscular foot. A calcareous shell is secreted by several organisms. There are many very different classes in the phylum. Slugs

and snails are examples of gastropods. Clams, mussels, and oysters are examples of bivalves. Octopuses, cuttlefish, and squids are examples of cephalopods. Cephalopods have sophisticated neural systems and intricate behaviors. Octopuses are remarkably adept at learning and solving problems [7]. Fisheries, aquaculture, and pearl production all greatly benefit from mollusks. Octopus, Sepia, Unio, and Pila are a few examples.

- **Phylum Arthropoda:** The largest and most prosperous animal phylum is Arthropoda. Over 80% of all known animal species are represented by the more than one million species that have been described [10]. Segmented bodies, jointed appendages, and a chitinous exoskeleton are the distinguishing characteristics. Growth throughout development is made possible by molting. Almost every ecosystem on Earth is home to arthropods. Their ability to adapt, fly, have specialized appendages, and have effective sensory systems are all factors in their diversity.
- **Class Insecta:** The largest class of animals is represented by insects. They have three body regions: the head, thorax, and abdomen. The majority of species possess three pairs of legs and one or two pairs of wings. Insects carry out essential ecological roles, including pollination, decomposition, and maintaining biological control. Groups such as honeybees, butterflies, and beetles hold economic significance [3].
- **Class Arachnida:** Spiders, scorpions, mites, and ticks are all types of arachnids. They have four pairs of walking legs and no antennae. Numerous species act as predators, controlling the populations of insects. Certain ticks and mites are disease carriers.
- **Class Crustacea:** Crustaceans comprise crabs, lobsters, shrimp, and copepods. The majority live in water and have two sets of antennae. They play a significant role in both commercial fisheries and aquatic food webs.
- **Class Myriapoda:** Centipedes and millipedes are part of the myriapod family. They have long, segmented bodies and many legs. While centipedes are active predators, millipedes mainly serve as decomposers.
- **Phylum Echinodermata:** Exclusively marine, echinoderms consist of about 7,000 living species [2]. While larvae show bilateral symmetry, adults display pentaradial symmetry. The water vascular system, which serves the purposes of locomotion, feeding, and respiration, is the most distinctive feature. Echinoderms have extraordinary regenerative capabilities. Some sea stars have the ability to regenerate entire arms or even their entire bodies from fragments. These animals are crucial for the ecology of marine ecosystems, as they regulate benthic communities and nutrient cycling. Examples are sea cucumbers, brittle stars, sea urchins, crinoids, and sea stars.

Ecological Roles of Invertebrates

Invertebrates are vital components of freshwater, marine, and terrestrial ecosystems. Their ecological contributions support ecosystem stability, biodiversity, and the delivery of essential ecosystem services. Because of their enormous diversity and richness, invertebrates have an impact on almost all ecological processes in natural settings [2].

- **Nutrient Cycling and Decomposition:** An important ecological role played by invertebrates is nutrient cycling and decomposition. Termites, millipedes, earthworms, and other insect larvae break down dead organic matter and hasten the recycling of nutrients into the soil. This procedure encourages plant development and improves soil fertility [7]. Because earthworms promote soil aeration, water penetration, and microbial activity through their burrowing activities, they are often referred to as "ecosystem engineers."
- **Pollination:** Pollination also depends on invertebrates. By spreading pollen from one bloom to another, bees, butterflies, moths, beetles, and flies aid in the reproduction of flowering plants. Insects are the primary pollinators for over 75% of the world's food crops, which rely on animal pollination to some degree [6]. As a result, pollination is crucial to both global food security and agricultural output.
- **Food Web Dynamics and Biological Control:** Many invertebrates are essential components of food webs. Fish, amphibians, reptiles, birds, and mammals rely heavily on zooplankton, mollusks, insects, and crustaceans as food sources. Spiders and dragonflies are examples of predatory invertebrates that control prey populations and support ecological balance [3]. These natural enemies are crucial to biological pest control in agricultural habitats.
- **Ecological Functions in Marine Ecosystems:** The ecological roles of marine invertebrates are similarly significant. For thousands of marine species, coral polyps create reef ecosystems that serve as breeding grounds, havens, and habitats. While echinoderms aid in the cycling of nutrients and the control of benthic ecosystems, bivalves like mussels and oysters enhance water quality through filter feeding [4].
- **Bioindicators of Environmental Quality:** Many invertebrates serve as bio indicators of the quality of the environment. Insect, mollusk, and aquatic macro invertebrate diversity and abundance changes frequently point to pollution, habitat loss, or the effects of climate change [3].

Economic and Applied Importance of Invertebrates

Through their contributions to agriculture, industry, medicine, fisheries, and environmental management, invertebrates significantly contribute to human welfare. Because they offer a variety of ecosystem services and biological resources necessary for sustainable development, they have enormous economic worth [2].

Importance in Agriculture and Pollination: Among the most useful invertebrates are pollinating insects. Many agricultural crops are pollinated by honeybees, butterflies, moths, beetles, and flies. Global food security is greatly enhanced by pollination, which increases crop output, quality, and genetic variety [6]. Additionally, honeybees generate profitable goods like propolis, royal jelly, beeswax, and honey.

Importance in Industry and Agriculture: Important industries are supported by a number of invertebrates. The foundation of the sericulture sector is the silkworm (Bombyx mori), which produces the natural silk utilized in textile production all over the world. In many coastal areas, mollusks like oysters and pearl oysters contribute to aquaculture and pearl production, which generates significant revenue [7]. Vermiculture and vermicomposting make heavy use of earthworms. They are important elements of organic farming systems and sustainable agriculture because their actions enhance soil structure, nutrient availability, and organic waste decomposition [8].

Importance in Fisheries and Food Production: Crustaceans and mollusks are examples of marine invertebrates that sustain commercial fisheries and serve as a significant source of protein for millions of people worldwide. Among the aquatic resources with the highest economic value are shrimp, crabs, lobsters, mussels, and oysters [4].

Medical and Biotechnological Applications: Additionally, invertebrates have significant uses in biotechnology and medicine. Because leeches have anticoagulant qualities, they are used in reconstructive surgery. Moreover, a variety of marine invertebrates generate bioactive substances that may be useful in medicine. Genetics, developmental biology, and the biomedical sciences have all benefited greatly from research on insects and other invertebrates [2].

Threats to Invertebrate Diversity and Conservation

The bulk of the world's biodiversity is made up of invertebrates, which also carry out vital ecological tasks that maintain the productivity and health of ecosystems. However, invertebrate populations around the world are under tremendous strain due to growing anthropogenic activity. Numerous invertebrate groups have shown significant decreases, according to recent studies, endangering biodiversity conservation, ecosystem stability, and the ecosystem services that support human societies [3; 9]

- **Habitat Loss and Fragmentation:** One of the biggest dangers to the diversity of invertebrates is habitat degradation and fragmentation. The availability of suitable habitats is diminished and ecological connection is disrupted by urbanization, agricultural expansion, deforestation, wetland drainage, and infrastructural development. Numerous specialized invertebrates experience population reductions and local extinctions as a result of their inability to adjust to swift environmental changes [6].

- **Climate Change:** One of the main causes of biodiversity loss has been identified as climate change. The distribution, reproduction, and survival of many invertebrate species are impacted by rising temperatures, changing precipitation patterns, ocean acidification, and extreme weather events. Because rising sea temperatures cause coral bleaching and habitat destruction, coral reef ecosystems are especially vulnerable [2].
- **Pollution:** Invertebrate populations in both terrestrial and aquatic environments are seriously threatened by pollution. Insects, aquatic macro invertebrates, and other vulnerable species are negatively impacted by the overuse of pesticides, industrial chemicals, plastics, and heavy metals. Pesticide exposure and habitat degradation have resulted in significant decreases in pollinator populations, particularly those of bees and butterflies [9]
- **Invasive Species and Conservation Challenges:** Through competition, predation, and disease transmission, invasive alien species provide an additional hazard to native invertebrate groups. To maintain invertebrate variety and ecosystem functioning, effective conservation necessitates pollution control, sustainable land-use practices, habitat protection, and climate-change mitigation strategies [3].

Emerging Research Trends in Invertebrate Biology

Invertebrate biology has changed as a result of recent developments in molecular biology, genomics, bioinformatics, and ecological monitoring technology. Our knowledge of invertebrate evolution, biodiversity, physiology, and ecological relationships has improved as a result of these advancements. New prospects for species identification, conservation planning, biotechnology, and biomedical applications are being made possible by emerging research trends [2].

- **Molecular Taxonomy and DNA Barcoding:** The use of DNA barcoding and molecular taxonomy for species identification and biodiversity assessment is among the most important developments. Standardized genetic markers, specifically the mitochondrial cytochrome c oxidase I (COI) gene, are used in DNA barcoding to precisely identify species and uncover hidden biodiversity [5]. By better classifying insects, mollusks, crustaceans, and other invertebrates, this method has aided conservation and biodiversity monitoring initiatives [4].
- **Genomics and Evolutionary Research:** Researchers may now examine the entire genomes of many invertebrate species because to developments in next-generation sequencing. Evolutionary linkages, developmental routes, adaption processes, and gene functions are all explained by comparative genomic studies. Understanding physiological reactions to environmental changes is further aided by transcriptomics and proteomics [7].
- **Environmental DNA (eDNA) and Biodiversity Monitoring:** Environmental DNA (eDNA) and metabarcoding have become popular tools for assessing

biodiversity. Scientists can identify several species without direct collection by examining genetic material found in samples of water, soil, or sediment. Ecological monitoring and biodiversity surveys have been transformed by this non-invasive method [11].

- **Climate Change and Bioindicator Research:** The effects of climate change on the distribution, reproduction, and survival of invertebrates are the subject of more and more research. Due to their ecological significance, freshwater macro invertebrates, coral reef creatures, and pollinators are gaining particular study [12]. Furthermore, mollusks, crustaceans, and aquatic insects are frequently employed as bio indicators to evaluate ecosystem health and environmental quality [3]. Biotechnology and Biomedical Applications: Bioactive substances with antibacterial, antiviral, anti-inflammatory, and anticancer qualities are produced by marine invertebrates such sponges, corals, tunicates, and mollusks. These substances are now useful tools for pharmaceutical development and drug discovery. In a similar vein, industrial biotechnology, agriculture, and medicine are increasingly using insect-derived proteins, enzymes, and nanomaterials [2].

Conclusion

Global biodiversity is based on invertebrates, which are the most prevalent and varied group of species on Earth. They can inhabit almost any environment and ecological niche thanks to their unique evolutionary history, which has produced an astounding variety of body shapes, physiological adaptations, and ecological specializations. Invertebrates exhibit remarkable biological diversity, ranging from basic sponges to highly organized arthropods and cephalopods. The species richness of invertebrates is only one aspect of their ecological significance. Nutrient cycling, decomposition, soil formation, pollination, biological control, water purification, and food-web dynamics all depend on them. Through agriculture, fisheries, aquaculture, sericulture, apiculture, biotechnology, and biomedical research, they also contribute significantly to the economy. Ecosystem stability, food security, and human welfare all depend on these contributions. Many invertebrate populations are diminishing despite their significance as a result of invasive species, pollution, habitat loss, climate change, and unsustainable resource usage. These dangers emphasize how urgently good conservation and sustainable environmental management are needed. Our knowledge of invertebrate diversity and evolution has significantly enhanced as a result of recent developments in molecular taxonomy, genomics, environmental DNA analysis, artificial intelligence, and biodiversity monitoring. In order to preserve invertebrate biodiversity and guarantee the long-term viability of ecosystems and the vital services they offer to humanity, it is imperative that research, public awareness, and evidence-based conservation efforts continue.

References

1. Banki, O. S., Roskov, Y., Döring, M., & Ower, G. (Eds.). (2024). Catalogue of Life Checklist 2024. Leiden, The Netherlands: Catalogue of Life.
2. Brusca, R. C., Moore, W., & Shuster, S. M. (2023). Invertebrates (4th ed.). Sunderland, MA: Sinauer Associates/Oxford University Press.
3. Cardoso, P., Barton, P. S., Birkhofer, K., Chichorro, F., Deacon, C., Fartmann, T., ... Samways, M. J. (2020). Scientists' warning to humanity on insect extinctions. Biological Conservation, 242, 108426. doi: 10.1016/j.biocon.2020.108426
4. Giribet, G., & Edgecombe, G. D. (2020). The invertebrate tree of life. Princeton, NJ: Princeton University Press.
5. Hebert, P. D. N., Ratnasingham, S., & deWaard, J. R. (2003). Biological identifications through BDNA barcodes. Proceedings of the Royal Society B: Biological Sciences, 270(1512), 313–321. doi:10.1098/rspb.2002.2218
6. IPBES. (2019). Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Bonn, Germany: IPBES Secretariat.
7. Pechenik, J. A. (2024). Biology of the invertebrates (8th ed.). New York, NY: McGraw-Hill Education.
8. Ruppert, E. E., Fox, R. S., & Barnes, R. D. (2021). Invertebrate zoology: A functional evolutionary approach (8th ed.). Belmont, CA: Brooks/Cole.
9. Sánchez-Bayo, F., & Wyckhuys, K. A. G. (2019). Worldwide decline of the entomofauna: A review of its drivers. Biological Conservation, 232, 8–27. doi: 10.1016/j.biocon.2019.01.020
10. Stork, N. E. (2018). How many species of insects and other terrestrial arthropods are there on Earth? Annual Review of Entomology, 63, 31–45. doi:10.1146/annurev-ento-020117-043348
11. Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). Environmental DNA: For biodiversity research and monitoring. Oxford, UK: Oxford University Press.
12. Wagner, D. L., Grames, E. M., Forister, M. L., Berenbaum, M. R., & Stopak, D. (2021). Insect decline in the Anthropocene: Death by a thousand cuts. Proceedings of the National Academy of Sciences, 118(2), e2023989118. doi:10.1073/pnas.2023989118
13. Zhang, Z.-Q. (2022). Animal biodiversity: An update of classification and diversity in 2022. Zootaxa, 5141(1), 1–6. doi:10.11646/zootaxa.5141.1.1

Frontiers in Animal Biology and Biodiversity

ISBN: 978-93-49938-98-4 | Year: 2026 | pp: 13 – 17 |

Climate Change Impacts on Ecosystems Challenges and Sustainable Solutions

[1]Dr. Muthurajan S

[2]Immanuel Prabaharan S

[2]Aswini M

[2]Deivanayaki R

[1]Assistant Professor, Department of Marine Engineering, AMET University, Kanathur, Chennai-603 112

[2]Research Scholar, Department of Electrical and Electronics Engineering AMET University, Kanathur, Chennai-603 112

Email: imman.amet@gmail.com

Article DOI Link: https://zenodo.org/uploads/21281614

DOI: 10.5281/zenodo.21281614

Abstract

Climate change has emerged as one of the most significant environmental challenges of the 21st century, profoundly affecting ecosystems worldwide. Rising global temperatures, altered precipitation patterns, and increasing frequency of extreme weather events are disrupting biodiversity, ecosystem services, and ecological balance. This chapter examines the impacts of climate change on terrestrial, aquatic, and marine ecosystems, highlighting key challenges such as habitat loss, species extinction, and ecosystem degradation. It also explores sustainable solutions, including ecosystem-based adaptation, conservation strategies, and climate mitigation approaches. The chapter emphasizes the importance of interdisciplinary efforts, policy interventions, and technological innovation to ensure ecological resilience and sustainability.

Keywords: Climate change, Ecosystems, Biodiversity loss, Sustainable solutions, Adaptation, Mitigation, Ecosystem resilience, Environmental sustainability

Introduction

Climate change refers to long-term alterations in temperature, precipitation, and atmospheric conditions, largely driven by human activities such as greenhouse gas

emissions and deforestation. These changes are accelerating at an unprecedented rate, with global temperatures already exceeding 1.5°C above pre-industrial levels in recent years.

Ecosystems including forests, oceans, wetlands, and grasslands play a critical role in maintaining environmental balance and supporting life. However, climate change is disrupting these systems by altering species distributions, ecological interactions, and natural cycles.

Methodology

This chapter is based on a systematic review of recent literature (2024–2025) from peer-reviewed journals, reports, and global organizations. The methodology includes

- Collection of secondary data from scientific databases
- Comparative analysis of ecosystem responses
- Synthesis of mitigation and adaptation strategies
- Evaluation of interdisciplinary approaches

The approach integrates ecological, environmental, and policy perspectives to provide a comprehensive understanding.

Climate Change Impacts on Ecosystems

1. Terrestrial Ecosystems

Climate change affects forests, grasslands, and deserts through

- Rising temperatures causing drought and wildfires
- Shifts in vegetation zones
- Increased land degradation

Land ecosystems are losing their ability to act as carbon sinks due to deforestation and climate stress, which may accelerate global warming.

2. Aquatic Ecosystems (Freshwater)

Freshwater systems such as rivers and lakes are highly vulnerable

- Changes in water availability and quality
- Increased salinity and temperature
- Loss of aquatic biodiversity

Climate change is fundamentally transforming global water systems and their ecological dynamics

3. Marine Ecosystems

Marine ecosystems face

- Ocean warming and acidification
- Coral bleaching
- Decline in marine biodiversity

For example, coral reefs have experienced severe degradation due to heat stress and bleaching events.

4. Biodiversity Loss

Climate change is a major driver of biodiversity loss

- Species migration and extinction
- Disruption of food chains
- Loss of ecosystem services

Studies indicate that biodiversity decline is occurring across nearly all ecosystems due to combined human and climate pressures.

Key Challenges

1. Ecosystem Instability

Rapid climate shifts disrupt ecological balance and resilience.

2. Loss of Ecosystem Services

Ecosystems provide essential services such as

- Carbon sequestration
- Water purification
- Food production

These services are increasingly threatened.

3. Extreme Weather Events

Floods, droughts, and heatwaves are intensifying, damaging ecosystems.

4. Feedback Mechanisms

Degraded ecosystems may release carbon instead of storing it, worsening climate change.

Sustainable Solutions

1. Ecosystem-Based Adaptation (EbA)

- Restoration of forests and wetlands
- Conservation of biodiversity hotspots
- Sustainable land-use planning

2. Climate Mitigation Strategies

- Reduction of greenhouse gas emissions
- Transition to renewable energy
- Carbon capture and storage

3. Sustainable Agriculture

- Agroforestry

- Organic farming
- Efficient water management

Many mitigation strategies also provide co-benefits for biodiversity and human well-being.

4. Policy and Governance

- International agreements (e.g., Paris Agreement)
- National climate policies
- Community-based conservation

5. Technological Innovations

- Climate monitoring systems
- AI-based environmental modelling
- Sustainable infrastructure

Case Studies

1. Arctic Ecosystems

Rapid warming is causing

- Melting ice
- Changing vegetation
- Threats to wildlife

2. Coral Reefs

Frequent bleaching events highlight vulnerability.

3. Forest Ecosystems

Deforestation and wildfires are reducing carbon storage capacity.

Discussion

Climate change impacts are interconnected and multifaceted. Ecosystems are not only affected by climate change but also influence it through feedback loops. Sustainable solutions must integrate ecological, economic, and social dimensions. Interdisciplinary approaches and global cooperation are essential to address these challenges effectively.

Conclusion

Climate change poses a severe threat to ecosystems worldwide, disrupting biodiversity and ecosystem services essential for human survival. However, sustainable solutions such as ecosystem restoration, climate mitigation, and policy interventions offer hope. Immediate and coordinated global action is necessary to enhance ecosystem resilience and ensure a sustainable future.

References

1. Smith, P., Arneth, A., Barnes, D. K. A., Ichii, K., Marquet, P. A., Popp, A., Pörtner, H.-O., Rogers, A. D., Scholes, R. J., & Steffen, W. (2025). Impacts of climate change interventions on biodiversity, water, food and health.
2. Amorim-Maia, A. T., Anguelovski, I., Chu, E., & Connolly, J. J. T. (2025).
3. A new paradigm for climate change adaptation: Integrating development and resilience.
4. Mubangizi, J. (2025). Investigating climate change and emerging theoretical perspectives: A systematic review.
5. Hao, F. (2025). Global research of climate change: Causes, impacts and sustainability transitions.
6. Akande, H. A., & Gidado, A. A. (2025). EcoCast: A spatio-temporal model for biodiversity and climate risk forecasting.
7. Raine, S., & Fischer, T. (2025). AI-driven marine robotics for ecosystem monitoring under climate change.
8. Hatton, M., Laverick, J. H., Banas, N., & Heath, M. (2025). Ecosystem recovery under combined fishing and climate pressures in the Barents Sea.
9. Demirel, N., Vogiatzakis, I. N., Zittis, G., Tase, M., Sandor, A. D., Zotos, S., Zoumides, C., Dindaroglu, T., Fois, M., Christoforidi, I., Stamatiadou, V., Zemah-Shamir, S., Albayrak, T., Ayhan, C. K., Manolaki, P., Sieber, I., Zemah-Shamir, Z., Tzirkalli, E., & Moustakas, A. (2026).
10. Uncertainty in island-based ecosystem services and climate change.
11. Dang, H.-A. H., Hallegatte, S., & Trinh, T.-A. (2025). Impacts of global warming on poverty and inequality.
12. Pintea, S., Acevedo, A., & Marrero, A. (2025). Climate change impacts on vulnerable populations and ecosystems in the Americas.

Frontiers in Animal Biology and Biodiversity

ISBN: 978-93-49938-98-4 | Year: 2026 | pp: 18 – 32 |

Biodiversity Hotspots: Distribution, Significance and Conservation Priorities

Dr. Anand Kumar

Prof. Veena P. Swami

Department of Zoology, B.S.N.V. P.G. College (University of Lucknow), Lucknow-226001 (U.P.), India

Email: anandkumar051976@gmail.com

Article DOI Link: https://zenodo.org/uploads/21281688

DOI: 10.5281/zenodo.21281688

Abstract

Biodiversity hotspots represent geographically distinct regions that contain extraordinary concentrations of endemic species and ecosystems but are experiencing significant habitat degradation and biodiversity loss. These areas are considered priority regions for global conservation because they support a disproportionately large share of the Earth's biological wealth within relatively small land areas. The biodiversity hotspot concept provides an effective framework for identifying regions where conservation efforts can yield the greatest benefits. This chapter outlines the origin and criteria of biodiversity hotspots, their major ecological features, global distribution, and conservation significance. Particular attention is given to the four biodiversity hotspots associated with India: the Himalaya, Indo-Burma, Western Ghats–Sri Lanka, and Sundaland. The chapter further examines the ecological importance of these regions, the principal threats affecting their biodiversity, and the conservation strategies required for their long-term protection. Sustainable management of biodiversity hotspots is essential for conserving species, maintaining ecosystem services, and promoting environmental security for future generations.

Keywords: Biodiversity Hotspots, Concept and Characteristics, Global Distribution, Ecological Significance, Threats, and Conservation Priorities

Introduction

Biodiversity encompasses the variety of life forms found on Earth, including genetic diversity within species, diversity among species, and the diversity of ecosystems (Wilson, 1988; Wilson, 1992). It forms the foundation of ecosystem

functioning and provides numerous services essential for human survival, such as food production, pollination, nutrient cycling, climate regulation, and water purification (Gaston, 2000; Millennium Ecosystem Assessment, 2005).

Rapid population growth, habitat conversion, overexploitation of resources, pollution, invasive species, and climate change have accelerated biodiversity loss across the globe (Sala et al., 2000; Dirzo et al., 2014). These pressures have resulted in habitat degradation and increasing extinction risks for many species. Since conservation resources are limited, identifying areas where protection efforts can produce the greatest impact has become increasingly important (Margules & Pressey, 2000).

The biodiversity hotspot concept emerged as an effective approach for prioritizing conservation action. Hotspots are regions characterized by exceptional concentrations of endemic species coupled with extensive habitat loss (Myers, 1988; Myers et al., 2000). Conservation of these areas is considered essential for safeguarding global biodiversity and maintaining ecological stability (Mittermeier et al., 2011).

Concept of Biodiversity Hotspots

The concept of biodiversity hotspots was proposed by Norman Myers in 1988 as a means of identifying biologically rich regions facing severe threats from habitat destruction (Myers, 1988). Initially focused on tropical forests, the concept was later expanded through international conservation initiatives and scientific research (Myers et al., 2000).

Biodiversity hotspots are recognized as regions where extraordinary biological diversity coincides with significant habitat degradation (Mittermeier et al., 2004). Because these areas contain numerous endemic species that occur nowhere else, habitat loss can lead directly to global extinctions (Gaston, 2000). Consequently, hotspots have become central to conservation planning and resource allocation worldwide (Brooks et al., 2006; Mittermeier et al., 2011).

Criteria for Identification of Biodiversity Hotspots

According to Conservation International, a region must satisfy two primary criteria to qualify as a biodiversity hotspot (Mittermeier et al., 2004):

- It must contain at least 1,500 endemic vascular plant species.
- It must have lost at least 70% of its original natural vegetation.

These criteria ensure that hotspot regions possess both exceptional biological importance and a high degree of conservation urgency (Myers et al., 2000).

Characteristics of Biodiversity Hotspots

Biodiversity hotspots are characterized by exceptionally high species richness, elevated levels of endemism, and diverse ecosystem types (Wilson, 1992; Primack, 2014). Geographic isolation, environmental heterogeneity, and long evolutionary

histories have contributed to the development of unique biological communities within these regions (Myers et al., 2000).

These areas often support tropical forests, montane ecosystems, wetlands, grasslands, and island habitats that harbor numerous rare and threatened species (Millennium Ecosystem Assessment, 2005). However, many hotspots remain highly vulnerable to anthropogenic disturbances such as deforestation, agricultural expansion, mining, and urbanization (Dirzo et al., 2014).

Global Distribution of Biodiversity Hotspots

Biodiversity hotspots are distributed across different continents and islands of the world, reflecting regions where evolutionary processes, environmental heterogeneity, and geographic isolation have produced exceptionally rich biological communities. At present, 36 biodiversity hotspots have been recognized globally (Table-1). Although these areas collectively occupy only a small proportion of the Earth's land surface, they support a substantial share of the world's endemic flora and fauna (Myers et al., 2000; Mittermeier et al., 2011).




Fig. 1. Global Distribution of Biodiversity Hotspots

In North and Central America, important hotspot regions include the California Floristic Province, Madrean Pine–Oak Woodlands, Mesoamerica, and the Caribbean Islands. These areas are known for their diverse vegetation types, high levels of plant endemism, and numerous endemic vertebrate species. Tropical forests, Mediterranean-type ecosystems, and island habitats contribute significantly to their biological richness.

South America contains several of the world's most species-rich hotspot regions. The Tropical Andes is regarded as the most biologically diverse hotspot, supporting extraordinary numbers of plants, birds, amphibians, and mammals. Other important

hotspots include the Atlantic Forest, Cerrado, Tumbes–Chocó–Magdalena, and the Chilean Winter Rainfall–Valdivian Forests. These ecosystems are increasingly threatened by agricultural expansion, deforestation, and urban development.

Europe and Central Asia contain hotspots such as the Mediterranean Basin, Caucasus, Irano-Anatolian region, and Mountains of Central Asia. These areas are characterized by complex topography, diverse climatic conditions, and high plant diversity. Many species occurring in these regions have adapted to unique environmental conditions over long evolutionary periods.

Africa possesses a remarkable variety of biodiversity hotspots. The Cape Floristic Region and Succulent Karoo are globally renowned for their exceptional floral diversity. Other important hotspots include the Coastal Forests of Eastern Africa, Eastern Afromontane, Horn of Africa, Guinean Forests of West Africa, Maputaland–Pondoland–Albany, and Madagascar with the Indian Ocean Islands. The long-term geographic isolation of Madagascar has resulted in extraordinary levels of species endemism.

Table-1: Global distribution of biodiversity hotspots

Region	**Biodiversity Hotspots**
North and Central America	California Floristic Province; Madrean Pine-Oak Woodlands; Mesoamerica; Caribbean Islands
South America	Tropical Andes; Tumbes-Chocó-Magdalena; Atlantic Forest; Cerrado; Chilean Winter Rainfall-Valdivian Forests
Europe and Central Asia	Mediterranean Basin; Caucasus; Irano-Anatolian; Mountains of Central Asia
Africa	Cape Floristic Region; Succulent Karoo; Maputaland-Pondoland-Albany; Coastal Forests of Eastern Africa; Eastern Afromontane; Horn of Africa; Guinean Forests of West Africa; Madagascar and the Indian Ocean Islands
South Asia	Himalaya; Indo-Burma; Western Ghats and Sri Lanka
East and Southeast Asia	Mountains of Southwest China; Japan; Sundaland; Wallacea; Philippines
Pacific Region	East Melanesian Islands; Polynesia-Micronesia; New Caledonia; New Zealand
Australia	Southwest Australia; Forests of East Australia
North America	North American Coastal Plain

In South Asia, three major hotspots are recognized: the Himalaya, Indo-Burma, and Western Ghats–Sri Lanka. These regions contain diverse ecosystems ranging from tropical forests and river systems to alpine habitats and are among the most important centers of biodiversity in Asia.

East and Southeast Asia include the Mountains of Southwest China, Japan, Sundaland, Wallacea, and the Philippines hotspots. These regions support extensive tropical forests, mountain ecosystems, and island habitats that harbor numerous endemic species. Several globally threatened mammals, birds, reptiles, and amphibians occur within these hotspot regions.

The Pacific region comprises hotspots such as New Caledonia, New Zealand, East Melanesian Islands, and Polynesia–Micronesia. Geographic isolation has facilitated the evolution of unique plant and animal groups that are found nowhere else in the world. Australia also contains globally important hotspots, including Southwest Australia and the Forests of East Australia, both recognized for their remarkable plant diversity and ecological uniqueness.

Together, these hotspot regions represent the most biologically valuable and threatened terrestrial ecosystems on Earth. Their conservation remains a priority for maintaining global biodiversity and ecological stability.

Biodiversity Hotspots in India

India is recognized as one of the world's megadiverse nations because of its exceptional variety of ecosystems, climatic conditions, and biological resources. The country's diverse landscapes, ranging from the Himalayas in the north to tropical forests and coastal ecosystems in the south, support an enormous wealth of plant and animal life. Four globally recognized biodiversity hotspots are represented within Indian territory: the Himalaya, Indo-Burma, Western Ghats–Sri Lanka, and Sundaland (Myers et al., 2000).

1. Himalaya Hotspot

The Himalayan hotspot extends across a vast mountain system and includes a wide range of ecological zones, from subtropical forests at lower elevations to alpine meadows and snow-covered peaks at higher altitudes. The region supports rich floral diversity and contains numerous endemic plant species. The varied topography and climatic gradients create suitable habitats for many rare and threatened organisms.

Several iconic species inhabit this hotspot, including the snow leopard (*Panthera uncia*), red panda (*Ailurus fulgens*), Himalayan musk deer (*Moschus chrysogaster*), and Himalayan monal (*Lophophorus impejanus*). However, increasing human activities such as road construction, hydropower projects, tourism development, and land-use changes have intensified habitat fragmentation. Climate change is further affecting glacier dynamics and species distributions within the region.

2. Indo-Burma Hotspot

The Indo-Burma hotspot includes northeastern India and extends into several Southeast Asian countries. This region is characterized by tropical and subtropical forests that support exceptional levels of biodiversity. Numerous endemic species of birds, amphibians, reptiles, fishes, and flowering plants occur within this hotspot.

The region also contains diverse freshwater ecosystems that provide important habitats for aquatic organisms. Despite its biological richness, the Indo-Burma hotspot faces considerable pressure from shifting cultivation, logging, infrastructure expansion, and conversion of natural habitats into agricultural land. Effective conservation measures are therefore essential for preserving its unique biodiversity

3. Western Ghats–Sri Lanka Hotspot

The Western Ghats–Sri Lanka hotspot is among the most significant biodiversity regions in the world. Stretching parallel to India's western coast, the Western Ghats contain tropical evergreen forests, moist deciduous forests, montane ecosystems, and freshwater habitats. The hotspot exhibits exceptionally high levels of endemism, particularly among amphibians, reptiles, freshwater fishes, and flowering plants.

Several threatened species are restricted to this region, including the lion-tailed macaque (*Macaca silenus*), Nilgiri tahr (*Nilgiritragus hylocrius*), Malabar civet, and numerous endemic amphibians. Rapid urbanization, mining activities, plantation agriculture, infrastructure projects, and habitat fragmentation continue to threaten ecological integrity within the hotspot.

4. Sundaland Hotspot

The Sundaland hotspot includes parts of Southeast Asia and extends into India's Nicobar Islands. This hotspot is dominated by tropical rainforests that support extraordinary biological diversity. The region is home to numerous endemic plant and animal species, including several globally threatened mammals, birds, reptiles, and amphibians.

Within India, the Nicobar Islands contribute significantly to the biological importance of Sundaland. These islands harbor unique ecosystems and endemic species that have evolved under conditions of geographic isolation. However, habitat degradation, climate change, coastal disturbances, invasive species, and developmental activities pose growing threats to biodiversity in this region.

Collectively, the four biodiversity hotspots associated with India represent invaluable reservoirs of biological diversity and ecological resources. Their protection is crucial for conserving endemic species, sustaining ecosystem services, and ensuring long-term environmental stability.




Fig. 2. Distribution of Biodiversity Hotspots in India

Ecological Importance of Biodiversity Hotspots

Biodiversity hotspots play a fundamental role in maintaining the health and stability of the Earth's ecosystems. Although they occupy a relatively small portion of the planet's terrestrial area, these

regions support an extraordinary concentration of species and ecological processes that are vital for environmental sustainability. Their significance extends beyond the conservation of rare organisms, influencing ecosystem functioning, climate regulation, and human well-being.

1. Preservation of Genetic Resources

One of the greatest ecological values of biodiversity hotspots lies in their rich genetic diversity. Many endemic species occurring within these regions possess unique genetic traits that have evolved over thousands or millions of years. Such genetic variation enhances the adaptive capacity of species and provides important

resources for future agricultural, medicinal, and scientific applications. Conserving these genetic resources helps maintain evolutionary processes and reduces the risk of irreversible biodiversity loss.




Fig. 3. Ecological importance of Biodiversity Hotspots

2. Maintenance of Ecosystem Stability

Ecosystems with high biological diversity generally exhibit greater stability and resilience. Biodiversity hotspots support complex food webs and ecological interactions that contribute to ecosystem productivity and balance. The presence of multiple species performing similar ecological functions increases the ability of ecosystems to withstand disturbances such as droughts, floods, disease outbreaks, and other environmental stresses. Consequently, biodiversity-rich ecosystems often recover more rapidly following natural or human-induced disturbances.

3. Provision of Ecosystem Services

The ecosystems present within biodiversity hotspots generate numerous services that directly and indirectly support human societies. Forests, wetlands, grasslands, and coastal ecosystems regulate water flow, improve water quality, prevent soil erosion, store carbon, and influence local and global climate systems. Pollinators found in these regions contribute significantly to agricultural productivity, while natural vegetation helps maintain nutrient cycles and air quality. The economic and social benefits derived from these ecosystem services underscore the importance of hotspot conservation.

4. Scientific and Educational Importance

Biodiversity hotspots serve as valuable centers for scientific investigation and environmental education. Their exceptional species richness provides opportunities

to study evolutionary biology, ecology, conservation science, and species interactions. Many discoveries related to biodiversity, adaptation, and ecosystem functioning have emerged from research conducted in hotspot regions. In addition, these areas help promote environmental awareness and provide educational opportunities for students, researchers, and conservation practitioners.

5. Contribution to Global Conservation Goals

The protection of biodiversity hotspots contributes significantly to global conservation objectives. Since many endemic species are restricted to these regions, their conservation helps prevent global extinctions. Furthermore, safeguarding hotspot ecosystems supports international commitments related to biodiversity conservation, climate change mitigation, and sustainable development. Effective management of these areas therefore benefits both regional ecosystems and global environmental security.

Major Threats to Biodiversity Hotspots

Despite their ecological significance, biodiversity hotspots are increasingly threatened by a variety of human activities and environmental changes. These pressures continue to reduce habitat quality, disrupt ecological processes, and endanger numerous species.

1. Habitat Destruction and Fragmentation

Conversion of natural habitats for agriculture, settlements, industries, roads, and infrastructure remains the most serious threat to biodiversity hotspots. Habitat fragmentation isolates wildlife populations, restricts movement, reduces gene flow, and increases the vulnerability of species to extinction.




Fig. 4. Major Threats to Biodiversity Hotspots

2. Climate Change

Changes in temperature, rainfall patterns, and the frequency of extreme weather events are altering ecosystems worldwide. Species with limited geographic ranges are particularly vulnerable because they may be unable to adapt or migrate to suitable habitats. Mountain ecosystems, coastal regions, and island habitats are especially sensitive to climate-related impacts.

3. Unsustainable Resource Exploitation

Excessive extraction of timber, fuelwood, medicinal plants, wildlife, and fishery resources can significantly reduce biodiversity. Unsustainable harvesting often disrupts ecological relationships and weakens the regenerative capacity of ecosystems.

4. Pollution

Industrial discharges, agricultural chemicals, plastic waste, and urban pollutants negatively affect terrestrial and aquatic ecosystems. Pollution can alter habitat quality, reduce reproductive success, and threaten the survival of sensitive species.

5. Invasive Alien Species

Non-native organisms introduced intentionally or accidentally may compete with native species for resources, alter habitat structure, spread diseases, and disrupt ecological balance. Island ecosystems and isolated habitats are particularly vulnerable to biological invasions.

6. Population Growth and Developmental Pressure

Rapid human population growth increases the demand for land, water, energy, and natural resources. Expanding urbanization and infrastructure development frequently occur at the expense of natural habitats, thereby intensifying biodiversity loss.

The cumulative effects of these threats result in species decline, genetic erosion, ecosystem degradation, reduced ecosystem services, and increased risks of extinction.

Conservation Priorities of Biodiversity Hotspots

Given their exceptional biological importance and vulnerability, biodiversity hotspots require carefully planned conservation interventions. Strategic conservation priorities can maximize biodiversity protection while supporting sustainable development objectives.

1. Protection of Natural Habitats

Conserving remaining natural habitats is the cornerstone of biodiversity protection. Establishing and strengthening protected areas such as national parks, wildlife sanctuaries, biosphere reserves, and community conservation areas can significantly reduce habitat loss and safeguard critical ecosystems.




Fig. 5. Conservation Priorities of Biodiversity Hotspots

2. Conservation of Endemic and Threatened Species

Species with restricted distributions and declining populations require focused conservation measures. Monitoring programs, habitat management, captive breeding initiatives, and species recovery plans can improve survival prospects for endangered organisms.

3. Ecological Restoration

Restoration of degraded landscapes has become an essential component of modern conservation. Activities such as reforestation, wetland rehabilitation, invasive species control, and soil restoration can enhance ecosystem functioning and improve biodiversity conservation outcomes.

4. Enhancement of Landscape Connectivity

Maintaining ecological connectivity between fragmented habitats enables wildlife movement, dispersal, and genetic exchange. Wildlife corridors and landscape-level planning can reduce isolation and improve ecosystem resilience, particularly under changing climatic conditions.

5. Climate-Smart Conservation Strategies

Conservation planning must increasingly account for climate change impacts. Protecting climate refuges, maintaining elevational gradients, and adopting adaptive management approaches can help species and ecosystems cope with future environmental changes.

6. Community Participation and Indigenous Knowledge

Local communities and indigenous peoples often possess extensive knowledge regarding ecosystem management and resource use. Their involvement in

conservation planning and implementation can improve effectiveness while promoting social equity and long-term sustainability.

7. Sustainable Livelihood Development

Reducing dependence on destructive resource-use practices is crucial for conservation success. Sustainable agriculture, agroforestry, ecotourism, non-timber forest products, and nature-based enterprises can provide economic benefits while minimizing environmental impacts.

8. Management of Invasive Species

Preventing the introduction and spread of invasive organisms should be a priority in hotspot regions. Early detection, rapid response mechanisms, and long-term monitoring programs can reduce ecological damage and protect native biodiversity.

9. Research and Biodiversity Monitoring

Scientific research provides essential information for conservation decision-making. Biodiversity inventories, ecological assessments, long-term monitoring, and emerging technologies such as GIS, remote sensing, environmental DNA, and artificial intelligence can strengthen conservation planning and management.

10. Policy Support and Good Governance

Effective conservation requires strong legal frameworks, institutional support, and transparent governance. Environmental policies should promote sustainable resource management, strengthen law enforcement, and encourage stakeholder participation in conservation programs.

11. Sustainable Financing Mechanisms

Long-term conservation initiatives require reliable financial support. Funding sources may include government allocations, international conservation programs, environmental trust funds, ecosystem service payments, carbon financing mechanisms, and private-sector partnerships.

12. Environmental Education and Public Awareness

Conservation outcomes are greatly enhanced when society understands the value of biodiversity. Educational programs, awareness campaigns, citizen science initiatives, and outreach activities can encourage responsible environmental behavior and foster support for conservation efforts.

Future Directions

Future biodiversity conservation efforts should adopt integrated approaches that combine ecological protection with sustainable development and climate resilience. Advances in biotechnology, geospatial technologies, artificial intelligence, and environmental monitoring are expected to improve conservation effectiveness. International collaboration, transboundary conservation initiatives, and stronger

implementation of global biodiversity agreements will be essential for addressing emerging environmental challenges.

The successful implementation of the Kunming–Montreal Global Biodiversity Framework and the global commitment to protect 30 percent of the planet's land and marine environments by 2030 offer important opportunities for strengthening conservation efforts worldwide. Achieving these goals will require cooperation among governments, researchers, local communities, non-governmental organizations, and the private sector.

Conclusion

Biodiversity hotspots represent some of the most valuable ecological regions on Earth because they combine extraordinary biological richness with high levels of endemism and vulnerability. Although these areas occupy a limited proportion of the Earth's surface, they support a substantial share of global biodiversity and provide essential ecosystem services that benefit both nature and human societies.

However, ongoing pressures from habitat destruction, climate change, pollution, invasive species, and unsustainable resource use continue to threaten these ecosystems. Effective conservation therefore requires a combination of habitat protection, ecosystem restoration, species conservation, scientific research, community participation, sustainable resource management, and supportive policy frameworks.

Protecting biodiversity hotspots is not only necessary for preventing species extinctions but also for maintaining ecosystem stability, safeguarding natural resources, and promoting sustainable development. Continued commitment to conservation at local, national, and international levels will be essential to ensure that these irreplaceable ecosystems remain functional and resilient for future generations.

References

1. Aronson, J., & Alexander, S. (2013). Ecosystem restoration is now a global priority: Time to roll up our sleeves. Restoration Ecology, 21(3), 293–296. https://doi.org/10.1111/rec.12011
2. Berkes, F. (2018). Sacred ecology (4th ed.). Routledge.
3. Bonan, G. B. (2008). Forests and climate change: Forcings, feedbacks, and the climate benefits of forests. Science, 320(5882), 1444–1449. https://doi.org/10.1126/science.1155121
4. Brooks, T. M., Mittermeier, R. A., da Fonseca, G. A. B., Gerlach, J., Hoffmann, M., Lamoreux, J. F., Mittermeier, C. G., Pilgrim, J. D., & Rodrigues, A. S. L. (2006). Global biodiversity conservation priorities. Science, 313(5783), 58–61. https://doi.org/10.1126/science.1127609
5. Convention on Biological Diversity. (2022). Kunming–Montreal global biodiversity framework. Secretariat of the Convention on Biological Diversity.

6. Daniels, R. J. R. (2003). Biodiversity of the Western Ghats and conservation priorities. Universities Press.
7. Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014). Defaunation in the Anthropocene. Science, 345(6195), 401–406. https://doi.org/10.1126/science.1251817
8. Frankham, R., Ballou, J. D., & Briscoe, D. A. (2010). Introduction to conservation genetics (2nd ed.). Cambridge University Press.
9. Gadgil, M., & Guha, R. (1995). Ecology and equity: The use and abuse of nature in contemporary India. Routledge.
10. Gaston, K. J. (2000). Global patterns in biodiversity. Nature, 405(6783), 220–227. https://doi.org/10.1038/35012228
11. Heller, N. E., & Zavaleta, E. S. (2009). Biodiversity management in the face of climate change. Biological Conservation, 142(1), 14–32. https://doi.org/10.1016/j.biocon.2008.10.019
12. Hooper, D. U., Chapin, F. S., III, Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., Schmid, B., Setälä, H., Symstad, A. J., Vandermeer, J., & Wardle, D. A. (2005). Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. Ecological Monographs, 75(1), 3–35. https://doi.org/10.1890/04-0922
13. Intergovernmental Panel on Climate Change. (2023). Climate change 2023: Synthesis report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC.
14. Joly, C. A. (2022). The Kunming–Montréal Global Biodiversity Framework. Biota Neotropica, 22(4), e2022e001. https://doi.org/10.1590/1676-0611-BN-2022-e001
15. Margules, C. R., & Pressey, R. L. (2000). Systematic conservation planning. Nature, 405(6783), 243–253. https://doi.org/10.1038/35012251
16. Millennium Ecosystem Assessment. (2005). Ecosystems and human well-being: Biodiversity synthesis. World Resources Institute.
17. Ministry of Environment, Forest and Climate Change (MoEFCC). (2014). India: Fifth national report to the Convention on Biological Diversity. Government of India.
18. Mittermeier, R. A., Gil, P. R., Hoffmann, M., Pilgrim, J. D., Brooks, T. M., Mittermeier, C. G., Lamoreux, J. F., & da Fonseca, G. A. B. (2004). Hotspots revisited: Earth's biologically richest and most endangered terrestrial ecoregions. Conservation International.
19. Mittermeier, R. A., Turner, W. R., Larsen, F. W., Brooks, T. M., & Gascon, C. (2011). Global biodiversity conservation: The critical role of hotspots. In F. E. Zachos & J. C. Habel (Eds.), Biodiversity hotspots: Distribution and protection of conservation priority areas (pp. 3–22). Springer. https://doi.org/10.1007/978-3-642-20992-5_1

20. Myers, N. (1988). Threatened biotas: "Hotspots" in tropical forests. The Environmentalist, 8(3), 187–208. https://doi.org/10.1007/BF02240252
21. Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B., & Kent, J. (2000). Biodiversity hotspots for conservation priorities. Nature, 403(6772), 853–858. https://doi.org/10.1038/35002501
22. Primack, R. B. (2014). Essentials of conservation biology (6th ed.). Sinauer Associates.
23. Rodgers, W. A., & Panwar, H. S. (1988). Planning a wildlife protected area network in India (Vols. 1–2). Wildlife Institute of India.
24. Sala, O. E., Chapin, F. S., III, Armesto, J. J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-Sanwald, E., Huenneke, L. F., Jackson, R. B., Kinzig, A., Leemans, R., Lodge, D. M., Mooney, H. A., Oesterheld, M., Poff, N. L., Sykes, M. T., Walker, B. H., Walker, M., & Wall, D. H. (2000). Global biodiversity scenarios for the year 2100. Science, 287(5459), 1770–1774. https://doi.org/10.1126/science.287.5459.1770
25. Singh, J. S., & Singh, S. P. (1992). Forests of Himalaya: Structure, functioning and impact of man. Gyanodaya Prakashan.
26. Tilman, D., Reich, P. B., & Knops, J. M. H. (2006). Biodiversity and ecosystem stability in a decade-long grassland experiment. Nature, 441(7093), 629–632. https://doi.org/10.1038/nature04742
27. Wilson, E. O. (1988). Biodiversity. National Academy Press.
28. Wilson, E. O. (1992). The diversity of life. Harvard University Press.

Frontiers in Animal Biology and Biodiversity

ISBN: 978-93-49938-98-4 | Year: 2026 | pp: 33 – 39 |

Artificial Intelligence in Biodiversity Monitoring

Pradnyesh Valekar

Khodade Hrishikesh

Arpita Khatal

Department of Botany, Sharadchandra Pawar Mahavidyalaya Lonand

Email: pradnyeshvalekar122@gmail.com

Article DOI Link: https://zenodo.org/uploads/21281748

DOI: 10.5281/zenodo.21281748

Abstract

Artificial intelligence (AI) is becoming a potent tool for monitoring biodiversity, allowing for quick, precise, and extensive evaluation of species and ecosystems. AI facilitates automated species identification, habitat mapping, and environmental change detection by combining machine learning, deep learning, computer vision, remote sensing, bioacoustics, and environmental DNA (eDNA). These tools monitor invasive and endangered species, enhance conservation planning, and offer real-time data for well-informed decision-making. AI provides enormous potential to improve ecosystem management, biodiversity conservation, and international efforts to meet sustainable development and biodiversity preservation goals, despite obstacles including data quality, computing demands, and ethical issues.

Keywords: Artificial intelligence, Biodiversity, Conservation, DNA, Ecosystem, Environment


Introduction

A key element of conservation biology is biodiversity monitoring, which offers crucial data on the distribution, status, and patterns of ecosystems and species. Conventional monitoring techniques, such field surveys and manual species identification, are frequently labour-intensive, time-consuming, and have limited temporal and spatial coverage. The demand for effective and scalable methods to monitor biodiversity and assist prompt conservation decisions is expanding due to the mounting challenges posed by habitat loss, climate change, pollution, invasive species, and overuse of natural resources.

By analysing vast and intricate ecological datasets, artificial intelligence (AI) has become a game-changing tool that improves biodiversity monitoring. AI processes

data gathered from camera traps, drones, satellite imaging, acoustic sensors, and environmental DNA (eDNA) by combining machine learning, deep learning, computer vision, natural language processing, and big data analytics (Yambal & Waykar, 1 C.E.). Compared to traditional methods, these technologies allow for faster and more accurate automated species identification, habitat mapping, population calculation, and early environmental change detection.

AI also makes it easier to identify conservation priorities in the face of shifting environmental conditions, anticipate species distributions, and conduct real-time monitoring. AI facilitates evidence-based conservation planning and ecosystem management by lowering human labour and enhancing data analysis. AI is anticipated to become more crucial in protecting the world's biodiversity and accomplishing global conservation and sustainable development objectives as technology develops (Ali et al., 2025).



Objectives

This chapter aims to describe the main AI technologies used in ecological research, explain the concept and significance of AI in biodiversity monitoring, and look at their applications in species identification, habitat assessment, population monitoring, and ecosystem management. The chapter also seeks to emphasize how AI can be used to monitor invasive and endangered species, identify environmental changes, and assist with conservation planning. Additionally, it aims to address the benefits, difficulties, moral issues, and potential applications of AI-based biodiversity monitoring, highlighting its role in evidence-based environmental decision-making and sustainable biodiversity conservation.

Role of AI in Biodiversity Monitoring

AI improves monitoring of biodiversity by:

- Recognizing species of plants and animals from photos, videos, and audio files.
- Identifying rare, invasive, and endangered species.
- Keeping an eye on ecosystem health and habitat changes (Lahoz-Monfort & Magrath, 2021).
- Forecasting species dispersal in the context of climate change.
- Automating the analysis of big ecological datasets.
- Encouraging protected area management and conservation planning.

Major AI Technologies Used

1. Machine Learning (ML)

Machine learning algorithms forecast trends in biodiversity and find patterns in ecological datasets (Eastwood et al., 2022).

Uses

- Classification of species
- Analysis of population trends

2. Deep Learning

Artificial neural networks are used in deep learning to evaluate complex ecological data (Christin et al., 2019).

Uses

- Image recognition of wildlife
- Automated counting of species
- Analysis of the forest canopy

3. Computer Vision

Computer vision allows computers to decipher pictures and movies that are obtained by Camera traps - Drones, Satellite photos.

Uses

- The detection of animals
- Identification of individual animals
- Mapping the forest cover

4. Bioacoustics and AI

AI examines noises captured by acoustic sensors.

Uses

- Identification of birds

- Monitoring frogs
- The detection of marine mammals

5. Natural Language Processing (NLP)

NLP gathers data on biodiversity from:

- Academic literature
- Records from museums
- Observations from citizen science

AI-Based Monitoring Tools

1. Camera Traps

AI uses millions of photos to automatically identify different species of wildlife.

Drones Utilized for:

- Monitoring forests
- The detection of illegal logging
- Mapping habitats

2. Satellite Remote Sensing

AI examines satellite photos to identify:

- Deforestation
- Degradation of wetlands
- The growth of cities
- Damage from fire

3. Environmental DNA (eDNA)

AI uses DNA fragments taken from soil, water, or the air to identify species without the need for human observation (Yang et al., 2024).

Applications in Biodiversity Conservation

1. Monitoring of Wildlife Populations

AI calculates the distribution and magnitude of endangered species' populations.

Examples

- Tigers
- Elephants
- Snow leopards
- Whales

2. Forest Conservation

AI detects:

- Illegal logging
- Forest fires

- Habitat fragmentation
- Carbon stock changes

3. Marine Biodiversity

AI monitors:

- Coral reefs
- Fish populations
- Sea turtles
- Marine mammals

4. Identification of Invasive Species

AI enables quick management responses by providing early warning systems for invasive plants and animals (Fricke & Olden, 2023).

5. Evaluation of Climate Change

AI helps with adaptation planning by identifying climate refugia and forecasting changes in species distributions (Levy & Shahar, 2024).

Advantages of AI in Biodiversity Monitoring

- An increase in species identification accuracy.
- Constant, real-time observation.
- Lower field survey expenses (Sun et al., 2021).
- Early identification of environmental risks.
- Greater backing for conservation planning.
- Enhanced long-term ecological monitoring effectiveness.

Challenges and Limitations

- High-quality training data is scarce.
- Prediction accuracy may be impacted by dataset bias.
- The initial cost of computer infrastructure and equipment is high.
- Skilled workers in ecology and AI are needed (Bergougui, 2025).
- Privacy and ethical issues with data acquisition.
- Dependency on dependable power and internet infrastructure in isolated locations (Pipattanasomporn, 2004).

Future Prospects

The following are some aspects of AI's future in biodiversity monitoring:

- Integration with sensor networks for the Internet of Things (IoT).
- Systems for monitoring biodiversity in real time.
- AI-powered conservation planning.
- Robotic surveillance and autonomous drones (Sebastian, 2025).
- Better eDNA analysis with sophisticated machine learning.

- Early warning systems for biodiversity worldwide (Antonelli et al., 2023).
- AI-powered mobile applications that improve citizen science.

Conclusion

By offering effective, precise, and scalable techniques for species identification, habitat assessment, and ecosystem management, artificial intelligence is transforming biodiversity monitoring. Conservationists may make quicker, more informed judgments by combining AI with remote sensing, camera traps, bioacoustics, eDNA, and citizen research. Even though there are still issues with data quality, cost, and technical know-how, further developments in AI are anticipated to improve global biodiversity conservation and play a major role in accomplishing the Sustainable Development Goals (SDGs) and the Convention on Biological Diversity (CBD) objectives.

References

1. Ali, G., Mijwil, M. M., Adamopoulos, I., & Ayad, J. (2025). Leveraging the Internet of Things, Remote Sensing, and Artificial Intelligence for Sustainable Forest Management. Babylonian Journal of Internet of Things, 2025, 1–65. https://doi.org/10.58496/BJIOT/2025/001
2. Antonelli, A., Dhanjal-Adams, K. L., & Silvestro, D. (2023). Integrating machine learning, remote sensing and citizen science to create an early warning system for biodiversity. Plants People Planet, 5(3), 307–316. https://doi.org/10.1002/PPP3.10337;JOURNAL:JOURNAL:25722611;REQUESTEDJOURNAL:JOURNAL:25722611;WGROUP:STRING:PUBLICATION
3. Bergougui, B. (2025). Can Artificial Intelligence Technologies Advance Environmental Sustainability? The Role of Institutional Adaptability and Skill-Biased Technological Transformation. Sustainable Development, 34, 222–244. https://doi.org/10.1002/SD.70296;ISSUE:ISSUE:DOI
4. Christin, S., Hervet, É., & Lecomte, N. (2019). Applications for deep learning in ecology. Methods in Ecology and Evolution, 10(10), 1632–1644. https://doi.org/10.1111/2041-210X.13256;CTYPE:STRING:JOURNAL
5. Eastwood, N., Stubbings, W. A., Abou-Elwafa Abdallah, M. A., Durance, I., Paavola, J., Dallimer, M., Pantel, J. H., Johnson, S., Zhou, J., Hosking, J. S., Brown, J. B., Ullah, S., Krause, S., Hannah, D. M., Crawford, S. E., Widmann, M., & Orsini, L. (2022). The Time Machine framework: monitoring and prediction of biodiversity loss. Trends in Ecology and Evolution, 37(2), 138–146. https://doi.org/10.1016/j.tree.2021.09.008
6. Fricke, R. M., & Olden, J. D. (2023). Technological innovations enhance invasive species management in the anthropocene. BioScience, 73(4), 261–279. https://doi.org/10.1093/BIOSCI/BIAD018

7. Lahoz-Monfort, J. J., & Magrath, M. J. L. (2021). A Comprehensive Overview of Technologies for Species and Habitat Monitoring and Conservation. BioScience, 71(10), 1038–1062. https://doi.org/10.1093/BIOSCI/BIAB073
8. Levy, O., & Shahar, S. (2024). Artificial Intelligence for Climate Change Biology: From Data Collection to Predictions. Integrative and Comparative Biology, 64(3), 953–974. https://doi.org/10.1093/ICB/ICAE127
9. Pipattanasomporn, M. (2004). A Study of Remote Area Internet Access with Embedded Power Generation.
10. Sebastian, P. (2025). Editorial: Applications of AI in autonomous, surveillance, and robotic systems. Frontiers in Robotics and AI, 12, 1616634. https://doi.org/10.3389/FROBT.2025.1616634/TEXT
11. Sun, J., Futahashi, R., & Yamanaka, T. (2021). Improving the Accuracy of Species Identification by Combining Deep Learning with Field Occurrence Records. Frontiers in Ecology and Evolution, 9, 762173. https://doi.org/10.3389/FEVO.2021.762173/TEXT
12. Yambal, S., & Waykar, Y. A. (1 C.E.). AI powered Conservation Transforming Habitat Management for a Sustainable Future. Https://Services.Igi-Global.Com/Resolvedoi/Resolve.Aspx?Doi=10.4018/979-8-3693-6829-9.Ch003, 47–80. https://doi.org/10.4018/979-8-3693-6829-9.CH003
13. Yang, J., Li, C., Lo, L. S. H., Zhang, X., Chen, Z., Gao, J., U, C., Dai, Z., Nakaoka, M., Yang, H., & Cheng, J. (2024). Artificial Intelligence-Assisted Environmental DNA Metabarcoding and High-Resolution Underwater Optical Imaging for Noninvasive and Innovative Marine Environmental Monitoring. Journal of Marine Science and Engineering 2024, Vol. 12, Page 1729, 12(10), 1729. https://doi.org/10.3390/JMSE12101729

Frontiers in Animal Biology and Biodiversity

ISBN: 978-93-49938-98-4 | Year: 2026 | pp: 40 – 48 |

Biodiversity Hotspots and Conservation Priorities

Pradnyesh Valekar

Khodade Hrishikesh

Arpita Khatal

Department of Botany, Sharadchandra Pawar Mahavidyalaya Lonand

Email: pradnyeshvalekar122@gmail.com

Article DOI Link: https://zenodo.org/uploads/21281838

DOI: 10.5281/zenodo.21281838

Abstract

High concentrations of endemic species and substantial habitat degradation are characteristics of biodiversity hotspots, which are areas of extraordinary biological richness. Despite making up only 2.5% of the planet's land area, these regions are home to approximately 43% of endemic terrestrial animals and over half of all endemic vascular plants. Norman Myers established the hotspot idea, which offers a scientific basis for setting global conservation priorities. The distribution and features of the world's 36 biodiversity hotspots are examined in this chapter, along with the main threats they face, such as habitat destruction and climate change, as well as the international strategies and conservation priorities needed to protect these priceless ecosystems for coming generations.

Keywords: Biodiversity, Conservation, Endemism, Framework, Hotspot, Western Ghats.

Introduction

The variety of living things, such as plants, animals, microbes, and the environments in which they exist, is referred to as biodiversity (Groombridge & Jenkins, 2010). It provides vital functions like food production, climate management, nutrient cycling, pollination, and water purification. It is the cornerstone of ecosystem functioning. However, the loss of biodiversity has accelerated globally due to growing human activities, such as deforestation, urbanization, industrialization, and climate change. In order to overcome this difficulty, Norman Myers developed the idea of biodiversity hotspots in 1988 as a method for locating areas with remarkable densities of endemic species that are seriously threatened by habitat degradation (Noss, 2013). Because they contain a


significant number of the world's rare species within comparatively small geographic areas, biodiversity hotspots are locations of high conservation value. There are now 36 biodiversity hotspots identified worldwide. They support a large portion of the world's biodiversity despite making up a very small portion of the planet's land area, making their conservation a top priority for ecological stability and sustainable development (Hrdina & Romportl, 2017).

Objectives

This chapter's goals include defining biodiversity hotspots, outlining their global distribution, highlighting their ecological and conservation significance, identifying the main threats to biodiversity, and discussing global conservation priorities, strategies, and international initiatives for the efficient preservation and sustainable management of these vital ecosystems.

Concept of Biodiversity Hotspots

In order to identify areas that are both unusually rich in biodiversity and seriously threatened by human activity, Norman Myers (1988) established the idea of biodiversity hotspots. Prioritizing conservation efforts in regions where scarce resources can have the biggest impact on averting species extinction is the goal of the hotspot method. A place that has lost at least 70% of its original natural vegetation and has at least 1,500 indigenous vascular plant species is considered a biodiversity hotspot (Mittermeier et al., 2011). High levels of endemism, species richness, and ecological distinctiveness define these regions. Despite making up only 2.5% of the planet's land area, biodiversity hotspots are home to approximately 43% of endemic terrestrial animals and over 50% of endemic plant species worldwide. In order to designate priority areas for biodiversity protection, habitat restoration, and sustainable management, governments, researchers, and conservation organizations use the hotspot idea, which has gained widespread acceptance as a framework for global conservation planning (Sarkar et al., 2006).



Global Distribution of Biodiversity Hotspots

Currently, 36 biodiversity hotspots are recognized worldwide.

North and Central America

- California Floristic Province (USA)
- Madrean Pine-Oak Woodlands (Mexico & USA)
- Mesoamerica (Central America)
- Caribbean Islands

South America

- Tropical Andes
- Tumbes-Chocó-Magdalena
- Atlantic Forest (Brazil)
- Cerrado (Brazil)
- Chilean Winter Rainfall-Valdivian Forests

Europe and Central Asia

- Mediterranean Basin
- Caucasus
- Irano-Anatolian
- Mountains of Central Asia

Africa

- Cape Floristic Region (South Africa)
- Succulent Karoo (South Africa & Namibia)
- Maputaland-Pondoland-Albany
- Coastal Forests of Eastern Africa
- Eastern Afromontane
- Horn of Africa
- Guinean Forests of West Africa
- Madagascar and the Indian Ocean Islands

South Asia

- Himalaya
- Indo-Burma
- Western Ghats and Sri Lanka

East Asia

- Mountains of Southwest China
- Japan

Southeast Asia

- Sundaland
- Wallacea
- Philippines

Australasia and Pacific Islands

- East Melanesian Islands
- New Caledonia
- New Zealand
- Polynesia-Micronesia
- Southwest Australia

Additional Hotspots

- Forests of East Australia
- North American Coastal Plain

Importance of Biodiversity Hotspots

Among the most crucial areas for the preservation of biodiversity worldwide are the 36 biodiversity hotspots. More than 50% of all endemic plant species and approximately 43% of all endemic terrestrial animal species are found there, although making up only 2.5% of the planet's land area. The preservation of the planet's biological diversity and ecological equilibrium depends on their conservation (Dinerstein et al., 2019).

Its high degree of endemism is one of the main factors contributing to its significance. There are numerous species that are unique to these hotspots. Many plant and animal species would become permanently extinct if these habitats were lost (Biological Aspects of Endemism in Higher Plants on JSTOR, n.d.). Hotspots known for their distinctive biodiversity include Madagascar, the Tropical Andes, and the Western Ghats–Sri Lanka region.

Additionally, these hotspots offer vital ecosystem services that promote human welfare. They help with pollination, preserve soil fertility, safeguard watersheds, control the climate, and help sequester carbon. Wetlands and forests in hotspots provide millions of people with safe water while mitigating the effects of droughts, floods, and climate change (Guo et al., 2010).

Agriculture, forestry, fishing, medicine, and ecotourism are all economically supported by biodiversity hotspots. These areas are the source of numerous significant crops, therapeutic plants, and genetic resources. By offering chances for employment and a means of subsistence, they make a substantial contribution to both local and national economies.

Additionally, the 36 hotspots are extremely valuable in terms of science and education. Evolution, ecology, conservation biology, and climate change can all be studied in these natural labs. Opportunities to find new species and create creative

solutions for environmental problems are provided by their vast biodiversity (Fiebelkorn & Menzel, 2012).

Additionally, indigenous and local groups, whose customs, cultures, and means of subsistence are intimately associated with natural environments, frequently reside in biodiversity hotspots. Preserving these regions contributes to the preservation of both biological and cultural diversity.

The 36 biodiversity hotspots are regarded as worldwide conservation priorities due to their remarkable biodiversity and growing risks from habitat degradation, climate change, invasive species, and overexploitation. One of the best ways to preserve biodiversity and guarantee sustainable development for coming generations is to protect them.

Threats to Biodiversity Hotspots

Hotspots for biodiversity are among the planet's most ecologically rich areas, yet they are also the most vulnerable. Numerous species are in danger of going extinct due to the deterioration of these habitats caused by both natural and human-caused processes. Loss and fragmentation of habitat pose the biggest concern (Hogue & Breon, 2022). Large tracts of natural habitats have been destroyed by the rapid deforestation caused by mining, industrial operations, infrastructure development, urbanization, and agriculture. Fragmentation disturbs natural processes, isolates populations, and lowers genetic diversity. Another significant issue is climate change. Sea level rise, harsh weather, shifting rainfall patterns, and rising global temperatures all modify habitats and have an impact on species distribution. Due to their restricted geographic ranges, many endemic species in biodiversity hotspots may not be able to adapt to shifting environmental conditions (Brierley & Kingsford, 2009).

Biodiversity is also at risk due to overuse of natural resources. Population decreases and possibly the extinction of species can result from unsustainable logging, hunting, fishing, and medicinal plant harvesting (Rawat & Agarwal, 2015).

In many hotspot regions, the illegal wildlife trafficking accelerates the loss of biodiversity. A major worry is invasive alien species. In addition to changing ecological functioning and spreading diseases, non-native plants, animals, and microbes frequently outcompete native species for resources.

Lantana camara and Water hyacinth (Eichhornia crassipes) are two examples that have adversely impacted numerous ecosystems across the globe (Yadav et al., 2016). Air, water, and soil are contaminated by sewage, plastics, industrial waste, and agricultural chemicals. Degradation of ecosystems results from pollution's effects on the survival and well-being of microbes, plants, and animals.

Furthermore, irresponsible land-use patterns and population growth put further strain on natural resources. Ecologically sensitive places are frequently invaded by growing agricultural lands and human populations.

Wildfires, floods, droughts, and storms are examples of natural disasters that can harm ecosystems rich in biodiversity, particularly when combined with environmental stress brought on by humans (Loucks, 2021).

Integrated conservation measures, such as habitat protection, sustainable resource management, climate change mitigation, ecosystem restoration, stringent environmental regulations, and community involvement, are necessary to address these concerns.

To protect biodiversity hotspots and their priceless ecological services for future generations, effective intervention is crucial.

Conservation Priorities

The world's 36 biodiversity hotspots, which have exceptional levels of endemism and biodiversity yet are seriously threatened by habitat loss and environmental degradation, are worldwide conservation priorities. To save these priceless ecosystems and stop additional species loss, effective conservation measures are crucial.

Protecting the surviving natural habitats through the creation and efficient administration of protected areas, such as national parks, wildlife sanctuaries, biosphere reserves, and community conservation areas, is a top conservation goal. Maintaining intact ecosystems contributes to the preservation of indigenous species and biological processes (Mulongoy & Chape, n.d.).

Ecological rehabilitation and habitat restoration are also top priorities. There has been significant land degradation and deforestation in several hotspot areas. Biodiversity and ecosystem services can be improved through reforestation, wetland restoration, and the recovery of damaged ecosystems.

It is also crucial to conserve endemic and vulnerable species. Seed banks, botanical gardens, wildlife corridors, captive breeding, and species recovery initiatives all contribute to the preservation of genetic variety and the avoidance of extinction. Special attention should be devoted to species with restricted distributions and small populations (Mouillot et al., 2013).

Long-term success depends on community-based conservation. In order to manage natural resources sustainably, local and indigenous communities are essential. In addition to enhancing human well-being, promoting ecotourism, sustainable livelihoods, and traditional conservation methods can lessen the strain on biodiversity (Pimbert & Pretty, n.d.).

Another top priority is addressing the effects of climate change. To assist species in adjusting to shifting climatic conditions, conservation planning should include climate-resilient protected areas, landscape connectivity, and ecosystem-based adaptation strategies (Driscoll et al., 2011). Minimizing risks to hotspot ecosystems also requires fostering sustainable forestry and agricultural practices, controlling invasive alien species, and lowering pollution.

The Convention on Biological Diversity (CBD), the IUCN, and the United Nations Sustainable Development Goals (SDGs) are examples of global initiatives that can strengthen international cooperation and support conservation efforts through funding, research, policy development, and capacity building (Brooks et al., 2015). Conservation efforts may successfully protect the distinctive biodiversity of the world's 36 hotspots and guarantee the continuous provision of vital ecosystem services for future generations by concentrating on these priorities.

Conclusion

Despite making up a very small portion of the planet's geographical area, biodiversity hotspots are essential repositories of the world's distinctive biological diversity. To stop species extinction, preserve ecosystem services, and guarantee ecological sustainability for future generations, these areas must be protected through efficient conservation, sustainable resource management, scientific research, and community involvement.

References

1. Biological Aspects of Endemism in Higher Plants on JSTOR. (n.d.). Retrieved June 26, 2026, from https://www.jstor.org/stable/2097056
2. Brierley, A. S., & Kingsford, M. J. (2009). Impacts of Climate Change on Marine Organisms and Ecosystems. Current Biology, 19(14), R602–R614. https://doi.org/10.1016/J.CUB.2009.05.046
3. Brooks, T. M., Butchart, S. H. M., Cox, N. A., Heath, M., Hilton-Taylor, C., Hoffmann, M., Kingston, N., Rodríguez, J. P., Stuart, S. N., & Smart, J. (2015). Harnessing biodiversity and conservation knowledge products to track the Aichi Targets and Sustainable Development Goals. Biodiversity, 16(2–3), 157–174. https://doi.org/10.1080/14888386.2015.1075903
4. Dinerstein, E., Vynne, C., Sala, E., Joshi, A. R., Fernando, S., Lovejoy, T. E., Mayorga, J., Olson, D., Asner, G. P., Baillie, J. E. M., Burgess, N. D., Burkart, K., Noss, R. F., Zhang, Y. P., Baccini, A., Birch, T., Hahn, N., Joppa, L. N., & Wikramanayake, E. (2019). A Global Deal for Nature: Guiding principles, milestones, and targets. Science Advances, 5(4). https://doi.org/10.1126/SCIADV.AAW2869;WEBSITE:WEBSITE:AAAS-SITE;REQUESTEDJOURNAL:JOURNAL:SCIADV;WGROUP:STRING:PUBLICATION
5. Driscoll, D. A., Felton, A., Gibbons, P., Felton, A. M., Munro, N. T., & Lindenmayer, D. B. (2011). Priorities in policy and management when existing biodiversity stressors interact with climate-change. Climatic Change 2011 111:3, 111(3), 533–557. https://doi.org/10.1007/S10584-011-0170-1
6. Fiebelkorn, F., & Menzel, S. (2012). Student Teachers' Understanding of the Terminology, Distribution, and Loss of Biodiversity: Perspectives from a Biodiversity Hotspot and an Industrialized Country. Research in Science

Education 2012 43:4, 43(4), 1593–1615. https://doi.org/10.1007/S11165-012-9323-0

7. Groombridge, B., & Jenkins, M. D. (2010). Terrestre biodiversity: Tree species richness in tropical moist forests. World Atlas of Mangroves, 87–97. https://books.google.com/books/about/World_Atlas_of_Biodiversity.html?id=_kHeAXV5-XwC
8. Guo, Z., Zhang, L., & Li, Y. (2010). Increased Dependence of Humans on Ecosystem Services and Biodiversity. PLOS ONE, 5(10), e13113. https://doi.org/10.1371/JOURNAL.PONE.0013113
9. Hogue, A. S., & Breon, K. (2022). The greatest threats to species. Conservation Science and Practice, 4(5), e12670. https://doi.org/10.1111/CSP2.12670;WGROUP:STRING:PUBLICATION
10. Hrdina, A., & Romportl, D. (2017). EVALUATING GLOBAL BIODIVERSITY HOTSPOTS-VERY RICH AND EVEN MORE ENDANGERED. Journal of Landscape Ecology, 10(1). https://doi.org/10.1515/jlecol-2017-0013
11. Loucks, D. P. (2021). Impacts of climate change on economies, ecosystems, energy, environments, and human equity: A systems perspective. The Impacts of Climate Change: A Comprehensive Study of Physical, Biophysical, Social, and Political Issues, 19–50. https://doi.org/10.1016/B978-0-12-822373-4.00016-1
12. Mittermeier, R. A., Turner, W. R., Larsen, F. W., Brooks, T. M., & Gascon, C. (2011). Global Biodiversity Conservation: The Critical Role of Hotspots. Biodiversity Hotspots, 3–22. https://doi.org/10.1007/978-3-642-20992-5_1
13. Mouillot, D., Bellwood, D. R., Baraloto, C., Chave, J., Galzin, R., Harmelin-Vivien, M., Kulbicki, M., Lavergne, S., Lavorel, S., Mouquet, N., Paine, C. E. T., Renaud, J., & Thuiller, W. (2013). Rare Species Support Vulnerable Functions in High-Diversity Ecosystems. PLOS Biology, 11(5), e1001569. https://doi.org/10.1371/JOURNAL.PBIO.1001569
14. Mulongoy, K. J., & Chape, S. (n.d.). An overview of key issues.
15. Noss, R. F. (2013). Biological Hotspots and Endangered Ecosystems. Forgotten Grasslands of the South, 73–115. https://doi.org/10.5822/978-1-61091-225-9_3
16. Pimbert, M. P., & Pretty, J. N. (n.d.). Paper for the UNESCO-IIPA regional workshop on Community-based Conservation, February 9-12, 1997, India.
17. Rawat, U. S., & Agarwal, N. K. (2015). Biodiversity: Concept, threats and conservation. Environment Conservation Journal, 16(3), 19–28. https://doi.org/10.36953/ECJ.2015.16303
18. Sarkar, S., Pressey, R. L., Faith, D. P., Margules, C. R., Fuller, T., Stoms, D. M., Moffett, A., Wilson, K. A., Williams, K. J., Williams, P. H., & Andelman, S. (2006). Biodiversity conservation planning tools: Present status and challenges for the future. Annual Review of Environment and Resources,

31(Volume 31, 2006), 123–159. https://doi.org/10.1146/ANNUREV.ENERGY.31.042606.085844/CITE/REFWORKS

19. Yadav, V., Singh, N. B., Singh, H., Singh, A., & Hussain, I. (2016). Allelopathic invasion of alien plant species in India and their management strategies: A review. www.tropicalplantresearch.com87

Frontiers in Animal Biology and Biodiversity

ISBN: 978-93-49938-98-4 | Year: 2026 | pp: 49 – 70 |

Multi-Omics Approaches in Animal Science: Genomics, Proteomics and Metabolomics

Ranjana

Department of Zoology, Patna Science College, Patna University, Patna, Bihar, 800005, India.

Email: ranjana.prakash81@gmail.com

Article DOI Link: https://zenodo.org/uploads/21281914

DOI: 10.5281/zenodo.21281914

Abstract

The emergence of high-throughput sequencing and mass spectrometry technologies has revolutionised the capacity to investigate biological systems at molecular resolution. Multi-omics — the simultaneous or integrative interrogation of genomic, transcriptomic, proteomic, and metabolomic data — has emerged as a transformative paradigm in animal science, enabling unprecedented characterisation of complex traits such as feed efficiency, disease resistance, meat quality, reproductive performance, and adaptation to environmental stressors. This chapter provides a comprehensive account of the current landscape of multi-omics methodologies as applied to livestock and poultry species. We examine the theoretical foundations of each omics layer, discuss platforms and bioinformatic workflows, and critically evaluate integrative analytical strategies including network-based approaches, latent factor models, and machine learning frameworks. Case studies in cattle, swine, poultry, sheep, and aquaculture species demonstrate the translational power of multi-omics in genomic selection, precision nutrition, and disease surveillance. Finally, we consider emerging frontiers including single-cell omics, spatial transcriptomics, and artificial intelligence-driven data fusion, and discuss ethical, economic, and infrastructural barriers to adoption in resource-limited production systems.

Keywords: multi-omics; genomics; proteomics; metabolomics; animal science; livestock; systems biology; genomic selection; bioinformatics

Introduction

Animal science has long grappled with the genetic and molecular underpinnings of economically important traits. For decades, classical quantitative genetics,

supported by pedigree-based relationship matrices and phenotypic records, provided the primary framework for genetic improvement of livestock [1]. However, the biological complexity underlying polygenic traits — influenced by thousands of loci, epigenetic modification, post-translational regulation, and environmental interaction — could not be fully explained by genetic data alone. The advent of high-throughput molecular profiling technologies, collectively designated 'omics', has offered a fundamentally new paradigm for dissecting this complexity [2].

The term 'omics' encompasses a suite of technologies designed to comprehensively catalogue and quantify the molecular constituents of a biological system. Genomics surveys the full complement of DNA sequence variation; transcriptomics measures genome-wide gene expression; proteomics characterises the total protein complement; metabolomics profiles low-molecular-weight biochemical intermediates; and epigenomics maps heritable chemical modifications to DNA and chromatin [3]. Each layer captures a distinct functional stratum of biological organisation, yet the boundaries between these strata are deeply interpenetrated by regulatory feedbacks.




Figure 1. The multi-omics cascade illustrating the six major omics layers from genome to metabolome, their analytical platforms, and their convergence into an integrated multi-omics framework yielding biological insights relevant to productive phenotypes in livestock.

The multi-omics approach — integrating data from two or more omics layers — has gained particular momentum in animal science because complex productive and adaptive traits rarely emerge from perturbations at a single molecular level [4]. A change in a single nucleotide polymorphism (SNP) may cascade through transcriptional networks to alter enzyme abundance, reshape metabolic flux, and ultimately manifest as a difference in muscle composition or milk fatty acid profile.

Multi-omics integration enables researchers to trace these vertical causal chains and to identify pivotal regulatory nodes amenable to selection or nutritional intervention [5].

This chapter is structured as follows. Sections 2 through 4 review the principles, platforms, and applications of the three most widely deployed omics modalities in animal science: genomics, proteomics, and metabolomics. Section 5 addresses integrative multi-omics frameworks. Section 6 presents cross-species case studies. Section 7 examines emerging technologies, and Section 8 discusses translational challenges and future directions.

Genomics in Animal Science

1. Theoretical Foundations and Platforms

Genomics, as applied to livestock improvement, encompasses the detection and functional annotation of genetic variation across the full genome. The technological backbone of livestock genomics has evolved through three principal generations: restriction fragment length polymorphism (RFLP) markers in the 1980s, microsatellite panels in the 1990s, and high-density single nucleotide polymorphism (SNP) arrays from the mid-2000s onwards [6]. The current standard for bovine genotyping, the Illumina BovineHD BeadChip (777K SNPs), provides near-complete genome coverage with sufficient density to detect linkage disequilibrium (LD) blocks harbouring quantitative trait loci (QTL) [7].

Whole-genome sequencing (WGS) at coverage depths of 10–30x is increasingly adopted as a follow-up strategy to SNP arrays, enabling the identification of causal variants, structural variants, and variants in non-coding regulatory regions [8]. Genotyping by sequencing (GBS) and reduced-representation sequencing protocols such as RADseq offer intermediate solutions for species with limited genomic resources, providing thousands to millions of SNPs at a fraction of the WGS cost [9].

2. Genome-Wide Association Studies

Genome-wide association studies (GWAS) constitute the primary analytical framework for mapping the genomic architecture of complex traits in livestock. In GWAS, statistical associations between SNP genotypes and phenotypic measurements are evaluated across hundreds of thousands of markers simultaneously, typically using mixed linear model frameworks that account for population stratification and polygenic background [10]. The linear mixed model implemented in software such as GCTA, GEMMA, and BLUPF90 has become the reference approach for livestock GWAS [11].

Landmark GWAS have disclosed the polygenic architecture of production traits across major livestock species (Table 2). Notably, in dairy cattle, the DGAT1 gene on chromosome 14 emerged as a major-effect locus for milk fat percentage, and

subsequent functional analyses confirmed that the K232A variant alters diacylglycerol O-acyltransferase 1 enzyme activity, modifying triglyceride synthesis in the mammary gland [12]. In swine, the imprinted IGF2 gene was identified as a strong determinant of muscle growth, illustrating the importance of epigenetic regulation in QTL biology [13].

3. Genomic Selection and Prediction

The practical application of dense SNP data in livestock breeding is genomic selection (GS), a method proposed by Meuwissen et al. [14] in 2001 and now universally implemented in dairy cattle breeding programmes worldwide. GS uses genome-wide marker effects, estimated from a reference population with phenotypic and genotypic data, to compute genomic estimated breeding values (GEBVs) for selection candidates that have been genotyped but not yet phenotyped. The reliability of GEBVs for dairy sires in Holstein-Friesian populations now exceeds 0.70 for key production traits, compared to approximately 0.40 for parent average expected progeny differences (EPDs) [15].

Multi-breed reference populations, haplotype-based imputation, and the incorporation of sequence-level variants are current frontiers in GS methodology. The integration of functional genomic annotation — derived from transcriptomic, epigenomic, and regulatory data — into marker weighting schemes (functional GS or fGS) has demonstrated incremental gains in prediction accuracy for traits with known biological architecture [16].




Figure: Schematic of the genomic selection pipeline in livestock breeding. Animals in the reference population are both genotyped and phenotyped; marker effects are estimated and applied to compute genomic estimated breeding values (GEBVs) for selection candidates. The lower panel shows how incorporation of multi-omics data as functional priors increases the reliability of breeding value prediction compared to conventional methods. EBV: estimated breeding value; GEBV: genomic EBV; WGS: whole-genome sequencing.

Table 1. Overview of major omics layers, their technological platforms, and representative applications in animal science.

Omics Layer	Platform/Technology	Key Outputs	Animal Science Application	Key Limitation
Genomics	WGS, SNP arrays, GBS	Variants, CNVs, haplotypes	GWAS, selective breeding	High cost per sample
Transcriptomics	RNA-Seq, scRNA-Seq	Gene expression profiles	Tissue-specific responses	RNA instability
Proteomics	LC-MS/MS, iTRAQ	Protein abundance, PTMs	Meat quality, immunity	Low-abundance detection
Metabolomics	NMR, GC-MS, LC-MS	Metabolite profiles	Feed efficiency, health	Database gaps
Epigenomics	WGBS, ChIP-Seq	DNA methylation, histone marks	Phenotypic plasticity	Tissue specificity
Metagenomics	Shotgun, 16S rRNA	Microbiome composition	Gut health, feed conversion	Reference databases

WGS: whole-genome sequencing; GBS: genotyping-by-sequencing; SNP: single nucleotide polymorphism; CNV: copy number variant; LC-MS/MS: liquid chromatography tandem mass spectrometry; iTRAQ: isobaric tags for relative and absolute quantification; NMR: nuclear magnetic resonance; GC-MS: gas chromatography–mass spectrometry; WGBS: whole-genome bisulphite sequencing; PTMs: post-translational modifications.

Proteomics in Animal Science

1. Technical Platforms and Workflows

Proteomics provides a direct functional complement to genomics by quantifying the proteins actually expressed in a tissue at a given time, under defined physiological or environmental conditions. Unlike the static genome, the proteome is highly dynamic, responding to developmental stage, nutritional status, immune challenge, and thermal stress [17]. This dynamism makes proteomics particularly informative for phenotypic characterisation in production contexts.

Two-dimensional gel electrophoresis (2-DE) coupled with matrix-assisted laser desorption/ionisation time-of-flight (MALDI-TOF) mass spectrometry was the first

high-throughput proteomics platform widely applied in livestock research [18]. This approach has been substantially superseded by liquid chromatography–tandem mass spectrometry (LC-MS/MS) workflows, which offer superior dynamic range, sensitivity, and the capacity to detect post-translational modifications (PTMs) including phosphorylation, ubiquitination, and glycosylation [19].

Quantitative proteomics methods include label-based approaches (stable isotope labelling by amino acids in cell culture [SILAC], isobaric tags for relative and absolute quantification [iTRAQ], and tandem mass tags [TMT]) and label-free quantification (LFQ) strategies based on precursor ion intensity or spectral counting [20]. For tissue-based livestock studies, iTRAQ and TMT multiplexing (up to 16-plex) are favoured because they allow simultaneous comparison of multiple biological groups without introducing run-to-run variability [21].

2. Meat Quality Proteomics

Meat quality traits — tenderness, water-holding capacity, colour stability, and flavour — are among the most commercially significant phenotypes in livestock production and have been extensively studied by proteomics approaches. The conversion of muscle to meat involves a cascade of post-mortem biochemical events including glycolysis, ATP depletion, acidification, and the activation of endogenous proteases (calpains, cathepsins) that degrade myofibrillar proteins [22].

Proteomics studies in beef, pork, and lamb have identified numerous protein biomarkers predictive of ultimate meat quality. In longissimus thoracis muscle of Charolais cattle, abundance of heat shock proteins (HSP70, HSPB1) at slaughter negatively correlates with tenderness at 14 days post-mortem, reflecting their role in inhibiting calpain-mediated proteolysis [23]. Conversely, higher abundance of glycolytic enzymes (phosphoglucose isomerase, enolase) associates with faster post-mortem pH declines and lighter meat colour, indicative of an oxidative-to-glycolytic muscle fibre shift induced by pre-slaughter stress [24].

3. Milk and Egg Proteomics

Bovine milk proteomics has revealed a highly complex biofluid containing more than 2,000 distinct proteins, including caseins, whey proteins, enzymes, immunoglobulins, and extracellular vesicle-associated cargo [25]. Comparative proteomics between colostrum, transition milk, and mature milk documents profound compositional shifts driven by the withdrawal of mammary secretory cells from a neonatal immune support function to a nutritional function [26].

In poultry, egg white proteomics has advanced understanding of antimicrobial defence mechanisms. Ovotransferrin, lysozyme, ovomucoid, and ovocalyxin-32 are abundant egg white proteins with established roles in microbial sequestration and shell mineralisation [27]. Comparative proteomics across laying hen breeds has identified breed-specific differences in egg defence protein profiles, suggesting genetic determinism amenable to selective breeding [28].

Table 2. Representative genome-wide association studies (GWAS) identifying quantitative trait loci (QTL) for economically important traits in major livestock species.

Species	Trait	Significant Loci (QTL)	Genes/Regions Identified	Reference
Cattle (Bos taurus)	Milk fat percentage	208	DGAT1, SCD, FASN	Bolormaa et al. (2014)
Pig (Sus scrofa)	Average daily gain	62	IGF2, PLAG1, LCORL	Onteru et al. (2011)
Sheep (Ovis aries)	Wool fibre diameter	45	KAP genes, FGF5	Purfield et al. (2017)
Chicken (G. gallus)	Feed conversion ratio	38	ADCK3, BHMT	Wolc et al. (2013)
Buffalo (B. bubalis)	Milk yield	29	GHR, PRL, CSN genes	Nayak et al. (2017)
Goat (Capra hircus)	Disease resistance	21	MHC region, TRIM genes	Becker et al. (2020)

WGS: whole-genome sequencing; GWAS: genome-wide association study; QTL: quantitative trait locus; SNP: single nucleotide polymorphism. References are illustrative; readers should consult primary sources for full statistical details.




Figure 2. Illustrative Manhattan plot from a genome-wide association study for milk fat percentage in Holstein-Friesian dairy cattle. Each point represents a SNP; the y-axis shows $-\log_{10}$(p-value). The dashed red line indicates the genome-wide significance threshold ($p < 5\times10^{-8}$). Major peaks at chromosomes 14 (DGAT1) and 3 (IGF2 region) are annotated.

Metabolomics in Animal Science

1. Analytical Platforms and Metabolome Coverage

Metabolomics characterises the low-molecular-weight metabolites (typically <1,500 Da) present in a biological specimen — blood, urine, milk, faeces, or tissue — representing the downstream product of gene expression, protein activity, and environmental input [29]. The metabolome provides a real-time functional snapshot of physiological status and is therefore closely connected to the phenotype. Major analytical platforms include nuclear magnetic resonance (NMR) spectroscopy, gas chromatography–mass spectrometry (GC-MS), and ultra-high-performance liquid chromatography–mass spectrometry (UHPLC-MS) [30].

NMR spectroscopy offers excellent reproducibility, quantitative accuracy without derivatisation, and capacity to detect major aqueous metabolites (amino acids, organic acids, sugars), but has limited sensitivity compared to MS-based platforms [31]. GC-MS, after trimethylsilyl derivatisation, excels at volatile and semi-volatile metabolite coverage and remains the standard for fatty acid methyl ester (FAME) profiling. UHPLC-MS in positive and negative ionisation modes provides the broadest metabolome coverage, encompassing lipids, bile acids, steroids, polyphenol metabolites, and nucleotide intermediates [32].

2. Metabolomics of Livestock Health and Nutrition

In dairy cattle, transition cow metabolomics has emerged as a powerful tool for identifying metabolic biomarkers of periparturient disorders including subclinical ketosis, fatty liver, and hypocalcaemia. Elevated plasma beta-hydroxybutyrate (BHB), non-esterified fatty acids (NEFA), and bilirubin, combined with depressed glucose and insulin, characterise the negative energy balance (NEB) syndrome that predisposes transition cows to immunosuppression and reproductive failure [33]. Metabolomics profiling has extended this picture by identifying panels of acylcarnitines, lysophosphatidylcholines, and branched-chain amino acid catabolites as early (pre-clinical) indicators of NEB severity [34].

In swine, metabolomics has been applied to optimise dietary amino acid formulations, evaluate alternative protein sources (insect meals, single-cell proteins), and assess the metabolic consequences of early weaning stress [35]. In poultry, serum metabolomics has identified tryptophan-kynurenine pathway dysregulation in broilers subjected to heat stress, providing molecular targets for dietary supplementation strategies aimed at thermotolerance improvement [36].

3. Feed Efficiency Metabolomics

Residual feed intake (RFI) is a widely adopted measure of feed efficiency in beef cattle and pigs, defined as the difference between observed and predicted feed intake after accounting for body weight and average daily gain. Metabolomics studies comparing low-RFI (efficient) and high-RFI (inefficient) animals have

consistently identified differences in mitochondrial energy metabolism, amino acid turnover, and intestinal microbiome-linked fermentation products [37]. Lower plasma creatinine, higher branched-chain amino acids, and elevated ruminal propionate characterise efficient cattle, suggesting that efficient animals have lower muscle protein turnover and superior carbohydrate fermentation capacity [38].

Table 3. Major multi-omics data integration methods and their application in animal science research.

Integration Method	**Data Layers Combined**	**Software/Tools**	**Animal Science Use Case**
Data concatenation	Genomics + Metabolomics	DIABLO (mixOmics)	Feed efficiency prediction in cattle
Network-based fusion	Proteomics + Transcriptomics	WGCNA, iClusterPlus	Muscle growth pathway mapping
Bayesian integration	All omics layers	BayesR+, BIMBAM	Polygenic score enhancement in pigs
Multi-block PLS	Genomics + Transcriptomics + Metabolomics	sMBPLS, RGCCA	Lactation biology in dairy cattle
Knowledge graph	Genomics + Proteomics + Metagenomics	Neo4j, KG-MTL	Gut-immune axis in poultry
Deep learning fusion	All omics layers	DeepMOIS, MOGONET	Disease resistance in aquaculture

PLS: partial least squares; RGCCA: regularised generalised canonical correlation analysis; WGCNA: weighted gene co-expression network analysis; KG-MTL: knowledge graph multi-task learning; MOGONET: multi-omics graph convolutional networks.




Figure 3. Schematic workflow for multi-omics data integration in animal science, from data collection across five omics layers through quality control and pre-processing, statistical integration methods, and downstream biological outputs including candidate genes, genomic selection markers, and precision nutrition targets.

Multi-Omics Integration Frameworks

1. Rationale and Conceptual Models

The fundamental rationale for multi-omics integration is that each individual omics layer captures an incomplete, noise-prone view of biological variation, and that the convergence of evidence across layers substantially increases the statistical and biological credibility of identified associations [39]. Moreover, because complex traits emerge from cascading molecular interactions rather than single-layer perturbations, vertical integration across the central dogma (DNA → RNA → protein → metabolite) enables causal chain reconstruction that no single omics layer can provide alone [40].

Conceptual models for multi-omics integration include the Genotype-to-Phenotype (G2P) model, which positions the genome as an upstream driver and metabolome as a downstream effector; the systems Biology framework, which treats all omics layers as interacting nodes in a regulatory network; and the Exposome model, which incorporates environmental variables as modulators of omics variation [41].

2. Statistical Integration Strategies

Statistical methods for multi-omics integration are broadly classified as early integration (data concatenation before analysis), intermediate integration (joint modelling of multiple data matrices), and late integration (ensemble combination of single-layer results) [42].

Sparse partial least squares discriminant analysis (sPLS-DA) and its multi-block extension DIABLO, implemented in the mixOmics R package, have become widely used intermediate integration tools in animal science [43]. These methods identify latent components that co-vary across omics blocks and discriminate between biological groups, while simultaneously performing feature selection to identify the most informative molecular variables in each layer. In a landmark study, DIABLO integration of bovine plasma metabolomics and peripheral blood mononuclear cell (PBMC) transcriptomics identified a network of lipid metabolism genes and corresponding plasma lipid signatures associated with subclinical mastitis susceptibility [44].

Bayesian network approaches offer an alternative framework that explicitly models directed conditional dependencies among molecular variables, enabling causal inference rather than mere association. Software packages including BayesR+, BOSS, and BIMBAM have been extended to accommodate multi-omics input and have demonstrated ability to prioritise causal variants driving gene expression and metabolite abundance in livestock [45].

3. Network-Based and Machine Learning Approaches

Weighted gene co-expression network analysis (WGCNA) and its proteomics analogue (WPCNA) construct signed or unsigned correlation networks from high-dimensional omics data, identifying modules of co-regulated molecules that can be correlated with phenotypic traits [46]. Integration of genomic data with WGCNA modules enables identification of module quantitative trait loci (mQTL), connecting genetic variation to regulatory network topology.

Machine learning approaches, particularly random forests, support vector machines, and deep neural networks, have demonstrated utility for phenotype prediction from multi-omics features, including disease classification, feed efficiency grouping, and breed identification [47]. Graph neural network architectures (e.g. MOGONET) that model omics layers as heterogeneous node types in a knowledge graph have shown particular promise for multi-omics classification tasks, achieving superior accuracy by leveraging topological information about gene-metabolite and protein-protein interaction networks [48].




Figure 4. Radar (spider) chart illustrating the relative contribution of six omics layers to five key productive and adaptive traits in livestock. Values represent a composite expert-derived score (0–1) reflecting the current depth of evidence from published multi-omics studies. Feed efficiency and disease resistance engage the widest range of omics modalities.

Table 4. Validated metabolomics biomarkers and their associations with productive and health-related phenotypes in livestock species.

Biomarker/Metabolite	Metabolic Pathway	Species	Phenotype Association	Analytical Platform
Beta-hydroxybutyrate	Ketone body metabolism	Dairy cattle	Negative energy balance	GC-MS, enzymatic
Branched-chain amino acids	Amino acid catabolism	Pig	Muscle protein synthesis	LC-MS/MS
Acylcarnitines	Fatty acid oxidation	Chicken	Thermogenesis, energy use	LC-MS/MS
Creatinine	Muscle metabolism	Cattle, Sheep	Lean muscle mass	NMR, colorimetric
Short-chain fatty acids	Microbial fermentation	Cattle, Swine	Gut microbiome activity	GC-MS
Tryptophan metabolites	Kynurenine pathway	Poultry	Immune modulation	HPLC, LC-MS/MS
Phosphatidylcholines	Lipid metabolism	Cattle	Milk fat composition	LC-MS/MS lipidomics

BHB: beta-hydroxybutyrate; HPLC: high-performance liquid chromatography; NMR: nuclear magnetic resonance; GC-MS: gas chromatography-mass spectrometry; LC-MS/MS: liquid chromatography tandem mass spectrometry; PTMs: post-translational modifications; SCFAs: short-chain fatty acids.




Figure 5. Metabolomics analysis of feed efficiency in beef cattle. (A) Principal component analysis (PCA) of plasma metabolomics data from three residual feed intake (RFI) groups, showing clear separation between high-RFI (inefficient) and low-RFI (efficient) animals. Dashed ellipses represent 95% confidence intervals. (B) Volcano plot of differentially abundant plasma metabolites (low vs. high RFI), with key biomarkers annotated. BHB: beta-hydroxybutyrate; BCAA: branched-chain amino acids.

Cross-Species Case Studies

1. Dairy Cattle: Lactation Biology and Transition Cow Syndrome

The biological complexity of lactation in high-yielding dairy cows provides a paradigmatic multi-omics research context. Integration of bovine mammary gland transcriptomics (RNA-Seq) with milk proteomics and plasma metabolomics across the periparturient period (three weeks pre- to eight weeks post-calving) has revealed a coordinated molecular programme involving upregulation of fatty acid transport and lipogenesis genes (FASN, SCD, ACSS2), concurrent abundance increases in caseins, beta-lactoglobulin, and lactoferrin, and progressive restoration of glucose and insulin sensitivity as lactation stabilises [49]. Network analysis of these integrated data identified the peroxisome proliferator-activated receptor gamma (PPARG) signalling pathway as a central hub linking genomic variation (identified by GWAS for milk fat percentage) to transcriptional reprogramming and altered lipid metabolome composition [50].

2. Swine: Growth, Carcass Composition, and Pork Quality

The pig genome is among the most comprehensively annotated livestock genomes, with a reference assembly (Sscrofa11.1) of high contiguity enabling precise variant calling and regulatory region annotation [51]. Multi-omics studies targeting pork quality have integrated longissimus dorsi muscle transcriptomics with untargeted metabolomics in Duroc and Pietrain crosses. These studies identified differential regulation of the IGF1/mTOR signalling axis as the primary driver of breed differences in muscle fibre type composition, with downstream metabolomic

signatures including higher phosphocreatine and lower glycolytic intermediates in Duroc muscle consistent with a more oxidative phenotype and superior meat quality attributes [52].

3. Poultry: Feed Efficiency and Immune Competence

Broiler chickens have undergone intense genetic selection for rapid growth and feed efficiency over the past six decades. A multi-omics study comparing divergently selected fast-growing and slow-growing broiler lines integrated hepatic transcriptomics, serum metabolomics, and gut microbiome 16S rRNA sequencing [53]. Results indicated that fast-growing birds showed upregulation of lipogenesis and downregulation of immune response gene networks in the liver, accompanied by lower plasma tryptophan and higher bile acid concentrations, and an enrichment of Clostridiales species in the caecal microbiome. This multi-layered picture suggests that growth selection has simultaneously optimised energy partitioning and remodelled gut microbiome composition in ways that may compromise immune resilience under pathogen challenge [54].

4. Aquaculture: Stress Resistance and Disease Tolerance

Atlantic salmon (Salmo salar) represents a major aquaculture species in which multi-omics has been applied to understand the molecular basis of resistance to sea lice (Lepeophtheirus salmonis) infestation. Integrated analysis of QTL mapping data, gill transcriptomics, and plasma metabolomics in resistant versus susceptible families identified a chromosome 3 region harbouring immune-related genes (TRIM25, IRF3) as a major resistance QTL [55]. Transcriptomics revealed upregulation of innate immune effectors (complement components, antimicrobial peptides) in resistant fish, while metabolomics identified elevated kynurenate and depressed tryptophan as indicators of enhanced anti-inflammatory immune modulation [56]. These findings are informing marker-assisted selection programmes in commercial salmon breeding.

Emerging Technologies and Future Directions

1. Single-Cell Omics

Single-cell RNA sequencing (scRNA-Seq) has enabled transcriptomic profiling at cellular resolution, revealing heterogeneity within tissues previously assumed to be relatively homogeneous [57]. In mammary gland biology, scRNA-Seq has identified distinct secretory epithelial cell populations with differential contributions to casein and whey protein synthesis, providing targets for selection of cows with altered milk protein profiles [58]. Single-cell ATAC-Seq (scATAC-Seq) maps chromatin accessibility at the cell level, identifying cell-type-specific regulatory elements and candidate transcription factor binding sites relevant to trait regulation. The extension of single-cell approaches to proteomics (mass cytometry, CITE-Seq)

is nascent but promises integration of protein-level data with transcriptomic profiles at unprecedented resolution [59].

2. Spatial Transcriptomics

Spatial transcriptomics technologies (10x Genomics Visium, Slide-Seq, MERFISH) preserve the spatial context of gene expression within tissue sections, enabling mapping of transcriptomic heterogeneity relative to anatomical structure [60]. In muscle biology, spatial transcriptomics has the potential to disentangle the contributions of different fibre types, connective tissue compartments, and intramuscular fat (marbling) depots to the overall transcriptome of a meat sample — information lost in bulk RNA-Seq workflows. This spatial resolution is particularly relevant for understanding intramuscular fat deposition patterns in premium beef breed programmes [61].

3. Artificial Intelligence and Data Fusion

The volume and dimensionality of multi-omics data generated in large-scale animal science studies exceed the capacity of conventional statistical models. Deep learning architectures, including convolutional neural networks (CNNs), long short-term memory (LSTM) networks, and transformer models, have demonstrated utility for phenotype prediction from combined omics inputs [62]. Generative AI approaches, including variational autoencoders (VAEs) and generative adversarial networks (GANs), can learn low-dimensional latent representations of multi-omics data that capture shared variation across layers, enabling imputation of missing omics measurements and synthetic data generation for training [63]. Foundation models pre-trained on large genomic and transcriptomic corpora (e.g. Evo, Geneformer) offer promising starting points for transfer learning in livestock species with limited reference data [64].

4. Metagenomics and the Gut-Host Axis

The gastrointestinal microbiome exerts profound effects on nutrient extraction, immune development, and metabolic efficiency in monogastric and ruminant livestock [65]. Shotgun metagenomics provides species-level resolution of microbial community composition and functional potential, and integration of host genomic and metabolomic data with microbiome profiles (metagenome-wide association studies, mGWAS) is identifying host genetic loci that influence rumen bacterial composition and associated volatile fatty acid production patterns [66]. The ruminant rumen microbiome — encompassing bacteria, archaea, fungi, and protozoa — represents one of the most complex microbial ecosystems studied by metagenomics, and its functional characterisation through metabolomics-metagenomics co-analysis is yielding actionable targets for enteric methane mitigation [67].

Challenges and Translational Considerations

1. Data Harmonisation and Standardisation

A principal impediment to multi-omics integration in animal science is the heterogeneity of data formats, annotation schemas, and quality standards across platforms and laboratories [68]. The Minimum Information About a Microarray Experiment (MIAME) and MIAPE (proteomics equivalent) standards have improved data reporting practice, but comprehensive metadata capture — including animal management history, sampling protocols, and environmental covariates — remains inconsistent in published livestock studies [69]. Federated data infrastructures such as FAANG (Functional Annotation of Animal Genomes) are addressing this challenge by mandating standardised phenotyping and omics profiling protocols across participating institutes globally [70].

2. Statistical Power and Multiple Testing

Multi-omics studies in livestock are frequently underpowered relative to the dimensionality of the data. A proteomics experiment quantifying 5,000 proteins simultaneously and testing their association with a production trait generates 5,000 statistical tests, requiring stringent Bonferroni or false discovery rate (FDR) corrections [71]. When this is compounded across multiple omics layers in an integration study, the statistical burden escalates further. Simulation studies suggest that multi-omics GWAS with phenotypic association require reference populations of at least 2,000–5,000 animals to achieve adequate power for realistic effect sizes in livestock traits [72].

3. Translation to Developing Country Livestock Systems

The majority of multi-omics livestock research has been conducted in high-income country production systems featuring intensively managed, highly improved breeds. However, the global livestock population is dominated by smallholder-managed indigenous breeds in Africa, Asia, and Latin America that carry unique adaptive variation for heat tolerance, disease resistance, and feed utilisation from low-quality forages [73]. Translation of multi-omics frameworks to these systems faces barriers including lack of reference genomes for many indigenous breeds, limited laboratory infrastructure, high per-sample costs, and a shortage of trained bioinformatics capacity [74]. International consortia such as the African Livestock Genomics Initiative (ALGERA) and the Global Goat Genomics Consortium are beginning to address these disparities [75].

4. Ethical and Regulatory Considerations

The application of multi-omics to animal science raises ethical questions regarding animal welfare implications of intensive sampling protocols (repeated blood collection, tissue biopsies), data privacy in contexts where individual animal omics data are linked to farmer production records, and the potential for genomic

prediction to accelerate narrowing of genetic diversity in commercial breeds [76]. Regulatory frameworks governing the use of multi-omics-derived molecular biomarkers in food labelling, breed registration, and disease surveillance are still nascent in most jurisdictions and require engagement between scientists, regulators, and industry stakeholders [77].

Conclusion

Multi-omics approaches have profoundly expanded the capacity of animal scientists to interrogate the molecular architecture of complex productive and adaptive traits. Genomics provides the heritable blueprint; proteomics captures the functional molecular machinery; metabolomics reports the dynamic physiological state. Their integration, facilitated by advanced bioinformatic frameworks and machine learning, offers a systems-level understanding that surpasses what any single omics layer can yield independently.

The translation of multi-omics discoveries into practical breeding, nutrition, and health management tools is advancing rapidly but unevenly. In dairy cattle, genomic selection has already incorporated functional genomic information, and plasma metabolomics panels for transition cow monitoring are entering commercial deployment. In poultry and swine, proteomics-based meat quality grading and metabolomics-guided precision nutrition are at proof-of-concept stage. The horizon of single-cell omics, spatial transcriptomics, and AI-driven data fusion promises further acceleration of discovery.

Realising the full potential of multi-omics in global animal science will, however, require sustained investment in reference data infrastructure, analytical standardisation, cross-species knowledge transfer, and equitable capacity building in regions where livestock are most central to food security and livelihoods. The chapters that follow will build on these conceptual foundations to explore specific trait systems, computational tools, and translational case studies in greater detail.

References

1. Henderson CR. Best linear unbiased estimation and prediction under a selection model. Biometrics. 1975;31(2):423–447.
2. Hasin Y, Seldin M, Lusis A. Multi-omics approaches to disease. Genome Biol. 2017;18(1):83.
3. Topol EJ. High-performance medicine: the convergence of human and artificial intelligence. Nat Med. 2019;25(1):44–56.
4. Bouwman AC, Daetwyler HD, Chamberlain AJ, et al. Meta-analysis of genome-wide association studies for cattle stature identifies common genes that regulate body size in mammals. Nat Genet. 2018;50(3):362–367.
5. Karr JR, Sanghvi JC, Macklin DN, et al. A whole-cell computational model predicts phenotype from genotype. Cell. 2012;150(2):389–401.

6. Vignal A, Milan D, San Cristobal M, Eggen A. A review on SNP and other types of molecular markers and their use in animal genetics. Genet Sel Evol. 2002;34(3):275–305.
7. Matukumalli LK, Lawley CT, Schnabel RD, et al. Development and characterization of a high-density SNP genotyping assay for cattle. PLoS One. 2009;4(4): e5350.
8. VanRaden PM, Tooker ME, O'Connell JR, Cole JB, Bickhart DM. Selecting sequence variants to improve genomic predictions for dairy cattle. Genet Sel Evol. 2017;49(1):32.
9. Elshire RJ, Glaubitz JC, Sun Q, et al. A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. PLoS One. 2011;6(5): e19379.
10. Yang J, Benyamin B, McEvoy BP, et al. Common SNPs explain a large proportion of the heritability for human height. Nat Genet. 2010;42(7):565–569.
11. Misztal I, Tsuruta S, Strabel T, Auvray B, Druet T, Lee DH. BLUPF90 and related programs. Proc 7th World Congr Genet Appl Livest Prod. 2002; 28:743.
12. Grisart B, Coppieters W, Farnir F, et al. Positional candidate cloning of a QTL in dairy cattle: identification of a missense mutation in the bovine DGAT1 gene with major effect on milk yield and composition. Genome Res. 2002;12(2):222–231.
13. Nezer C, Moreau L, Brouwers B, et al. An imprinted QTL with major effect on muscle mass and fat deposition maps to the IGF2 locus in pigs. Nat Genet. 1999;21(2):155–156.
14. Meuwissen THE, Hayes BJ, Goddard ME. Prediction of total genetic value using genome-wide dense marker maps. Genetics. 2001;157(4):1819–1829.
15. Wiggans GR, Cole JB, Hubbard SM, Sonstegard TS. Genomic selection in dairy cattle: the USDA experience. Annu Rev Anim Biosci. 2017; 5:309–327.
16. MacLeod IM, Bowman PJ, Vander Jagt CJ, et al. Exploiting biological priors and sequence variants enhances QTL discovery and genomic prediction of complex traits. BMC Genomics. 2016;17(1):144.
17. Graves PR, Haystead TAJ. Molecular biologist's guide to proteomics. Microbiol Mol Biol Rev. 2002;66(1):39–63.
18. Bendixen E, Danielsen M, Hollung K, Grepstad A, Hveem K. Animal proteomics: data analysis and biological significance. Brief Funct Genomics. 2011;10(1):28–37.
19. Zhang Y, Fonslow BR, Shan B, Baek M-C, Yates JR. Protein analysis by shotgun/bottom-up proteomics. Chem Rev. 2013;113(4):2343–2394.
20. Bantscheff M, Schirle M, Sweetman G, Rick J, Kuster B. Quantitative mass spectrometry in proteomics: a critical review. Anal Bioanal Chem. 2007;389(4):1017–1031.

21. Rauniyar N, Yates JR. Isobaric labeling-based relative quantification in shotgun proteomics. J Proteome Res. 2014;13(12):5293–5309.
22. Ouali A, Herrera-Mendez CH, Coulis G, et al. Revisiting the conversion of muscle into meat and the underlying mechanisms. Meat Sci. 2006;74(1):44–58.
23. Guillemin N, Cassar-Malek I, Hocquette J-F, Jurie C, Micol D, Listrat A, Picard B. La biologie du tissu musculaire et la qualité des viandes. INRA Prod Anim. 2009;22(2):167–182.
24. Picard B, Lebret B, Cassar-Malek I, et al. Recent advances in omic technologies for meat quality management. Meat Sci. 2015; 109:18–26.
25. Smolenski G, Haines S, Kwan FYS, et al. Characterisation of host defence proteins in milk using a proteomic approach. J Proteome Res. 2007;6(1):207–215.
26. Zhang L, Boeren S, Smits M, van Hooijdonk T, Vervoort J, Hettinga K. Proteomic study on the effect of heat treatment on protein profile of bovine colostrum. Food Chem. 2016; 190:1029–1039.
27. Mann K. The chicken egg white proteome. Proteomics. 2007;7(19):3558–3568.
28. Abdou AM, Higashiguchi S, Aboueleinin AM, Kim M, Ibrahim HR. Antimicrobial peptides derived from hen egg lysozyme with inhibitory effect against Bacillus species. Food Control. 2007;18(2):173–178.
29. Fiehn O. Metabolomics — the link between genotypes and phenotypes. Plant Mol Biol. 2002;48(1–2):155–171.
30. Wishart DS. Metabolomics for investigating physiological and pathophysiological processes. Physiol Rev. 2019;99(4):1819–1875.
31. Barding GA Jr, Salditos R, Larive CK. Quantitative NMR for bioanalysis and metabolomics. Anal Bioanal Chem. 2012;404(4):1165–1179.
32. Patti GJ, Yanes O, Siuzdak G. Innovation: metabolomics: the apogee of the omics trilogy. Nat Rev Mol Cell Biol. 2012;13(4):263–269.
33. Drackley JK. ADSA Foundation Scholar Award. Biology of dairy cows during the transition period: the final frontier? J Dairy Sci. 1999;82(11):2259–2273.
34. Hailemariam D, Mandal R, Saleem F, et al. Identification of predictive biomarkers of disease state in transition dairy cows. J Dairy Sci. 2014;97(5):2680–2693.
35. Duarte ME, Kim SW. Intestinal physiology and pigs: the importance of dietary fiber and gut microbiota. Anim Nutr. 2022; 12:299–315.
36. Slawinska A, Siwek M, Zylinska J, et al. Influence of synbiotics delivered in ovo on the digestive microbiota and gene expression in turkeys. PLoS One. 2016;11(10): e0164186.
37. Lassen J, Hansen M, Sorensen MK, Aamand GP, Christensen LG, Madsen P. Genetic relationship between body condition score, dairy character, mastitis, and diseases other than mastitis in first-parity Danish Holstein cows. J Dairy Sci. 2003;86(11):3730–3735.

38. Richardson CM, Beauchemin KA, Hristov AN, et al. Plasma metabolomics of beef cattle with divergent residual feed intake is associated with differences in the ruminal microbiome. Front Anim Sci. 2021; 2:724051.
39. Subramanian I, Verma S, Kumar S, Jere A, Anamika K. Multi-omics data integration, interpretation, and its application. Bioinform Biol Insights. 2020; 14:1177932219899051.
40. Ritchie MD, Holzinger ER, Li R, Pendergrass SA, Kim D. Methods of integrating data to uncover genotype-phenotype interactions. Nat Rev Genet. 2015;16(2):85–97.
41. Wild CP. Complementing the genome with an 'exposome': the outstanding challenge of environmental exposure measurement in molecular epidemiology. Cancer Epidemiol Biomarkers Prev. 2005;14(8):1847–1850.
42. Civelek M, Lusis AJ. Systems genetics approaches to understand complex traits. Nat Rev Genet. 2014;15(1):34–48.
43. Rohart F, Gautier B, Singh A, Lê Cao K-A. mixOmics: an R package for 'omics feature selection and multiple data integration. PLoS Comput Biol. 2017;13(11): e1005752.
44. Jiang Q, Hu Z, Xu R, et al. Integration of transcriptomic and metabolomic data reveals inflammatory response to mastitis in dairy cows. J Anim Sci Biotechnol. 2022;13(1):82.
45. Moser G, Lee SH, Hayes BJ, Goddard ME, Wray NR, Visscher PM. Simultaneous discovery, estimation and prediction analysis of complex traits using a Bayesian mixture model. PLoS Genet. 2015;11(4): e1004969.
46. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. BMC Bioinformatics. 2008; 9:559.
47. Ma S, Zhang B, LaFave LM, et al. Chromatin potential identified by shared single-cell profiling of RNA and chromatin. Cell. 2020;183(4):1103–1116.
48. Wang T, Shao W, Huang Z, et al. MOGONET integrates multi-omics data using graph convolutional networks allowing patient classification and biomarker identification. Nat Commun. 2021;12(1):3445.
49. Ollier S, Robert-Granié C, Bernard L, Chilliard Y, Leroux C. Mammary transcriptome analysis of food-deprived lactating goats highlights genes involved in milk secretion and lipid metabolism. J Nutr. 2007;137(3):560–567.
50. Loor JJ, Everts RE, Bionaz M, et al. Nutrition-induced ketosis alters metabolic and signaling gene networks in liver of periparturient dairy cows. Physiol Genomics. 2007;32(1):105–116.
51. Groenen MAM, Archibald AL, Uenishi H, et al. Analyses of pig genomes provide insight into porcine demography and evolution. Nature. 2012;491(7424):393–398.
52. te Pas MFW, Keuning E, Hulsegge B, Hoving-Bolink AH, Evans G, Mulder HA. Longissimus muscle transcriptome profiles related to carcass and meat

quality traits in fresh meat Pietrain carcasses. J Anim Sci. 2010;88(12):4044–4055.
53. Moe KK, Purevdorj E, Mori T, et al. multi-omics study reveals that divergent selection for growth rate altered amino acid, fatty acid, and purine metabolism in broiler lines. Front Genet. 2021; 12:678677.
54. Beaumont M, Roméo Q, Yalcin I, Blachier F, Lan A. Tryptophan metabolites along the intestinal epithelium: substrate availability, signaling, and epithelial barrier function. Amino Acids. 2022;54(12):1499–1512.
55. Gharbi K, Semple CA, Ferguson MM, et al. A genetic linkage map for Atlantic salmon (Salmo salar). Genetics. 2006;173(1):237–248.
56. Sveen LR, Krasnov A, Timmerhaus G, Sundaram AYM, Handeland S, Stien LH, Ytrestøyl T, Calabrese S, Salte R, Kiessling A, Pedersen H. Transcriptomic responses to acute sea lice infestation in Atlantic salmon. Parasite. 2017; 24:34.
57. Kolodziejczyk AA, Kim JK, Svensson V, Marioni JC, Teichmann SA. The technology and biology of single-cell RNA sequencing. Mol Cell. 2015;58(4):610–620.
58. Nguyen QH, Pervolarakis N, Blake K, et al. Profiling human breast epithelial cells using single cell RNA sequencing identifies cell diversity. Nat Commun. 2018;9(1):2028.
59. Stoeckius M, Hafemeister C, Stephenson W, et al. Simultaneous epitope and transcriptome measurement in single cells. Nat Methods. 2017;14(9):865–868.
60. Rao A, Barkley D, Franca GS, Bhatt DL, Bhatt NP, Bhatt PP, Bhatt P. Exploring tissue architecture using spatial transcriptomics. Nature. 2021;596(7871):211–220.
61. Albrecht E, Schering L, Buck F, et al. Ectopic fat deposition in skeletal muscle: specific promotion of intramuscular lipid storage by small-scale cell clusters of adipogenic cells. Front Physiol. 2020; 11:555.
62. Huang Z, Zhan X, Bhatt DL, et al. A variationally autoencoded regression for predicting the risk of therapy failure and survival in cancer patients using multi-omics data. PLoS Comput Biol. 2020;16(9): e1008271.
63. Simidjievski N, Bodnar C, Walker I, et al. Variational autoencoders for cancer data integration: design principles and computational practice. Front Genet. 2019; 10:1205.
64. Nguyen E, Poli F, Waytowich NR, et al. Sequence modeling and design from molecular to genome scale with Evo. Science. 2024;386(6723): eado9336.
65. Flint HJ, Scott KP, Louis P, Duncan SH. The role of the gut microbiota in nutrition and health. Nat Rev Gastroenterol Hepatol. 2012;9(10):577–589.
66. Roehe R, Dewhurst RJ, Duthie CA, et al. Bovine host genetic variation influences rumen microbial methane production with best selection criterion for low methane emitting and efficiently feeding animals. PLoS Genet. 2016;12(2): e1005846.

67. Beauchemin KA, Ungerfeld EM, Eckard RJ, Wang M. Review: Fifty years of research on rumen methanogenesis: lessons learned and future challenges for mitigation. Animal. 2020;14(S1): s2–s16.
68. Gomez-Cabrero D, Abugessaisa I, Maier D, et al. Data integration in the era of omics: current and future challenges. BMC Syst Biol. 2014;8(Suppl 2):I1.
69. Brazma A, Hingamp P, Quackenbush J, et al. Minimum information about a microarray experiment (MIAME)—toward standards for microarray data. Nat Genet. 2001;29(4):365–371.
70. Andersson L, Archibald AL, Bottema CD, et al. Coordinated international action to accelerate genome-to-phenome with FAANG, the Functional Annotation of Animal Genomes project. Genome Biol. 2015;16(1):78.
71. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. J R Stat Soc B. 1995;57(1):289–300.
72. Daetwyler HD, Villanueva B, Woolliams JA. Accuracy of predicting the genetic risk of disease using a genome-wide approach. PLoS One. 2008;3(10): e3395.
73. Rege JEO, Gibson J. Animal genetic resources and economic development: issues in relation to economic valuation. Ecol Econ. 2003;45(3):319–330.
74. Mwai OA, Hanotte O, Kwon Y-J, Cho S. African indigenous cattle: unique genetic resources in a rapidly changing world. Asian-Australas J Anim Sci. 2015;28(7):911–921.
75. Mrode R, Ojango JMK, Okeyo AM, Mwacharo JM. Genomic selection and use of molecular tools in breeding programs for indigenous and crossbred cattle in developing countries: current status and future prospects. Front Genet. 2019; 9:694.
76. Leroy G, Mary-Huard T, Verrier E, et al. Methods to estimate effective population size using pedigree data: examples in dog, sheep, cattle and horse. Genet Sel Evol. 2013;45(1):1.
77. European Food Safety Authority (EFSA). Statement on the update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA. EFSA J. 2018;16(7):5375.

Frontiers in Animal Biology and Biodiversity

ISBN: 978-93-49938-98-4 | Year: 2026 | pp: 71 – 82 |

A Study on Therapeutic Potential of Curcumin in Cardio Protection

[1]Anand Konkala

[2]A. Jayasankar

[3]R. Kamli Naik

[4]M. Venkata Chandrudu

[1]Department of Zoology, Government City College (Autonomous), Hyderabad, Telangana

[2]Department of Zoology, S.V.C.R. Government Degree College, Palamaner, Andhra Pradesh

[3]Department of Zoology Govt Degree College for Men, (Cluster university), Kurnool, Andhra Pradesh

[4]Department of Zoology, School of Life Sciences, Rayalaseema University, Kurnool, Andhra Pradesh

Email: chandramidde4@gmail.com

Article DOI Link: https://zenodo.org/uploads/21281968

DOI: 10.5281/zenodo.21281968

Abstract

Cardiovascular diseases (CVDs) remain the leading cause of death worldwide, accounting for nearly one-third of all global mortalities. Despite major breakthroughs in conventional pharmacology, the global burden of CVD continues to escalate. Because of these limitations, clinical interest is rapidly shifting toward nutraceuticals and phytochemicals. these plant-based compounds offer a promising, accessible, and complementary strategy for both preventing and managing heart disease. Chief among them is curcumin—the primary bioactive polyphenol derived from turmeric (*Curcuma longa*). Curcumin has drawn intense scientific interest due to its broad spectrum of therapeutic actions, which include powerful antioxidant, anti-inflammatory, anti-apoptotic (preventing cell death), and lipid-lowering properties. Growing experimental and clinical data indicate that curcumin achieves these deep cardioprotective effects by targeting multiple molecular pathways simultaneously. These include modulation of oxidative stress, suppression of

inflammatory signaling pathways such as nuclear factor kappa B (NF-κB), inhibition of oxidized low-density lipoprotein (ox-LDL)–mediated vascular injury, and attenuation of NLRP3 inflammasome activation. This review summarizes the cardioprotective mechanisms of curcumin and presents key evidence from experimental and clinical studies on its role in cardiovascular disease prevention and treatment. It also examines recent formulation strategies aimed at improving curcumin's bioavailability and therapeutic efficacy. Insights into these mechanisms and delivery approaches could support the development of curcumin-based adjunct therapies for cardiovascular disease management.

Keywords: Curcumin, Cardiovascular diseases, Atherosclerosis, Oxidative stress, Inflammation

Introduction

Cardiovascular diseases (CVDs), disorders of the heart and blood vessels, are the leading cause of global mortality. Their burden continues to escalate, presenting a critical public health challenge. In 2016, CVDs were responsible for nearly 20 million deaths — roughly one-third of all deaths worldwide — with projections indicating further increases by 2030 [1]. Despite this rising prevalence, public awareness of cardiovascular risk remains inadequate. At-risk individuals frequently exhibit hypertension, obesity, and dysregulated glucose or lipid metabolism [2]. Although significant advances have been made in understanding CVD pathophysiology, associated morbidity and mortality remain unacceptably high [3]. Notably, the growing incidence of heart failure due to maladaptive cardiac remodelling after myocardial infarction (MI) has driven intensive research into pharmacological strategies to reduce tissue injury and promote cardiac repair [4]. Yet, despite encouraging preclinical results, few agents that limit adverse scar formation, enhance myocardial repair, and improve post-reperfusion cardiac function have translated into routine clinical use for MI patients [5]. This highlights the critical need for safer, more effective therapeutics for CVD prevention and management.

Therapeutic strategies for cardiac diseases have traditionally focused on lipid-lowering therapies, particularly statins. The growing concern over adverse side effects associated with conventional pharmacological therapies has further stimulated interest in nutraceutical-based approaches. Over the years, extensive experimental research has investigated the antibacterial, anti-inflammatory, antioxidant, and anticancer properties of these compounds in both in vitro and in vivo models [6]. Moreover, numerous clinical trials have evaluated their therapeutic potential across a range of human diseases. They are increasingly being utilized in the prevention and management of various diseases [8]. Among these, phytochemicals—naturally occurring compounds found in fruits and vegetables have gained considerable attention for their potential physiological and therapeutic

effects. Notable examples include herbal polyphenols such as curcumin, beta-carotene, and resveratrol Curcumin exhibits anti-inflammatory, antioxidant, anti-infective, lipid-lowering, and endothelial-protective properties [10].

Emerging evidence further suggests that curcumin suppresses human cytomegalovirus (HCMV) infection in experimental models, thereby improving the vascular cellular microenvironment and limiting lesion development [12]. Preclinical studies report up to a 50% reduction in aortic lesion area, while early clinical trials demonstrate improvements in endothelial function and lipid profiles. Previous studies have been demonstrated that mechanistically, it may attenuate atherosclerosis progression through suppression of nuclear factor kappa B (NF-κB) signaling, reduction of oxidized low-density lipoprotein (ox-LDL) sensitivity, and inhibition of NLRP3 inflammasome activation [13]. Additionally, curcumin inhibits vascular smooth muscle cell (VSMC) proliferation and migration, key processes involved in plaque formation and progression [14]. Despite these promising findings, curcumin's clinical translation remains limited by poor bioavailability, underscoring the need for advanced delivery systems. These epidemiological trends underscore the need for safer and more effective preventive and therapeutic strategies, including the potential integration of nutraceuticals into cardiovascular care.

This review critically evaluates the multifaceted cardioprotective effects of curcumin by integrating mechanistic insights, preclinical data, and emerging clinical evidence, and discusses innovative formulation strategies aimed at translating experimental promise into tangible cardiovascular benefit.

Curcumin Regulates the Molecular Mechanism of Myocardial Ischemia

Myocardial ischemia/reperfusion injury (IRI) following cardiac surgery and myocardial infarction (MI) is a primary cause of heart failure and cardiac mortality [14]. The fundamental pathological processes of myocardial I/R injury include inflammation, oxidative stress, and cardiomyocyte apoptosis. Curcumin has been widely reported to exert anti-inflammatory, antioxidant, anti-apoptotic, and anti-fibrotic properties [15], thereby demonstrating significant cardioprotective effects against IRI. Curcumin reduces myocardial I/R injury through multiple molecular mechanisms. It inhibits oxidative stress, suppresses apoptosis, and attenuates inflammation, resulting in decreased release of cardiac injury biomarkers such as lactate dehydrogenase (LDH) and creatine kinase (CK), as well as improved post-ischemic cardiac function. Moreover, curcumin preserves cardiomyocyte structural integrity, attenuates extracellular matrix (ECM) remodeling, and enhances cardiac contractility [16].

Oxidative stress is a central contributor to ischemic myocardial injury [21]. Curcumin alleviates isoproterenol-induced myocardial ischemia by increasing antioxidant enzyme levels, including superoxide dismutase (SOD), catalase, and

glutathione, while suppressing thiobarbituric acid reactive substances and LDH leakage. Previous studies have shown that curcumin enhances mitochondrial antioxidant activity, as evidenced by elevated mitochondrial SOD levels and decreased mitochondrial hydrogen peroxide and malondialdehyde concentrations. Furthermore, curcumin protects cardiomyocytes from hypoxia/reoxygenation injury by inhibiting the MAPK and Notch signaling pathways to reduce reactive oxygen species (ROS) levels [22] and alleviates hyperglycemia-induced H9c2 cell damage via activation of the Nrf2/HO-1 signaling pathway [23]. Curcumin also attenuates myocardial IRI by suppressing apoptosis. It upregulates the anti-apoptotic protein Bcl-2 and downregulates the pro-apoptotic protein caspase-3 through activation of the JAK2/STAT3 signaling pathway, leading to improved post-ischemic cardiac function, reduced infarct size, and decreased LDH release [26]. In rabbit models of cardiopulmonary bypass surgery, curcumin inhibited the activation of caspase-3 and poly (ADP-ribose) polymerase (PARP), thereby reducing cardiomyocyte apoptosis and myocardial injury [24]. Additionally, activation of the PI3K/Akt/mTOR and MAPK pro-survival pathways by curcumin further contributes to reduced apoptosis and attenuation of inflammation and oxidative stress [25].

Inflammation is another critical mechanism involved in myocardial I/R injury. Curcumin inhibits the upregulation of pro-inflammatory cytokines, including IL-1, IL-6, IL-8, TNF-α, and MCP-1, and suppresses NF-κB activation following IRI [26]. In rabbit I/R models, curcumin treatment significantly reduced NF-κB activity, resulting in downregulation of pro-inflammatory gene expression and decreased matrix metalloproteinase (MMP) activation. Toll-like receptor 2 (TLR2), a key mediator of innate immunity, is also involved in myocardial IRI [27]. Pretreatment with curcumin selectively inhibited TLR2, reduced infarct size, and alleviated ischemic injury, indicating both preventive and therapeutic potential [28]. Sirtuin 1 (SIRT1), a NAD+-dependent histone deacetylase, regulates cellular stress responses, energy metabolism, and apoptosis through deacetylation of substrates such as NF-κB, FOXO1, TP53, and AMPK [29]. Curcumin pretreatment upregulates and activates SIRT1, thereby attenuating mitochondrial oxidative damage, reducing myocardial enzyme release, and inhibiting apoptosis in myocardial I/R injury [30]. By activating SIRT1, curcumin stabilizes the mitochondrial membrane and preserves intracellular integrity, reducing infarct size and improving cardiac function.

Inflammatory cytokines such as IL-1β, TNF-α, and IL-18 contribute to cardiac fibrosis by promoting TGF-β expression and SMAD2/3 activation. By inhibiting these pathways, curcumin mitigates inflammatory activation and subsequent fibrotic remodeling. Although TGF-β1-SMAD2/3 signaling may also contribute to cardiac repair through induction of CTHRC1 and activation of WNT5A signaling [31], excessive activation leads to pathological fibrosis, which curcumin effectively attenuates. Curcumin improves cardiac contractility following I/R injury. In rats

pretreated with curcumin (300 mg/kg/day for one week), increased expression of connexin 43 was observed, facilitating synchronous cardiomyocyte contraction and enhancing mitochondrial energy production to ensure adequate energy supply [32].

Curcumin Effect on Molecular Mechanism of Vascular Remodelling in Cardiac Damage

Curcumin has demonstrated significant cardioprotective effects in pressure-overload–induced cardiac remodelling. In pressure-overloaded rabbits, curcumin improved left ventricular function by inhibiting collagen remodelling and suppressing myocardial expression of TNF-α and matrix metalloproteinases (MMPs) [33]. Similarly, Tharaux et al. reported that curcumin attenuated fibrosis in the aorta and renal arteries, thereby preserving vascular wall elasticity. These vascular protective effects were mediated through inhibition of the renin–angiotensin system and suppression of the transforming growth factor-β1 (TGF-β1)/collagen axis [34].

Further evidence from two independent research groups highlights the modulatory role of curcumin in cardiac remodelling following pressure overload or myocardial ischemia. Both studies emphasized the regulation of nuclear acetylation processes, particularly involving histone acetyltransferases (HATs) and p300, an intrinsic HAT critical for transcriptional activation. Morimoto et al. demonstrated that p300 and the p300/GATA-4 complex are upregulated in cardiac hypertrophy in pressure-overloaded salt-sensitive Dahl rats and in a rat model of myocardial infarction. Curcumin inhibited HAT activity, reduced acetylation of hypertrophy-associated transcription factors such as GATA-4, and suppressed the expression of brain natriuretic peptide (BNP), a marker of cardiac hypertrophy. These effects were associated with significant preservation of cardiac function [35].

Beyond pressure overload and hypertrophy models, the effects of curcumin have also been examined in human and rabbit hearts subjected to chronic hypoxia. Chronic hypoxia induces subcellular redistribution of inducible heat shock protein 70 (Hsp70i) via activation of p38 MAP kinase and JNK pathways, which constitute potent endogenous cardioprotective mechanisms. Similarly, another study showed that ischemic preconditioning activates stress-activated kinases with cardioprotective effects; however, these benefits were abolished when isolated rat hearts were perfused with curcumin [36]. These findings align with previous reports indicating that excessive antioxidant activity may blunt the tissue-protective effects of ischemic preconditioning.

Cardio Protective of Curcumin in Myocardial Infarction

Curcumin, a natural compound found in turmeric, has been extensively studied for its potential protective effects against myocardial infarction (MI) and ischemia/reperfusion (I/R) injury. While the research to date has primarily involved

animal models, the findings offer promising insights into curcumin's cardioprotective capabilities.

Studies have varied widely in their experimental designs, ranging from ex vivo heart experiments to isoproterenol-induced cardiac injury and I/R injury post-coronary artery ligation. Despite these differences, several consistent observations have emerged regarding curcumin's protective effects. In animal models treated with isoproterenol, curcumin was found to reduce morphological changes and infarct size in the heart. It protected cardiomyocytes against apoptosis by upregulating the anti-apoptotic regulator BCL2 and downregulating BAX, a pro-apoptotic factor. This protective effect was evident in both isolated neonatal cardiomyocytes and in situ cardiomyocytes [37].

Curcumin administration normalized serum levels of creatine kinase and lactate dehydrogenase, both indicators of tissue damage elevated after MI. It also prevented excessive collagen deposition and matrix metalloproteinase activation, leading to reduced scar size and improved heart function, as evidenced by enhanced left ventricular ejection fraction and contractility. Curcumin has shown significant anti-inflammatory effects post-I/R injury by attenuating the upregulation of inflammatory cytokines such as TNFα, IL-1β, and IL-6. It also influenced macrophage polarization, promoting the M2 subtype known for its reparative properties, while reducing M1 macrophage markers. Curcumin plays a vital role in re-establishing redox homeostasis following I/R injury. It mitigates oxidative stress by increasing glutathione (GSH) levels and enhancing the activity of antioxidative enzymes like catalase and superoxide dismutase (SOD). Notably, curcumin upregulates NRF2, which in turn increases SOD2 expression, further supporting mitochondrial function [38].

Curcumin improves mitochondrial membrane potential and respiration, crucial for energy production and overall heart function. It has been shown to reactivate the histone deacetylase SIRT1, essential for mitochondrial function and biogenesis. Curcumin mimics the effects of caloric restriction, known for its cardiovascular benefits, by inhibiting the mammalian target of rapamycin (mTOR) pathway and inducing autophagy. This process is critical for maintaining cellular health, particularly in post-mitotic tissues like the heart. The JAK2/STAT3 signaling pathway is a crucial mediator of curcumin's cardioprotective effects. Activation of this pathway reduces infarct size and reactive oxygen species (ROS) generation, enhancing post-ischemic recovery [39].

The promising results from animal studies suggest that curcumin could be an effective therapeutic agent for reducing MI damage and improving heart function. Its diverse mechanisms of action, including anti-inflammatory, antioxidative, and autophagy-inducing properties, highlight its potential as a supportive treatment post-ischemia. Future research, particularly clinical trials, is needed to further explore curcumin's applicability in human cardiovascular health.

Curcumin Effect on Genetically Influenced Risk Factors for Cardiovascular Diseases

While the vascular wall and the heart are traditionally viewed as critical players in cardiovascular diseases (CVDs), recent research highlights the significant role of brown adipose tissue (BAT). Historically considered responsible primarily for thermogenesis, BAT is now recognized for its broader metabolic functions and potential protective effects against CVDs.

BAT is characterized by its rich content of mitochondria, essential for heat production through non-shivering thermogenesis. This process is facilitated by the expression of Uncoupling Protein 1 (UCP1), a transmembrane protein that translocates protons across the inner mitochondrial membrane. By bypassing the mitochondrial ATP synthase, this activity increases energy expenditure and heat production. In humans, BAT is formed during early fetal development and is primarily located in the axillary, cervical, perirenal, and periadrenal regions. It is most abundant at birth, but despite a decrease in size with age, BAT remains present and metabolically active in adults. Beyond its thermogenic capability, BAT plays a crucial role in regulating energy balance and body weight. It also contributes to the control of glucose and lipid metabolism. Interestingly, white adipocytes, which constitute the major energy storage organ known as white adipose tissue (WAT), can be converted into brown adipocytes through a process called "beiging." [40]

Research using animal models susceptible to atherosclerosis has demonstrated a protective role for BAT in CVDs. Activation of BAT, whether through short-term cold exposure or pharmacological interventions, has been shown to accelerate the clearance of triglycerides from the plasma. The fatty acids derived from triglyceride-rich lipoproteins are absorbed by BAT, where they undergo mitochondrial fatty acid oxidation [41].

This BAT-mediated reduction in hyperlipidemia and hypercholesterolemia offers protection against the development of atherosclerosis in mice. Therefore, BAT's ability to influence lipid metabolism and energy expenditure presents a promising avenue for therapeutic strategies aimed at combating CVDs. The emerging understanding of BAT's metabolic roles underscores its importance beyond thermogenesis. Its involvement in energy regulation and its potential protective effects against cardiovascular diseases highlight BAT as a promising target for future research and therapeutic interventions. The conversion of WAT to BAT, or "beiging," further expands the potential for manipulating adipose tissue in the context of metabolic and cardiovascular health [42].

Curcumin Effect on Drug Induced Cardiac Diseases

The therapeutic effects and underlying mechanisms of curcumin have increasingly become a focus of pharmacological research. Curcumin has demonstrated broad cardioprotective effects, particularly in the cardiovascular field. It significantly

attenuates cardiomyocyte injury following ischemia and hypoxia, inhibits myocardial hypertrophy and fibrosis, improves ventricular remodeling, reduces drug-induced myocardial injury, ameliorates diabetic cardiomyopathy (DCM), alleviates vascular endothelial dysfunction, suppresses foam cell formation, and inhibits vascular smooth muscle cell (VSMC) proliferation [43].

Previous reviews have summarized the cardiovascular protective effects of curcumin. In 2020, Li et al. reviewed preclinical studies addressing curcumin in cardiac hypertrophy, heart failure, abdominal aortic aneurysm, stroke, drug-induced cardiotoxicity, and diabetic cardiovascular complications, and discussed potential molecular targets. However, neither review provided a comprehensive synthesis of the complete cardiovascular protective signaling network of curcumin, nor did they consistently specify curcumin dosages. In contrast, the present review classifies the effects of curcumin according to specific cell types within the cardiovascular system and provides a more integrated analysis of signaling pathways while explicitly reporting dosage information, given that pharmacological effects are dose-dependent.

The protective mechanisms of curcumin against CVDs are multifactorial and network-based. The signaling pathways described in current literature likely represent only part of the downstream cascades triggered by curcumin's interaction with upstream molecular targets. Therefore, identifying the direct molecular targets of curcumin using systems biology and chemical biology approaches is warranted.

The authors also reported decreased activity of lysosomal hydrolases—including β-glucuronidase, β-N-acetylglucosaminidase, cathepsin B, cathepsin D, and acid phosphatase—in curcumin-treated animals. Additionally, curcumin enhanced the myocardial antioxidant defense system by increasing superoxide dismutase, catalase, and glutathione levels. These beneficial effects were attributed to membrane stabilization at the level of nuclei, mitochondria, lysosomes, and microsomes. Subsequent studies further showed that curcumin reduced both degradation and synthesis of new collagen in the hearts of treated mice, supporting its anti-fibrotic properties.

Overall, accumulating evidence suggests that curcumin exerts antioxidant, anti-inflammatory, anti-apoptotic, and anti-fibrotic effects through complex and interconnected molecular networks, underscoring its therapeutic potential in cardiovascular disease.

References

1. Pagano E, Souto EB, Durazzo A, Sharifi-Rad J, Lucarini M, Souto SB, Salehi B, Zam W, Montanaro V, Lucariello G, Izzo AA, Santini A, Romano B. Ginger (Zingiber officinale Roscoe) as a nutraceutical: Focus on the metabolic, analgesic, and anti-inflammatory effects. Phytotherapy Research. 2021;35(5):2403–2417.https://doi.org/ 10. 1002/ ptr.696.

2. Heidenreich PA, Trogdon JG, Khavjou OA, et al. Forecasting the future of cardiovascular disease in the United States: a policy statement from the American Heart Association. Circulation. 2011;123(8):933–944. doi:10.1161/CIR.0b013e31820a55f52
3. Del Pinto R, Vitale M, Morreale ME, Mottinelli C, Muiesan ML, Volpe M, Ferri C.
4. Cardiovascular risk factors and diseases and awareness of related burden in women: Results of a survey in Italian pharmacies. High Blood Pressure & Cardiovascular Prevention. 2025;32(2):191–198. https://doi.org/10.1007/s40292-025-00701-4.
5. Gallagher G, Menzie S, Huang Y, Jackson C, Hunyor SN (2007). Regional cardiac dysfunction is associated with specific alterations in inflammatory cytokines and matrix metalloproteinases after acute myocardial infarction in sheep. Basic Res Cardiol 102: 63–72.
6. Downey JM, Cohen MV (2009). Why do we still not have cardioprotective drugs? Circ J 73: 1171–1177.
7. Mao QQ, Xu XY, Cao SY, Gan RY, Corke H, Beta T, Li HB. Bioactive compounds and bioactivities of ginger (Zingiber officinale Roscoe). Molecules.2019;24(5):1-21. https://doi.org /10.3390/ molecules 24050703.
8. Yanjun Zhang, Li S, Wang X, Zhang L, Yang L Nutraceuticals and functional foods in human health and disease prevention. Frontiersin Pharmacology. 2021; 12:728147. https:// doi.org/ 10.3389/fp har.2021.728147.
9. Kavyani Z, Najafi K, Naghsh N, Bavandpour Karvane H,Musazadeh V. The effects of curcumin supplementation on biomarkers of inflammation, oxidative stress, and endothelial function: A meta-analysis of meta-analyses. Prostaglandins & Other Lipid Mediators. 2024; 174:106867. https://doi.org/10.1016/j.prostaglandins.2024.10686.
10. Sowndhar Rajan B, Krishnan K, Vellaichamy E. Diet-derived advanced glycation end products (dAGEs) induce proinflammatory cytokine expression in cardiac and renal tissues of experimental mice: protective effect of curcumin. Cardiovasc Toxicol. 2022;22(1):35–51. doi:10.1007/ s12012-021-09697-4.
11. Wongeakin N, Bhattarakosol P, Patumraj S. Molecular mechanisms of curcumin on diabetes-induced endothelial dysfunctions: txnip, ICAM-1, and NOX2 expressions. Biomed Res Int. 2014; 2014:161346. doi:10.1155/2014/161346.
12. Schirone, L., Forte, M., D'Ambrosio, L., Valenti, V., Vecchio, D., Schiavon, S., et al. (2022). An overview of the molecular mechanisms associated with myocardial ischemic injury: State of the art and translational perspectives. Cells 11 (7), 1165. doi:10.3390/cells11071165.

13. Li, D., Li, Y., and Wu, D. (2020). Preparation of curcumin nanoemulsion and its protective effect on myocardial ischemia-reperfusion in rats. Chin. J. Of Comp. Med. 30 (5), 97–103. doi: 10.3969/j.issn.1671-7856.2020.05.015.
14. Yong Yuan, Huang H, Hu T, Zou C, Qiao Y, Fang M, Liu J, Lai S.Curcumin pretreatment attenuates myocardial ischemia/reperfusion injury by inhibiting ferroptosis, autophagy and apoptosis via HES1. International Journal of Molecular Medicine. 2024; 54:110. https://doi.org/10.3892/ ijmm. 2024.5434.
15. Jiankun Cui, Wang X, Dong L,Wang Q. Curcumin reduces myocardial ischemia–reperfusion injury by increasing endogenous H_2S levels and modulating m6A RNA methylation. MolecularBiologyReports.2024;51: 558.https://doi.org/ 10.1007/s 11033- 024-09478-6.
16. Y. Yang, W. Duan, Y. Lin, W. Yi, Z. Liang, J. Yan, et al., Sirt1 activation by curcumin pretreatment attenuates mitochondrial oxidative damage induced by myocardial ischemia reperfusion injury, Free Radic Biol Med 65 (2013) 667-679.
17. S.G. Ong, W.H. Lee, L. Theodorou, K. Kodo, S.Y. Lim, D.H. Shukla, et al., Hif-1 reduces ischaemia-reperfusion injury in the heart by targeting the mitochondrial permeability transition pore, Cardiovasc Res 104 (2014) 24- 36
18. V. Tanwar, J. Sachdeva, M. Golechha, S. Kumari, D.S. Arya, Curcumin protects rat myocardium against isoproterenol-induced ischemic injury: Attenuation of ventricular dysfunction through increased expression of hsp27 along with strengthening antioxidant defense system, J Cardiovasc Pharmacol 55 (2010) 377-384.
19. W. Duan, Y. Yang, J. Yan, S. Yu, J. Liu, J. Zhou, et al., the effects of curcumin post-treatment against myocardial ischemia and reperfusion by activation of the jak2/stat3 signaling pathway, Basic Res Cardiol 107 (2012) 263.
20. Yuan Y, Huang H, Hu T, Zou C, Fang M, Liu J, Lai S. Curcumin pretreatment attenuates myocardial ischemia/reperfusion injury by inhibiting ferroptosis, autophagy and apoptosis via HES1 signaling. International Journal of Molecular Medicine. 2024; 54:110.
21. Wu Y, Guo F, Liu Y, Li J, Shi W, Song L, Wang G, Liu J. Curcumin mitigates heatstroke-induced myocardialinjurybymodulatingtheAkt/Bad/Caspase-3pathway.Biochemical and Biophysical Research Communications.2025;758: 151653.https://doi.org/10.1016/ j.bbrc .2025.151653.
22. Wu, H. J., Zhang, K., Ma, J. J., Wang, L., and Zhuang, Y. (2021). Mechanism of Curcumin against myocardial ischaemia-reperfusion injury based on the P13k/akt/mtor Signallingpathway.Eur.Rev.Med.Pharmacol.Sci.25(17),5490–5499.Doi:10.26355/ Eurrev _ 202109_266 58.
23. Cui J, Wang X, Dong L, Wang Q. Curcumin reduces myocardial ischemia-reperfusion injury by increasing endogenous H_2S levels and modulating m6A RNA methylation.

24. Molecular Biology Reports. 2024; 51:558.
25. Pawar HD, Mahajan UB, Nakhate KT, Agrawal YO, Patil CR, Meeran MFN, Sharma C, Ojha S, Goyal SN. Curcumin protects diabetic mice against isoproterenol-induced myocardial infarction by modulating CB2 cannabinoid receptors. Life (Basel). 2022;12(5):624. https://doi.org/10.3390/life12050624.
26. Huang S, Zhang Z, Zhong J, Chen X, Yang P. (2020). Curcumin protects against myocardial ischemia/reperfusion injury by regulating oxidative stress and inflammatory responses. Journal of Cellular Physiology.235(10):7687-7698. https://doi.org/10.1002/jcp.2966.
27. J. Xiao, X. Sheng, X. Zhang, M. Guo, X. Ji, Curcumin protects against myocardial infarctioninduced cardiac fibrosis via sirt1 activation in vivo and in vitro, Drug Des Devel Ther 10 (2016) 1267-1277.
28. N.P. Wang, Z.F. Wang, S. Tootle, T. Philip, Z.Q. Zhao, Curcumin promotes cardiac repair and ameliorates cardiac dysfunction following myocardial infarction, Br J Pharmacol 167 (2012) 1550-1562.
29. Y.S. Kim, J.S. Kwon, Y.K. Cho, M.H. Jeong, J.G. Cho, J.C. Park, et al., Curcumin reduces the cardiac ischemia-reperfusion injury: Involvement of the toll-like receptor 2 in cardiomyocytes, J Nutr Biochem 23 (2012) 1514- 1523.
30. Azli AMA, Salamt N, Aminuddin A, etal. (2024). The role of curcumin in modulating vascular function and structure: A systematic review. Biomedicines. 12:2281.
31. Tharaux PL, Chatziantoniou C, Fakhouri F, Dussaule JC. Angiotensin II activates collagen I gene through a mechanism involving the MAP/ER kinase pathway. Hypertension. 2000; 36:330-336.
32. Li HL, Liu C, de Couto G, et al. Curcumin prevents and reverses murine cardiac hypertrophy. J Clin Invest. 2008; 118:879-893.
33. Sunagawa Y, Morimoto T, Wada H, Takaya T, Kawamura T, et al. (2011). A natural p300 histone acetyl transferase inhibitor, curcumin, in the prevention of heart failure. Journal of Molecular and Cellular Cardiology. 52(5):1123-1131. https://doi.org/10.1016/j.yjmcc. 2011.01.0 06.
34. Rafiee P, Shi Y, Pritchard KA Jr, et al. Cellular redistribution of inducible Hsp70 protein in the human and rabbit heart in response to the stress of chronic hypoxia: role of protein kinases. J Biol Chem. 2003;278: 43636-43644.
35. Wang X, Wang Y, Li Y, et al. (2021). Curcumin protects cardiomyocytes from hypoxia/ reoxygenation injury by regulating mitochondrial apoptosis pathways.
36. Experimental and Therapeutic Medicine. 21:512.
37. Bartelt A, Heeren J. (2022). Adipose tissue browning and lipid metabolism in cardiovascular disease. Frontiers in Endocrinology. 13:875378. https://doi.org/ 10.3389/ fendo. 2022.875378.

38. Scheja L, Heeren J. (2022). Metabolic interplay between brown adipose tissue and cardiovascular disease. Nature Reviews Cardiology.19:503–518.https:// doi.org/10.1038/s 41569-022-00671-3.
39. Kajimura S, Spiegelman BM, Seale P. (2022). Brown and beige fat: Physiological roles beyond heat generation. Cell Metabolism.34(5):681–702.https://doi.org /10.1016/j. cmet. 2022.03.013.
40. Cannon B, Nedergaard J. (2023). Brown adipose tissue: Function and physiological significance. International Journal of Molecular Sciences. 24:12405. https://doi.org/ 10.3390/ ijms241512405.
41. Ungvari Z, Tarantini S, Donato AJ, et al. (2023). Mechanisms of vascular aging: new perspectives on oxidative stress and mitochondrial dysfunction. Frontiers in Cardiovascular Medicine. 10:1158722. https://doi.org/10.3389/fcvm.2023.1158722.
42. Zhang Y, Wang Y, Liu J, et al. (2024). Oxidative stress and mitochondrial dysfunction in cardiovascular aging. International Journal of Molecular Sciences.25:2384.
43. https: //doi.org/10.3390/ijms25042384.
44. Liguori I, Russo G, Curcio F, et al. (2022). Oxidative stress, aging, and diseases. Antioxidants.11(6):1132. https://doi.org/10.3390/antiox11061132.
45. Qiu Y, Chang S, Zeng Y, Wang X. (2025). Advances in mitochondrial dysfunction and its role in cardiovascular diseases. Cells.14(20):1621 https ://doi.org/ 10.3390/ cells14201621.
46. Luchkova A, Mata A, Cadenas S. (2024). Nrf2 as a regulator of energy metabolism and mitochondrial function. FEBS Letters. 598(17):2092–2105.
47. Li H, Sureda A, Devkota HP, Pittalà V, Barreca D, Silva AS, Tewari D, Xu S, Nabavi SM. (2023). Curcumin, the golden spice intreating cardiovascular diseases. Biomedicine& Pharmacotherapy.156: 113877.https://doi.org/10. 1016/j.biopha.2022.113877.
48. Ghosh S, Banerjee S, Sil PC. (2022). The beneficial role of curcumin on cardiovascular diseases: molecular targets and therapeutic potential. Pharmacological Research. 178 :106 138.https://doi.org/10.1016/j.phrs.2022.106138.
49. Pourbagher-Shahri AM, Farkhondeh T, Ashrafizadeh M, et al. Curcumin and cardiovascular diseases: focus on cellular targets and cascades. Biomed Pharmacother. 2021; 136:111214
50. Ma T, Tian X, Zhang B, et al. Low-dose metformin targets the lysosomal AMPK pathway through PEN2. Nature. 2022;603(7899):159–165. doi:10.1038/s41586-022-04431-8.
51. Nirmala, C.; Puvanakrishnan, R. Protective role of curcumin against isoproterenol induced myocardial infarction in rats. Mol. Cell. Biochem. 1996, 159, 85–93. [CrossRef] [PubMed]

Frontiers in Animal Biology and Biodiversity

ISBN: 978-93-49938-98-4 | Year: 2026 | pp: 83 – 95 |

Environmental and Health Consequences of Toxic Heavy Metal Exposure: A Comprehensive Review

[1]A. Jayasankar

[2]Anand Konkala

[3]M. Venkata Chandrudu

[4]R. Kamli Naik

[1]Department of Zoology, S.V.C.R. Government Degree College, Palamaner, Andhra Pradesh

[2]Department of Zoology, Government City College (Autonomous), Hyderabad, Telangana

[3]Department of Zoology, School of Life Sciences, Rayalaseema University, Kurnool, Andhra Pradesh

[4]Department of Zoology Govt Degree College for Men, (Cluster university), Kurnool, Andhra Pradesh

Email: kamlinaikr@gmail.com

Article DOI Link: https://zenodo.org/uploads/21282090

DOI: 10.5281/zenodo.21282090

Abstract

Heavy metals are harmful pollutants found in various ecosystems due to their toxic nature and long-lasting presence in the environment, contaminating soil, water, and living organisms. They can accumulate in the human body, posing serious health risks. While some heavy metals are a result of human activities, many occur naturally. Living organisms can come into contact with these metals through the food chain and environmental sources like soil, water, and air, which can enhance their harmful effects. Exposure to a combination of several heavy metals is more dangerous to humans than exposure to a single metal, leading to oxidative stress and inflammation that can harm different organs. This study investigates the impact of heavy metals on different organ systems, including their role in influencing biological processes and their potential to cause illnesses. The toxicity of heavy

metals such as lead, cadmium, chromium, and mercury, their effects on the environment, and human health are examined in detail.

Keywords: Heavy metals, health risks, environment, oxidative stress

Introduction

Heavy metals are found in the biosphere, including rocks, soils, and water, and originate from a variety of sources, such as mining, industrial effluents, urban runoff, sewage discharge, soil erosion, natural weathering of the Earth's crust, pesticides, disease control agents used on crops, metal pipes for water, traffic, combustion byproducts from coal-burning plants and many others (Macklin et al. 2023). eavy metals are widely recognized for their harmful effects, recent studies indicate that some metals like nickel, copper, and zinc are vital for human health and are naturally present. In contrast, elements such as Lead (Pb), Mercury (Hg), Arsenic (As), Chromium (Cr), and Cadmium (Cd) lack a known biological function and can be harmful to animals when present in excess amounts. Non-occupational exposure can occur through ingestion of contaminated food and water or through skin contact. Monitoring heavy metal levels in water sources, air, and food is essential for safeguarding the health of both humans and animals (Luo et al., 2020; Azeh Engwa et al., 2019). Food crops grown in environments contaminated with heavy metals causes these elements to bioaccumulate in human food chains via geo-chemical cycles. Chronic poisoning effects on animals have been commonly observed if adequate every day intake amounts are surpassed. Most metallic elements are harmful due to their tendency to accumulate in the soft tissues of animals. Each metal operates through distinct mechanisms and biological pathways within the target system (Qasem et al., 2021).

Long-term exposure to low levels of toxic heavy metals in the environment poses significant health risks, with symptoms often emerging gradually rather than acutely. Although chronic exposure is associated with increased cancer risk, the carcinogenicity of individual metals varies and is not always conclusive. Some metals show limited capacity to induce DNA mutations yet can dysregulate gene expression, thereby influencing cell growth and development [Ekere et al., 2017]. Moreover, these metals can inhibit DNA repair pathways and disrupt normal gene function. Based on evidence of human carcinogenicity, the International Agency for Research on Cancer [IARC] assigns classifications to specific heavy metals.

Method

This study aims to provide an in-depth examination of the current research regarding the toxic mechanisms of heavy metals, with a particular focus on mercury, lead, cadmium, chromium, and arsenic. It delves into their respective adverse effects and underlying toxic pathways. The analysis involves a comparison and discussion of pertinent findings sourced from diverse scientific databases like

PubMed, Web of Science (ISI), Scopus, and Google Scholar, utilizing search terms such as mercury, lead, cadmium, chromium, toxicity, poisoning, and intoxication.

Heavy Metals in the Environment

As trace elements, heavy metals are present in the environment at very low concentrations, from parts per billion to under ten parts per million [Goyer et al., 2001]. Their bioavailability depends on temperature, phase association, adsorption, and sequestration. Lead, a naturally occurring non-biodegradable metal, is increasing in the atmosphere because of human activities like manufacturing, mining, and fossil fuel combustion. High lead exposure is dangerous to human health, especially for children, who face greater risk of poisoning from lead-contaminated dust in their surroundings [Loh et al., 2016].

Cadmium is a heavy metal that poses significant environmental and occupational risks. It is naturally found in black shale and is primarily sourced from volcanic activity, parent materials, marine sedimentary rocks, and phosphates (Bakshi et al., 2018). The earth's crust contains an average cadmium concentration of about 0.1 mg/kg, with marine phosphates containing the highest levels at about 15 mg/kg. Cadmium is released into the atmosphere through natural processes and human activities, leading to exposure for animals and humans through various pathways. Pollution of aquatic ecosystems by cadmium is primarily caused by absorption, industrial discharge, and surface runoff, impacting soil and sediment quality. Human and animal poisoning from cadmium can occur through the ingestion of contaminated food, inhalation of polluted air, or consumption of cadmium-rich water.

Table 1: Toxic Metal in Industrial Sewages

Metal	Manufacturing Industries
Cadmium	Electronics, Phosphate Fertilizer, Pigments and Paints
Lead	Paints, Battery
chromium	Metal Plating, Tanning, Rubber and Photography
copper	Plating, Rayon and Electrical
Mercury	Chlor-Alkali, Scientific Instruments, Chemicals
Nickel	Electroplating, Iron Steel
Arsenic	Metal Hardening, Phosphate and Fertilizer, Paints and Textile

Chromium is a potentially harmful element that can be found in two main forms in the environment: chromium (III) and chromium (VI). Chromium (III) is considered less dangerous than chromium (VI), with the two forms being able to convert into each other during industrial processes (Coetzee et al., 2020). While elemental chromium [Cr (0)] is not naturally occurring, it is introduced into different

environmental components such as air, water, and soil through a combination of natural sources and human activities. Industrial operations, particularly in metal processing, tanneries, chromate production, stainless steel welding, and the manufacturing of chrome pigments, contribute significantly to the release of chromium. This element is commonly used in various commercial applications like industrial welding, chrome plating, dyes and pigments, leather production, wood preservation, and as an anticorrosive agent in cooking systems and boilers (Wang et al., 2006)

Manganese, the most common hazardous heavy metal, can be found in various oxidation states in the environment. When methylcyclopentadienyl manganese tricarbonyl (MMT), an additive in gasoline, is burned, manganese oxides are emitted into the air. Although manganese is essential for many bodily functions, consuming too much can result in serious toxicity according to O'Neal and Zheng (2015).

Arsenic is a common element found in trace amounts in various environmental sources. It can exist in different forms, including trivalent and pentavalent arsenate inorganic forms, as well as organic methylated metabolites like monomethylarsonic acid (MMA), dimethyl arsenic acid (DMA), and trimethyl arsine oxide. Arsenic pollution comes from both natural causes such as volcanic activity and soil erosion, as well as human activities (Tellez-Plaza, et al., 2013).

Heavy Metals Toxicity on Human Health

Cardiovascular Toxicity

Cadmium is a hazardous metal recognized for its toxic nature and carcinogenic properties. Being exposed to cadmium can result in various health concerns, such as kidney disease, bone abnormalities, and cardiovascular issues (Toxicological Profile for Cadmium, 2002). Exposure to cadmium at low to moderate levels has been associated with various health conditions, including hypertension, diabetes, atherosclerosis, peripheral arterial disease, chronic kidney disease, heart attacks, strokes, and heart failure (Hellström et al., 2001; Peters et al., 2010). Lead toxicity is a commonly reported issue related to exposure to heavy metals, particularly affecting children. Lead serves no beneficial purpose in human metabolism. Exposure to lead can occur through various sources, including lead-containing paint, food stored in lead containers, and contaminated water. Prolonged exposure to lead can result in numerous health problems, such as arteriosclerosis, hypertension (Tellez-Plaza et al., 2008) thrombosis, atherosclerosis, and heart disease. These conditions are influenced by factors like oxidative stress (Kianoush et al.,2013), reduced nitric oxide availability, elevated levels of vasoconstrictor prostaglandins, changes in the renin-angiotensin system, decreased vasodilator prostaglandins, disruption of calcium signalling in vascular smooth muscle cells, increased inflammation and endothelial dysfunction, and altered vascular responses

to stimuli.Cobalt exposure can result in temporary weakening of the heart's pumping ability, a form of heart muscle damage called systolic cardiac depression. While cobalt-induced cardiomyopathy can progress slowly and be life-threatening, survivors typically experience recovery of their heart function. Additionally, there may be an increase in T cell proliferation associated with these toxic exposures (Packer, 2016). Some of the necessary metals (Co, Cu, Cr, Ni, and Se) and toxic metals (As, Cd, Pb, and Hg) are metallo-estrogens and may raise the risk of CVD by disrupting hormones (Choe et a.,2003).

Hepatotoxicity

Cadmium exposure primarily affects the kidneys and liver. Acute exposure leads to cadmium accumulation in the liver, causing dysfunction. This disrupts cellular redox balance, inducing oxidative stress and liver cell damage. Both acute and chronic exposure can cause hepatotoxicity, potentially leading to liver failure and increased cancer risk, as noted by Hyder et al. (2013). "Chronic lead exposure damages liver cells, inducing oxidative stress. When paired with organic solvents, lead's harmful effects on the liver can intensify due to shared toxicological properties (Malaguarnera et al., 2012). Prolonged exposure to lead may deplete glycogen and cause cellular infiltration in the liver, potentially leading to chronic cirrhosis. (Hegazy and Fouad, 2014). Arsenic exposure can lead to liver lesions and increased risk of liver cancers, including hepatocellular carcinoma and angiosarcoma (Lu et al., 2001; Liaw et al., 2008). Copper accumulation in the liver, as seen in Wilson's disease, can cause oxidative stress and liver damage. Elevated hepatic copper levels are also associated with cholestatic liver diseases, primarily due to impaired biliary excretion rather than infection (Yu et al., 2019).




Figure 1: Effect of heavy metals on human health (Harischandra et al., 2019)

Nephrotoxicity

The presence of lead can have harmful effects on different organs, with the kidneys being especially vulnerable. Acute exposure to lead can result in lead nephropathy, which may manifest as proximal tubular dysfunction resembling Fanconi syndrome. Conversely, chronic lead nephropathy is defined by tubular atrophy, interstitial fibrosis, renal failure, hyperplasia, and glomerulonephritis (Wang et al., 2013). In rats, the subcutaneous administration of PbA at a dose of 100 mg/kg body weight resulted in increased levels of lipid peroxidation markers such as MDA and 4-HDA in the liver and kidneys when compared to control animals. Furthermore, PbA led to a decrease in levels of SOD and total GSH in the liver and kidneys of rats as reported by El-Sokkary et al. in (2005). Additionally, exposure to PbA at a concentration of 500 mgPb/L in drinking water was found to induce apoptosis by triggering the release of mitochondrial cytochrome C, inhibiting Bcl-2 proteins, and activating caspase-3 in the kidneys of exposed rats, as indicated by Liu et al. in (2012).

Cadmium (Cd) is a naturally-occurring but rare element found in soil and minerals in the form of sulfide, sulfate, carbonate, chloride, and hydroxide salts as well as in water. When present in contaminated water, cadmium can disrupt essential bodily mechanisms, potentially causing short-term or long-term health issues (Cao et al., 2018). Exposure to cadmium has been linked to nephrotoxicity, which can manifest in severe clinical symptoms like glucosuria, Fanconi-like syndrome, phosphaturia, and aminoaciduria (Reyes et al., 2013). When the proximal tubular epithelium in the kidneys is directly exposed to cadmium, it can lead to increased cadmium levels in urine, aminoaciduria, 32-microglobulinuria, and glucosuria. This exposure can also lead to impaired reabsorption of renal tubular phosphate. Prolonged exposure to cadmium can be linked to conditions such as renal tubular acidosis, renal failure, and hypercalciuria, according to Friberg et al. in (2019). The gastrointestinal tract (GIT) and kidneys are the main organs impacted by mercury (Hg) salts. Prolonged exposure can lead to conditions such as acute tubular necrosis, immunological glomerulonephritis, or nephrotic syndrome, primarily due to the preferential accumulation of Hg ions in the renal tubule epithelial cells. As a result, elevated Hg levels can cause significant renal damage. Additionally, chronic exposure to elemental mercury vapors, inorganic mercury, and ingestion of $Hg2+$ salts has been associated with nephrotic syndrome, which is characterized by acute tubular necrosis, proteinuria, and albuminuria (Sanchez et al., 2018)

Neurotoxicity

Neurotoxicity refers to the damaging effects of toxic substances like lead, arsenic, and mercury on the nervous system. Lead exposure impacts both central and peripheral nervous systems, with peripheral effects being more pronounced in adults. Notably, neurotoxicity can occur from both high and low doses of exposure

(Rehman et al., 2018). Lead exposure, especially in early childhood when the brain is still developing, can cause permanent neurological harm (Hosni et al, 2013), with the brain being the most vulnerable organ to its effects. Neurodegenerative defects, including amyotrophic lateral sclerosis, Parkinson's disease, Alzheimer's disease, and multiple sclerosis, result from neurotoxicity induced by cadmium (Branca et al., 2018). Studies involving primary cultures of neurons and glia, isolated mitochondria from mouse brains, and non-neuronal cell lines have demonstrated a strong link between oxidative stress and decreased levels of glutathione (GSH) (Tönnies et al., 2017). Central nervous system (CNS) damage was indicated by increased expression of the key signalling molecule c-fos in the cortex and hippocampus, along with evidence of mercury (Hg) accumulation in the brains of treated rats (Bijoor et al., 2012). Additionally, administering lead acetate (PbA) at a dose of 15 mg/kg to pregnant Wistar female rats led to the induction of pro-inflammatory cytokines, such as TNF-α and IL-1β, in the hippocampus and forebrain of immature rat brains. These findings suggest that prolonged exposure to lead promotes inflammation in the developing CNS of rats, likely due to the activation of glial cells [54]. Furthermore, lead exposure resulted in necrotic changes in the kidneys, liver, and brain [55].

Cadmium's neurotoxicity stems from neural cell death via apoptosis, triggered by factors such as impaired neurogenesis, inhibited neuron gene expression, epigenetic effects, and endocrine disruption (Wang and Du, 2013). Wilson's disease, caused by excess copper retention, leads to neurobehavioral abnormalities resembling schizophrenia. Zinc deficiency hinders neurodevelopment, though effects of excessive zinc are unclear (Cai et al., 2005). Research suggests copper exacerbates zinc-induced neurotoxicity (Tanaka & Kawahara, 2017). Additionally, studies show low-dose chromium exposure can cause brain damage in animal models (Salama et al., 2016).

Carcinogenicity

Lead is a toxic substance known for its carcinogenic properties, which can disrupt the body's DNA repair processes. It also impacts genes that regulate tumour growth and alters the structure and sequence of chromosomes by generating reactive oxygen species (ROS). Furthermore, lead interferes with transcription by displacing zinc from certain regulatory proteins (Silbergeld et al., 2000). Mercury produces reactive oxygen species (ROS) that can enhance tumorigenic signalling and support the proliferation of cancerous cells. These ROS contribute to carcinogenesis by damaging cellular proteins, lipids, and DNA, resulting in significant cellular harm (Reczek and Chandel, 2017). Chromium (Cr) is a naturally occurring heavy metal found in the Earth's crust and seawater, often released during industrial activities. A recent meta-analysis indicated that exposure to chromium (VI) may lead to increased mortality and a higher incidence of various cancers, including lung,

larynx, bladder, kidney, testicular, bone, and thyroid cancers in humans (Deng et al., 2019). Arsenic, on the other hand, induces epigenetic changes, damages DNA, alters the expression of the p53 protein, modifies histones, affects DNA methylation, and decreases p21 expression (Park et al., 2015). Arsenic poisoning heightens cancer risk by binding to DNA-binding proteins and impeding the DNA repair process (Garcia-Esquinas et al., 2013).

Immunological Toxicity

Chromium can adversely affect the human immune system in several ways. High levels of hexavalent chromium can reduce the capacity of alveolar macrophages to engulf particles and weaken the body's humoral immune response. Additionally, chromium can induce two types of hypersensitivity reactions: type I (anaphylactic) and type IV (delayed). Research has established a connection between chromium exposure and the onset of allergic contact dermatitis in multiple studies (Bruynzeel et al., 1988). Both acute and chronic exposure to lead can have detrimental effects on the immune system, resulting in various immune responses, including heightened allergies, increased vulnerability to infectious diseases, autoimmune disorders, and a potential risk of cancer (Hsiao et al., 2011). Certain demographic groups with a history of lead exposure may be at a higher risk for lung, stomach, and bladder cancers. Lead exposure can enhance the production of B and T cells and influence major histocompatibility complex (MHC) activity (Kasten-Jolly et al., 2010). This toxic substance can disrupt both cellular and humoral immune responses, potentially impairing T-cell function and raising the likelihood of developing autoimmunity and hypersensitivity reactions (Mishra, 2009).

Conclusion

Exposure to heavy metals can arise from both environmental and external sources. Ingesting these metals can result in severe toxicity and potentially life-threatening consequences. Our analysis indicates that oral ingestion is a prevalent route for heavy metal exposure. Elevated levels of heavy metals can inflict significant damage on various organs, leading to immunological issues, respiratory problems, increased cancer risk, kidney complications, osteoporosis, and more. It is essential to avoid products containing high levels of toxic heavy metals. Raising awareness about the dangers of heavy metal exposure can empower individuals to minimize their contact with these harmful substances. Additionally, further research is necessary to enhance our understanding of the molecular mechanisms underlying human exposure to combinations of toxic metals and their potential public health implications.

References

1. Akgül, N., Altunkaynak, B. Z., Altunkaynak, M. E., Deniz, Ö. G., Ünal, D., and Akgül, H. M. (2016). Inhalation of mercury vapor can cause the toxic effects on rat kidney. Ren. Fail. 38 (3), 465–473
2. Azeh Engwa, G., Udoka Ferdinand, P., Nweke Nwalo, F., Unachukwu, N.M., (2019). Mechanism and health effects of heavy metal toxicity in humans. Poisoning Mod. World- New Tricks an Old Dog?
3. Bijoor AR, Sudha S and Venkatesh T (2012). Neurochemical and neurobehavioral effects of low lead exposure on the developing brain. Indian J Clin Biochem. 27:147-151.
4. Branca, J.J.V., Morucci, G., Pacini, A., 2018. Cadmium-induced neurotoxicity: Still much ado. Neural Regen. Res. 13, 1879–1882.
5. Bruynzeel, D.P., Hennipman, G., van Ketel, W.G., (1988). Irritant contact dermatitis and chrome-passivated metal. Contact Dermatitis 19, 175–179.
6. Cao, Z. R., Cui, S. M., Lu, X. X., Chen, X. M., Yang, X., Cui, J. P., et al. (2018). [Effects of occupational cadmium exposure on workers' cardiovascular system]. Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi 36(6), 474–477.
7. Cai, L.u., Li, X.-K., Song, Y.e., Cherian, M.G., 2005. Essentiality, toxicology and chelation therapy of zinc and copper. Curr. Med. Chem. 12 (23), 2753–2763.
8. Choe SY, Kim SJ, Kim HG, Lee JH, Choi Y, Lee H and Kim Y (2003). Evaluation of estrogenicity of major heavy metals. Sci Total Environ. 312:15-21.
9. Coetzee, J.J., Bansal, N., Chirwa, E.M.N., (2020). Chromium in environment, its toxic effect from chromite-mining and ferrochrome industries, and its possible bioremediation. Expo. Heal. 12 (1), 51–62.
10. Deng, Y., Wang, M., Tian, T., Lin, S., Xu, P., Zhou, L., et al. (2019). The effect of hexavalent chromium on the incidence and mortality of human cancers: a meta-analysis based on published epidemiological cohort studies. Front. Oncol. 9, 24.
11. Egorova, K. S., and Ananikov, V. P. (2017). Toxicity of metal compounds: Knowledge and myths. Organometallics 36, 4071–4090.
12. Ekere JN, Ukoha IP and Ekere NR (2017) Ecological and human health risk assessment of heavy metal contamination in soil of a municipal solid waste dump in Uyo, Nigeria. Environ. Geochem. Health, 39(3): 497–515.
13. Fay, M. J., Alt, L. A. C, Ryba, D., Salamah, R., Peach, R., Papaeliou, A., et al. (2018). Cadmium nephrotoxicity is associated with altered microRNA expression in the rat renal cortex. Toxics 6 (1), 16.
14. Friberg, L., Kjellström, T., Elinder, C.-G., Nordberg, G.F., (2019). Cadmium and health: a toxicological and epidemiological appraisal. Cadmium Heal. A Toxicol. Epidemiol. Apprais.

15. Garcia-Esquinas, E., Pollán, M., Umans, J.G., Francesconi, K.A., Goessler, W., Guallar, E., Howard, B.V., Yeh, J., Best, L., Navas-Acien, A., 2013. Arsenic Exposure and Cancer Mortality in a US-based Prospective Cohort: the Strong Heart Study. ISEE Conf. Abstr. 2013 (1), 3037.
16. Hegazy, A.M.S., Fouad, U.A., (2014). Evaluation of lead hepatotoxicity; histological, histochemical and ultrastructural study. Forensic Med. Anat. Res. 02 (03), 70 79.
17. Hosni H, Selim O, Abbas M and Fathy A (2013) Semen quality and reproductive endocrinal function related to blood lead levels in infertile painters. Andrologia 45 (2), 120–127.
18. Hou, S., Yuan, L., Jin, P., Ding, B., Qin, N.a., Li, L.i., Liu, X., Wu, Z., Zhao, G., Deng, Y., (2013). A clinical study of the effects of lead poisoning on the intelligence and neurobehavioral abilities of children. Theor. Biol. Med. Model. 10 (1).
19. Hsiao, C.-L., Wu, K.-H., Wan, K.-S., (2011). Effects of environmental lead exposure on T-helper cell-specific cytokines in children. J. Immunotoxicol. 8 (4), 284–287.
20. Hyder, O., Chung, M., Cosgrove, D., Herman, J.M., Li, Z., Firoozmand, A., Gurakar, A., Koteish, A., Pawlik, T.M., (2013). Cadmium exposure and liver disease among US adults. J. Gastrointest. Surg. 17 (7), 1265–1273.
21. Kabata- Pendia A 3rd, editor. Trace Elements in Soils and Plants. Boca Raton, FL: CRC Press; 2001
22. Karbowska, B., (2016). Presence of thallium in the environment: sources of contaminations, distribution and monitoring methods. Environ. Monit. Assess. 188 (11).
23. Kasten-Jolly, J., Heo, Y., Lawrence, D.A., (2010). Impact of developmental lead exposure on splenic factors. Toxicol. Appl. Pharmacol. 247 (2), 105–115.
24. Liaw, J., Marshall, G., Yuan, Y., Ferreccio, C., Steinmaus, C., and Smith, A. H. (2008). Increased childhood liver cancer mortality and arsenic in drinking water in northern Chile. Cancer Epidemiol. Biomarkers Prev. 17 (8), 1982–1987
25. Liu, C.-M., Ma, J.-Q., and Sun, Y.-Z. (2012). Puerarin protects rat kidney from leadinduced apoptosis by modulating the PI3K/Akt/eNOS pathway. Toxicol. Appl. Pharmacol. 258 (3), 330–342.
26. Loh, N., Loh, H.-P., Wang, L.K., Wang, M.H.S. (2016). Health effects and control of toxic lead in the environment. Nat. Resour. Control Process. 233–284.
27. Luo, L., Wang, B., Jiang, J., Huang, Q., Yu, Z., Li, H., et al. (2020). Heavy metal contaminations in herbal medicines: determination. comprehensive risk assessments. Front. Pharmacol. 11, 595335. doi:10.3389/fphar.2020.595335
28. Macklin MG, Thomas CJ, Mudbhatkal A et al (2023) Impacts of metal mining on river systems: a global assessment. Science 381:1345–1350.

29. Malaguarnera, G., Cataudella, E., Giordano, M., Nunnari, G., Chisari, G., Malaguarnera, M., (2012). Toxic hepatitis in occupational exposure to solvents. World J. Gastroenterol. 18, 2756–2766.
30. Mishra, K.P., (2009). Lead exposure and its impact on immune system: A review. Toxicol. Vitr. 23 (6), 969–972.
31. O'Neal, S.L., Zheng, W., (2015). Manganese toxicity upon overexposure: a decade in review. Curr. Environ. Heal. reports 2 (3), 315–328.
32. Packer, M., (2016). Cobalt cardiomyopathy: a critical reappraisal in light of a recent resurgence. Circ. Hear. Fail. 9 (12).
33. Park, Y.-H., Kim, D., Dai, J., Zhang, Z., 2015. Human bronchial epithelial BEAS-2B cells, an appropriate in vitro model to study heavy metals induced carcinogenesis. Toxicol. Appl. Pharmacol. 287 (3), 240–245.
34. Peters, J.L., Perlstein, T.S., Perry, M.J., McNeely, E., Weuve, J., (2010). Cadmium exposure in association with history of stroke and heart failure. Environ. Res. 110 (2), 199–206.
35. Qasem NA, Mohammed RH and Lawal DU (2021). Removal of heavy metal ions from wastewater: A comprehensive and critical review. Npj Clean Water. 4:1-15.
36. Reczek, C.R., Chandel, N.S., (2017). The two faces of reactive oxygen species in cancer. Annu. Rev. Cancer Biol. 1 (1), 79–98.
37. Rehman K, Fatima F, Waheed I and Akash M S (2018) Prevalence of exposure of heavy metals and their impact on health consequences. J. Cell. Biochem. 119, 157-184.
38. Reyes, J.L., Molina-Jijón, E., Rodríguez-Muñoz, R., Bautista-García, P., Debray-García, Y. Namorado, M.D.C., (2013). Tight junction proteins and oxidative stress in heavy metals-induced nephrotoxicity. Biomed Res. Int. 1–14.
39. Salama, A., Hegazy, R., and Hassan, A. (2016). Intranasal chromium induces acute brain and lung injuries in rats: assessment of different potential hazardous effects of environmental and occupational exposure to chromium and introduction of a novel pharmacological and toxicological animal model. PLoS One 11 (12), e0168688.
40. Sanchez T (2018). Effects of mercury, lead, arsenic and zinc to human renal oxidative stress and functions: a review. Arch Med. 4.
41. Silbergeld, E.K., Waalkes, M., Rice, J.M., (2000). Lead as a carcinogen: Experimental evidence and mechanisms of action. Am. J. Ind. Med. 38, 316–323
42. Sun, H., Brocato, J., Costa, M., (2015). Oral chromium exposure and toxicity. Curr. Environ. Heal. Rep. 2 (3), 295–303.

43. Tanaka, K.I., Kawahara, M., 2017. Copper enhances zinc-induced neurotoxicity and the endoplasmic reticulum stress response in a neuronal model of vascular dementia. Front. Neurosci. 11.
44. Tchounwou PB, Wilson B, Ishaque A (1999). Important considerations in the development of public health advisories for arsenic and arsenic-containing compounds in drinking water. Rev Environ Health. 14(4):211–229.
45. Tchounwou PB, Yedjou, CG, Patlolla AK, and Sutton DJ (2012). Heavy metal toxicity and the environment. Mol Clin Environ Toxicol.133-164.
46. Tellez-Plaza, M., Guallar, E., Howard, B.V., Umans, J.G., Francesconi, K.A., Goessler, W., Silbergeld, E.K., Devereux, R.B., Navas-Acien, A., (2013). Cadmium exposure and incident cardiovascular disease. Epidemiology 24 (3), 421–429.
47. Tellez-Plaza, M., Navas-Acien, A., Crainiceanu, C.M., Guallar, E., (2008). Cadmium exposure and hypertension in the 1999–2004 National Health and Nutrition Examination Survey (NHANES). Environ. Health Perspect. 116 (1), 51–56.
48. Tönnies E and Trushina E (2017). Oxidative stress, synaptic dysfunction, and Alzheimer's disease. J Alzheimer's Dis. 57:1105-1121.
49. Toxicological Profile for Cadmium, (2002). ATSDR's Toxicol. Profiles.
50. Velma V, Vutukuru SS, Tchounwou PB (2009). Ecotoxicology of hexavalent chromium in freshwater fish: a critical review. Rev Environ Health. 24(2):129–145.
51. Wang XF, Xing ML, Shen Y, Zhu X, Xu LH (2006). Oral administration of Cr (VI) induced oxidative stress, DNA damage and apoptotic cell death in mice. Toxicology. 228:16–23.
52. Wang, J., Zhu, H., Yang, Z., and Liu, Z. (2013). Antioxidative effects of hesperetin against lead acetate-induced oxidative stress in rats. Indian J. Pharmacol. 45 (4), 395–398.
53. Wang Y, Tang Y, Li Z, Hua Q, Wang L, Song X, Zou B, Ding M, Zhao J and Tang C (2020). Joint toxicity of a multi-heavy metal mixture and chemoprevention in sprague dawley rats. Int J Environ Res Pub Health. 17:1451.
54. Wilson DN Association Cadmium. Cadmium - market trends and influences; London. Cadmium 87 Proceedings of the 6th International Cadmium Conference; 1988. pp. 9–16.
55. Wu X, Cobbina SJ, Mao G, Xu H, Zhang Z, Yang L. A review of toxicity and mechanisms of individual and mixtures of heavy metals in the environment. Environ Sci Pollut Res Int. 2016;23(9):8244-59.
56. Yoshizawa, K., Rimm, E.B., Morris, J.S., Spate, V.L., Hsieh, C.C., Spiegelman, D., Stampfer, M.J., 2002. Mercury and the risk of coronary heart disease in men. N. Engl. J. Med. 347, 1755–1760.

57. Yu, L., Liou, I.W., Biggins, S.W., Yeh, M., Jalikis, F., Chan, L.-N., Burkhead, J., 2019. Copper deficiency in liver diseases: a case series and pathophysiological considerations. Hepatol. Commun. 3 (8), 1159–1165

Frontiers in Animal Biology and Biodiversity

ISBN: 978-93-49938-98-4| Year: 2026 | pp: 96 – 108

Emerging and Re-emerging Zoonotic Diseases: A One Health Perspective

[1]Isai M

[2]Annadurai T

[3]Nagaraj K

[4]Subavathy P

[1]Assistant Professor, Department of Zoology, Seethalakshmi Ramaswami College (Autonomous), Affiliated to Bharathidasan University, Tiruchirappalli, Tamilnadu

[2]Associate Professor, Centre for Excellence in Nanobio Translational Research, Department of Pharmaceutical Technology, University college of Engineering, Anna University- BIT campus, Tiruchirappalli - 620024, Tamilnadu

[3]Professor, Biomedical & Nano-Drug Formulation Laboratory, Department of General Medicine, Saveetha Medical College and Hospital, Saveetha Institute of Medical and Technical Sciences (SIMATS), Chennai-602105, Tamilnadu.

[4]Assistant Professor, PG and Research Department of Zoology, St. Mary's College (Autonomous), Thoothukudi, Tamilnadu

Email: mathivananisai@gmail.com

Article DOI Link: https://zenodo.org/uploads/21282200

DOI: 10.5281/zenodo.21282200

Abstract

Zoonotic diseases — infectious diseases transmitted between vertebrate animals and humans — account for approximately 60% of all emerging infectious diseases and pose one of the most pressing challenges to global health security. The One Health framework, which recognizes the inextricable interconnectedness of human, animal, and ecosystem health, provides the most coherent conceptual and operational paradigm for addressing these threats. This chapter examines the epidemiological, ecological, and socioeconomic drivers that facilitate the emergence and re-emergence of zoonotic pathogens, with reference to landmark outbreaks including Ebola virus disease, SARS-CoV-2, Nipah virus, and highly pathogenic avian influenza. We analyze the structural components of the One Health approach —

integrated surveillance, cross-sectoral governance, interdisciplinary research, and community-based prevention — and critically evaluate evidence for their effectiveness. The chapter further discusses the role of anthropogenic pressures such as land-use change, climate disruption, biodiversity loss, and intensified animal agriculture in amplifying spillover risk. We conclude that a fully operationalized One Health architecture, embedded in national and international policy, represents the most viable pathway toward pandemic preparedness and sustainable zoonotic disease control in the twenty-first century.

Keywords: zoonotic diseases, One Health, emerging infectious diseases, spillover, pandemic preparedness, eco health, surveillance, antimicrobial resistance

Introduction

Throughout recorded history, infectious diseases of animal origin have profoundly shaped human civilizations. From the Justinian Plague of the sixth century to the 1918 influenza pandemic, and from the AIDS epidemic to the COVID-19 pandemic of 2019–2023, pathogens originating in animal reservoirs have repeatedly demonstrated their capacity to destabilize societies, economies, and public health systems. Taylor et al. (2001) established in a landmark analysis that zoonoses constitute approximately 61% of all known human pathogens and account for 75% of emerging infectious diseases, a figure that has been widely corroborated by subsequent surveillance data. More recently, Woolhouse and Gowtage-Sequeria (2005) identified wildlife as the primary reservoir for the majority of emerging zoonoses, underscoring the centrality of the human–wildlife interface to global health risk.

The concept of One Health — formally articulated in the 2004 Manhattan Principles and subsequently endorsed by the World Health Organization (WHO), the Food and Agriculture Organization (FAO), and the World Organisation for Animal Health (WOAH, formerly OIE) — posits that human health, animal health, and ecosystem integrity are interdependent and must be managed in an integrated fashion (One Health High-Level Expert Panel [OHHLEP], 2022). This framework has gained renewed institutional traction following the COVID-19 pandemic, which exposed catastrophic gaps in global preparedness infrastructure and illustrated the consequences of siloed sectoral thinking in infectious disease management.

This chapter provides a comprehensive synthesis of the epidemiology, ecology, and public health dimensions of emerging and re-emerging zoonotic diseases through a One Health lens. We review the principal pathogens and disease categories of concern, analyze the complex web of anthropogenic drivers accelerating emergence risk, and evaluate the institutional and operational frameworks available for integrated response. The chapter draws on peer-reviewed literature, WHO and FAO situation reports, and case studies from Africa, Asia, and the Americas to illustrate key concepts.

Defining Zoonotic Diseases: Classification and Epidemiological Scope

The World Health Organization defines zoonoses as diseases and infections that are naturally transmitted between vertebrate animals and humans (WHO, 2020). This broad category encompasses bacterial, viral, parasitic, and prion diseases, and includes both direct transmission (contact with infected animals or their secretions) and indirect transmission via arthropod vectors, contaminated food, or water. The distinction between emerging and re-emerging zoonoses is epidemiologically and policy-relevant: emerging zoonoses are those appearing in a human population for the first time or spreading to new geographic regions, while re-emerging zoonoses are those that had previously declined in incidence but are resurging, often in new epidemiological contexts (Daszak et al., 2000).

Viruses, particularly RNA viruses, predominate among newly emerging zoonotic pathogens due to their high mutation rates and capacity for rapid host adaptation. Key viral families implicated include Coronaviridae (SARS, MERS, SARS-CoV-2), Filoviridae (Ebola, Marburg), Paramyxoviridae (Nipah, Hendra), and Orthomyxoviridae (influenza A). Among bacteria, Brucella spp., Coxiella burnetii (Q fever), and Leptospira spp. remain globally significant zoonotic burdens, particularly in low- and middle-income countries with close human–livestock interfaces. Parasitic zoonoses, including toxoplasmosis and echinococcosis, disproportionately affect immunocompromised populations and communities dependent on subsistence agriculture (Hotez et al., 2014).

Table 1

Disease	Reservoir Host	Transmission Route	Geographic Distribution
Ebola Virus Disease	Bats (fruit bats)	Direct contact with body fluids	Sub-Saharan Africa
SARS-CoV-2	Bats (probable)	Respiratory droplets/aerosols	Global (pandemic)
Nipah Virus	Pteropus bats	Bat-to-human via date palm sap	South/Southeast Asia
H5N1 Avian Influenza	Wild birds/poultry	Direct animal contact	Asia, Africa, Europe
Monkeypox (Mpox)	Rodents, primates	Close contact, respiratory	Global (2022 outbreak)
Lyme Disease	Ixodes ticks/deer	Vector-borne (tick bite)	North America, Europe
Rabies	Dogs, bats, wildlife	Animal bite/scratch	Asia, Africa (endemic)
Brucellosis	Cattle, goats, swine	Ingestion of raw dairy/meat	Mediterranean, Middle East

Table 1. Selected emerging and re-emerging zoonotic diseases: reservoir hosts, transmission routes, and geographic distribution. Adapted from WHO (2023) and Jones et al. (2008).

Anthropogenic Drivers of Zoonotic Disease Emergence

1. Land-Use Change and Deforestation

The encroachment of human settlements and agricultural activities into wildlife habitats represents perhaps the most potent proximate driver of zoonotic spillover. Deforestation for agriculture, logging, mining, and infrastructure development reduces the geographic buffering between wildlife reservoirs and human communities, increasing contact rates at the human–animal interface. Wolfe et al. (2005) demonstrated that bushmeat hunting and wildlife trade in forested regions of Central Africa serve as principal mechanisms for the introduction of novel simian retroviruses into human populations. The Nipah virus outbreaks in Malaysia (1998–1999), in which displaced Pteropus bats shed virus into pig farming operations as a result of forest loss driven by fires and drought, provide a canonical example of this pathway (Chua et al., 2002).

2. Climate Change and Ecological Disruption

Anthropogenic climate change is reshaping the geographic distribution, abundance, and phenology of disease vectors, reservoir hosts, and pathogens. Rising temperatures are expanding the range of Aedes and Culex mosquitoes, increasing the transmission windows for arboviruses including dengue, Zika, and West Nile virus. Climate-driven changes in precipitation patterns and flooding events are amplifying conditions favorable to leptospirosis and waterborne pathogens. Altizer et al. (2013) argued that climate change represents a fundamental threat to wildlife health and, by extension, to the zoonotic disease interface, through disrupting host–pathogen–vector relationships that have been evolutionarily stabilized over millennia. Seasonal shifts also affect the migratory routes of wild birds, altering the geographic scope of highly pathogenic avian influenza (HPAI H5N1) dissemination (Gilbert et al., 2008).

3. Intensified Animal Agriculture and Antimicrobial Resistance

Industrial-scale livestock production creates ecologically dense environments that amplify pathogen transmission, facilitate genetic reassortment of influenza viruses, and select for antimicrobial resistance. The global poultry industry, which confines billions of birds in high-density facilities, has been epidemiologically linked to repeated emergence and amplification events of HPAI strains. Furthermore, prophylactic and sub-therapeutic antibiotic use in livestock — estimated at over 73,000 tonnes annually (Van Boeckel et al., 2014) — is a primary driver of the global antimicrobial resistance (AMR) crisis, which itself constitutes a re-emerging threat of existential public health consequence. The convergence of zoonotic disease emergence and AMR within the One Health framework has been explicitly recognized by the Tripartite Alliance (WHO, FAO, WOAH) in their 2021 joint plan of action.

4. Global Mobility and Trade

International travel and global trade networks serve as force multipliers for zoonotic pathogen dispersal. The 2003 SARS epidemic demonstrated how a novel coronavirus could traverse continental boundaries within days via commercial aviation, transforming a localized wildlife spillover event in Guangdong Province into a 29-country outbreak (Peiris et al., 2003). The live animal trade — both legal and illicit — has been implicated in the introduction of monkeypox to North America in 2003, via African rodents, and continues to represent a major pathway for pathogen importation (Reed et al., 2004). Digital surveillance tools and genomic epidemiology have substantially improved the capacity to trace transboundary pathogen movement, yet regulatory frameworks governing wildlife trade remain fragmented and inadequate.

Figure 1




Figure 1. The One Health Triad: conceptual framework illustrating the intersecting domains of human health, animal health, and ecosystem health, with zoonotic diseases occupying the shared interface of all three. Adapted from OHHLEP (2022).

The One Health Framework: Principles, Architecture, and Implementation

1. Conceptual Foundations

The One Health concept is grounded in the recognition that the health of humans, animals, and ecosystems are not merely analogous but functionally interdependent. The OHHLEP (2022) definition formally adopted by the Tripartite Plus organizations specifies One Health as "an integrated, unifying approach that aims to sustainably balance and optimize the health of people, animals and ecosystems". This definition explicitly incorporates ecosystem health and environmental sustainability alongside the more traditional human–animal dyad, reflecting

growing scientific consensus that biodiversity conservation and ecological integrity are foundational prerequisites for long-term zoonotic disease control (Keesing et al., 2010).

The intellectual antecedents of One Health include the One Medicine concept proposed by veterinarian Calvin Schwabe in 1964, which emphasized the biological continuum between human and veterinary medicine. However, One Health extends beyond this clinical equivalence to encompass population-level surveillance, policy integration, and the ecological determinants of health (Zinsstag et al., 2011). The framework has found its most robust institutional expression in joint FAO–WHO–WOAH initiatives including the Global Early Warning System for major animal diseases (GLEWS) and the Joint External Evaluation (JEE) process for national health security capacities.

2. Integrated Surveillance Systems

Effective zoonotic disease surveillance requires simultaneous monitoring at the human–animal–environment interface. Traditional epidemiological surveillance, designed around human clinical presentations, consistently generates delayed outbreak detection when the primary amplification occurs in animal populations. The PREDICT project (2009–2019), funded by USAID, screened over 164,000 animals and people across 31 countries and identified 931 novel viruses, demonstrating the feasibility and value of proactive zoonotic pathogen discovery (Becker et al., 2020). Event-based surveillance systems, including the WHO's Global Outbreak Alert and Response Network (GOARN) and WOAH's World Animal Health Information System (WAHIS), have improved real-time information sharing, though integration across the human and animal sectors remains structurally incomplete in most national health architectures.

3. Cross-Sectoral Governance and Policy Integration

One Health operationalization requires institutional mechanisms that transcend traditional ministerial silos. National One Health Platforms, established in countries including Uganda, Sierra Leone, and Vietnam, provide interministerial coordination structures that facilitate joint surveillance, response planning, and resource allocation across the human health, animal health, and environment ministries. The International Health Regulations (IHR, 2005) provide the primary international legal framework for zoonotic event notification, but their implementation has been shown to be uneven, with many low- and middle-income countries lacking the core capacities required for timely and transparent reporting (Gostin et al., 2020). The Pandemic Treaty negotiations initiated by the WHO in 2021 represent a significant opportunity to strengthen the international governance architecture for zoonotic disease prevention, though their trajectory remains subject to geopolitical negotiation.

Table 2

Pillar	Core Activities	Key Mechanisms	Expected Outcomes
Surveillance	Integrated disease monitoring across sectors	Tri-sectoral data sharing (WHO, FAO, OIE)	Early outbreak detection
Prevention	Vaccination, biosecurity, habitat protection	Cross-species vaccine platforms	Reduced spillover events
Response	Coordinated multi-agency rapid response	Emergency Operation Centers	Contained outbreaks
Research	Interdisciplinary zoonotic disease studies	EcoHealth Alliance, PREDICT	Evidence-based policy
Governance	Policy integration across ministries	National One Health plans	Institutional collaboration
Capacity Building	Training across health disciplines	Field Epidemiology Programs	Workforce preparedness

Table 2. Core pillars of the One Health framework: activities, mechanisms, and expected outcomes for zoonotic disease prevention and control. Adapted from OHHLEP (2022) and Zinsstag et al. (2011).

Illustrative Case Studies: Lessons from Major Outbreaks

1. SARS-CoV-2 and the COVID-19 Pandemic

The COVID-19 pandemic, caused by the novel coronavirus SARS-CoV-2, represents the most consequential zoonotic emergence event of the twenty-first century, resulting in over 7 million confirmed deaths globally and trillions of dollars in economic losses (WHO, 2023). Genomic evidence strongly supports a bat origin for SARS-CoV-2, with Rhinolophus bats harboring the closest known viral relatives (Zhou et al., 2020). The precise pathway of spillover to humans, whether through an intermediate host or direct bat–human transmission, remains under active scientific investigation. The pandemic exposed the inadequacy of existing surveillance systems at the wildlife–human interface, the insufficient integration of veterinary and public health intelligence, and the consequences of delayed transparent reporting. It catalyzed unprecedented investment in One Health infrastructure and

research, including the establishment of the WHO Scientific Advisory Group for the Origins of Novel Pathogens (SAGO).

2. Nipah Virus: A Recurrent Regional Threat

Nipah virus, a paramyxovirus maintained in Pteropus fruit bats, has caused repeated outbreaks in Bangladesh (annually since 2001) and India, with case fatality rates ranging from 40% to 100% (Luby et al., 2009). The primary transmission pathway in Bangladesh involves human ingestion of raw date palm sap contaminated by bat saliva and urine. Limited human-to-human transmission has been documented in healthcare settings. The Nipah outbreaks illustrate the critical role of behavioral and ecological risk factors — particularly agricultural practices that create bat–human food-sharing opportunities — in sustaining zoonotic spillover, and demonstrate the value of community-based One Health interventions including targeted behavior change communication around date palm sap consumption.

3. Highly Pathogenic Avian Influenza

H5N1 highly pathogenic avian influenza (HPAI) has been detected in over 80 countries since 2003, causing hundreds of confirmed human infections with a case fatality rate exceeding 60% (WHO, 2024). The HPAI H5N1 clade 2.3.4.4b emerged in 2021 as a globally dominant strain, driving mass mortality in wild birds and poultry across multiple continents and raising pandemic concern due to evidence of spillover to mammals, including dairy cattle in the United States in 2024. A One Health response to HPAI requires coordinated wildlife surveillance, rapid culling and compensation schemes for poultry farmers, occupational health protection for agricultural workers, and sustained investment in influenza pandemic preparedness including cross-reactive vaccine platforms.

Figure 2




Figure 2. Conceptual framework illustrating the five major anthropogenic drivers of zoonotic disease emergence and their convergence at the human–animal–environment interface.

Challenges and Critical Perspectives

Despite its conceptual elegance and institutional momentum, the One Health framework faces substantial implementation challenges. First, there is a persistent resource asymmetry between human and animal health sectors, with most national health budgets heavily skewed toward human clinical infrastructure and offering minimal investment in veterinary epidemiology, wildlife disease monitoring, or environmental health surveillance (Grace, 2015). In many low- and middle-income countries, veterinary services are critically understaffed and underequipped, rendering the aspirations of integrated One Health surveillance practically unattainable without substantial external investment.

Second, governance fragmentation across ministerial portfolios creates coordination failures that impede timely joint responses. The entities responsible for human health, livestock, wildlife, and environment are often organizationally and politically siloed, with competing mandates and institutional incentives. Achieving genuine tri-sectoral integration requires political will at the highest levels of government, backed by dedicated joint financing mechanisms and legally mandated collaboration protocols (Rüegg et al., 2018).

Third, the growing burden of antimicrobial resistance at the human–animal interface constitutes a slow-burning zoonotic emergency that demands One Health governance with the same urgency as acute outbreak response. The WHO Global Action Plan on AMR (2015) and subsequent national action plans have made progress in acknowledging the livestock–human AMR nexus, but translating policy commitments into measurable reductions in veterinary antibiotic usage remains challenging in jurisdictions where agricultural economic interests are politically powerful (Van Boeckel et al., 2019).

Future Directions and Policy Recommendations

The future of zoonotic disease prevention lies in proactive rather than reactive paradigms. Early warning systems capable of detecting novel pathogen spillover events before they achieve epidemic potential require sustained investment in wildlife disease surveillance networks, environmental DNA monitoring, and bioinformatic capacity in endemic countries. The Global Virome Project, which aims to characterize the full diversity of zoonotic viruses in wildlife before they emerge in humans, represents an ambitious horizon for proactive discovery (Carroll et al., 2018).

At the policy level, the international community must capitalize on the post-COVID institutional momentum to embed One Health principles in binding international agreements, including the proposed Pandemic Accord, CITES regulations governing wildlife trade, and the Convention on Biological Diversity's Kunming-Montreal Global Biodiversity Framework. Biodiversity conservation must be explicitly framed and funded as a public health investment, given robust evidence

that intact, species-rich ecosystems provide regulatory services that dilute spillover risk (Keesing & Ostfeld, 2021).

Community engagement and social science research represent underinvested dimensions of One Health. Sustainable zoonotic disease control ultimately requires behavioral change at the community level — modified hunting practices, improved biosecurity at livestock–wildlife interfaces, safe food handling, and reduced antibiotic demand — that can only be achieved through culturally competent, co-designed public health programmes. Incorporating Indigenous and local ecological knowledge into surveillance and response frameworks can provide irreplaceable insights into wildlife ecology and human–animal contact patterns at hyperlocal scales (Mather & Marshall, 2011).

Conclusion

Emerging and re-emerging zoonotic diseases are not random, unpredictable misfortunes but the predictable consequence of accelerating disruptions to the human–animal–environment interface wrought by habitat destruction, climate change, industrial food systems, and global mobility. The One Health framework provides the most epistemologically coherent and operationally promising response architecture available. However, translating its principles into durable institutional arrangements, equitably financed surveillance systems, and community-anchored prevention programmes requires political commitment, interdisciplinary collaboration, and investment levels commensurate with the scale of the threat. The COVID-19 pandemic demonstrated, at catastrophic cost, the price of neglecting zoonotic interfaces. The imperative now is to treat pandemic prevention as a global public good and to invest in One Health infrastructure with the urgency and sustained commitment that the existential risk profile of zoonotic disease demands.

References

1. Altizer, S., Ostfeld, R. S., Johnson, P. T. J., Kutz, S., & Harvell, C. D. (2013). Climate change and infectious diseases: From evidence to a predictive framework. Science, 341(6145), 514–519. https://doi.org/10.1126/science.1239401
2. Becker, D. J., Albery, G. F., Sjodin, A. R., Poisot, T., Dallas, T. A., Eskew, E. A., Farrell, M. J., Guth, S., Han, B. A., Simmons, N. B., Carlson, C. J. (2020). Predicting wildlife hosts of betacoronaviruses for SARS-CoV-2 conservation risk assessment. eLife, 10, e65306. https://doi.org/10.7554/eLife.65306
3. Carroll, D., Daszak, P., Wolfe, N. D., Gao, G. F., Morel, C. M., Morzaria, S., Pablos-Méndez, A., Tomori, O., & Mazet, J. A. K. (2018). The Global Virome Project. Science, 359(6378), 872–874. https://doi.org/10.1126/science.aap7463
4. Chua, K. B., Bellini, W. J., Rota, P. A., Harcourt, B. H., Tamin, A., Lam, S. K., Ksiazek, T. G., Rollin, P. E., Zaki, S. R., Shieh, W.-J., Goldsmith, C. S., Gubler, D. J., Roehrig, J. T., Eaton, B., Gould, A. R., Olson, J., Field, H.,

Daniels, P., Ling, A. E., . . . Rota, P. A. (2002). Nipah virus: A recently emergent deadly paramyxovirus. Science, 288(5470), 1432–1435. https://doi.org/10.1126/science.288.5470.1432

5. Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2000). Emerging infectious diseases of wildlife — Threats to biodiversity and human health. Science, 287(5452), 443–449. https://doi.org/10.1126/science.287.5452.443
6. Gilbert, M., Xiao, X., Domenech, J., Lubroth, J., Martin, V., & Slingenbergh, J. (2008). Anatidae migration in the western Palearctic and spread of highly pathogenic avian influenza H5N1 virus. Emerging Infectious Diseases, 12(11), 1650–1656. https://doi.org/10.3201/eid1211.060223
7. Gostin, L. O., Habibi, R., & Mason Meier, B. (2020). Has global health law risen to meet the COVID-19 challenge? Revisiting the International Health Regulations to prepare for future threats. Journal of Law, Medicine & Ethics, 48(2), 376–381. https://doi.org/10.1177/1073110520935354
8. Grace, D. (2015). Review of evidence on antimicrobial resistance and animal agriculture in developing countries. Evidence on Demand. https://doi.org/10.12774/eod_cr.june2015.graced
9. Hotez, P. J., Alvarado, M., Basáñez, M.-G., Bolliger, I., Bourne, R., Boussinesq, M., Brooker, S. J., Brown, A. S., Buckle, G., Budke, C. M., Carabin, H., Coffeng, L. E., Fèvre, E. M., Fürst, T., Halasa, Y. A., Jasrasaria, R., Johns, N. E., Keiser, J., King, C. H., . . . Naghavi, M. (2014). The global burden of disease study 2010: Interpretation and implications for the neglected tropical diseases. PLOS Neglected Tropical Diseases, 8(7), e2865. https://doi.org/10.1371/journal.pntd.0002865
10. Jones, K. E., Patel, N. G., Levy, M. A., Storeygard, A., Balk, D., Gittleman, J. L., & Daszak, P. (2008). Global trends in emerging infectious diseases. Nature, 451(7181), 990–993. https://doi.org/10.1038/nature06536
11. Keesing, F., Belden, L. K., Daszak, P., Dobson, A., Harvell, C. D., Holt, R. D., Hudson, P., Jolles, A., Jones, K. E., Mitchell, C. E., Myers, S. S., Bogich, T., & Ostfeld, R. S. (2010). Impacts of biodiversity on the emergence and transmission of infectious diseases. Nature, 468(7324), 647–652. https://doi.org/10.1038/nature09575
12. Keesing, F., & Ostfeld, R. S. (2021). Impacts of biodiversity and biodiversity loss on zoonotic diseases. Proceedings of the National Academy of Sciences, 118(17), e2023540118. https://doi.org/10.1073/pnas.2023540118
13. Luby, S. P., Gurley, E. S., & Hossain, M. J. (2009). Transmission of human infection with Nipah virus. Clinical Infectious Diseases, 49(11), 1743–1748. https://doi.org/10.1086/647951
14. Mather, C., & Marshall, A. (2011). Living with disease: Biosecurity and avian influenza in humans. Sociology of Health & Illness, 33(2), 167–179. https://doi.org/10.1111/j.1467-9566.2010.01253.x

15. One Health High-Level Expert Panel (OHHLEP), Adisasmito, W. B., Almuhairi, S., Behravesh, C. B., Bilivogui, P., Bukachi, S. A., Casas, N., Becerra, N. C., Charron, D. F., Chaudhary, A., Dóniz, J. C., Doyon, J. F., Franka, R., Gao, G. F., Hayman, D. T. S., Khammavong, K., Kumaresan, J., . . . Zumla, A. (2022). One Health: A new definition for a sustainable and healthy future. PLOS Pathogens, 18(6), e1010537. https://doi.org/10.1371/journal.ppat.1010537
16. Peiris, J. S. M., Lai, S. T., Poon, L. L. M., Guan, Y., Yam, L. Y. C., Lim, W., Nicholls, J., Yee, W. K. S., Yan, W. W., Cheung, M. T., Cheng, V. C. C., Chan, K. H., Tsang, D. N. C., Yung, R. W. H., Ng, T. K., & Yuen, K. Y. (2003). Coronavirus as a possible cause of severe acute respiratory syndrome. The Lancet, 361(9366), 1319–1325. https://doi.org/10.1016/S0140-6736(03)13077-2
17. Reed, K. D., Melski, J. W., Graham, M. B., Regnery, R. L., Sotir, M. J., Wegner, M. V., Kazmierczak, J. J., Stratman, E. J., Li, Y., Fairley, J. A., Swain, G. R., Olson, V. A., Sargent, E. K., Kehl, S. C., Frace, M. A., Kline, R., Foldy, S. L., Davis, J. P., & Damon, I. K. (2004). The detection of monkeypox in humans in the Western Hemisphere. New England Journal of Medicine, 350(4), 342–350. https://doi.org/10.1056/NEJMoa032299
18. Rüegg, S. R., Nielsen, L. R., Buttigieg, S. C., Santa, M., Aragrande, M., Canali, M., Ehlinger, T., Chantziaras, I., Boriani, E., Radeski, M., Mindekem, R., Bhatt, V., & Häsler, B. (2018). A systems approach to evaluate One Health initiatives. Frontiers in Veterinary Science, 5, 23. https://doi.org/10.3389/fvets.2018.00023
19. Taylor, L. H., Latham, S. M., & Woolhouse, M. E. (2001). Risk factors for human disease emergence. Philosophical Transactions of the Royal Society B: Biological Sciences, 356(1411), 983–989. https://doi.org/10.1098/rstb.2001.0888
20. Van Boeckel, T. P., Brower, C., Gilbert, M., Grenfell, B. T., Levin, S. A., Robinson, T. P., Teillant, A., & Laxminarayan, R. (2014). Global antibiotic consumption 2000 to 2010: An analysis of national pharmaceutical sales data. The Lancet Infectious Diseases, 14(8), 742–750. https://doi.org/10.1016/S1473-3099(14)70780-7
21. Van Boeckel, T. P., Pires, J., Silveira, R., Zhao, C., Song, J., Criscuolo, N. G., Gilbert, M., Bonhoeffer, S., & Laxminarayan, R. (2019). Global trends in antimicrobial resistance in animals in low- and middle-income countries. Science, 365(6459), eaaw1944. https://doi.org/10.1126/science.aaw1944
22. Wolfe, N. D., Dunavan, C. P., & Diamond, J. (2007). Origins of major human infectious diseases. Nature, 447(7142), 279–283. https://doi.org/10.1038/nature05775

23. Woolhouse, M. E., & Gowtage-Sequeria, S. (2005). Host range and emerging and reemerging pathogens. Emerging Infectious Diseases, 11(12), 1842–1847. https://doi.org/10.3201/eid1112.050997
24. World Health Organization. (2020). Zoonoses (Fact sheet). https://www.who.int/news-room/fact-sheets/detail/zoonoses
25. World Health Organization. (2023). COVID-19 dashboard: Global situation. https://covid19.who.int/
26. World Health Organization. (2024). Influenza (avian and other zoonotic). https://www.who.int/news-room/fact-sheets/detail/influenza-(avian-and-other-zoonotic)
27. Zhou, P., Yang, X.-L., Wang, X.-G., Hu, B., Zhang, L., Zhang, W., Si, H.-R., Zhu, Y., Li, B., Huang, C.-L., Chen, H.-D., Chen, J., Luo, Y., Guo, H., Jiang, R.-D., Liu, M.-Q., Chen, Y., Shen, X.-R., Wang, X., . . . Shi, Z.-L. (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature, 579(7798), 270–273. https://doi.org/10.1038/s41586-020-2012-7
28. Zinsstag, J., Schelling, E., Waltner-Toews, D., & Tanner, M. (2011). From "one medicine" to "one health" and systemic approaches to health and well-being. Preventive Veterinary Medicine, 101(3–4), 148–156. https://doi.org/10.1016/j.prevetmed.2010.07.003

Frontiers in Animal Biology and Biodiversity

ISBN: 978-93-49938-98-4| Year: 2026 | pp: 109 – 118

Heavy Metals and Emerging Contaminants in Drinking Water: Ecological Risks and Health Implications

Anup Kumar Verma

Dr. Abhinav Singh

Department of Zoology, Acharya Narendra Dev Kisan P.G. College, Babhnan, Gonda, U.P, India

Email: anupk.verma6@gmail.com

Article DOI Link: https://zenodo.org/uploads/21282272

DOI: 10.5281/zenodo.21282272

Abstract

The presence of heavy metals and emerging contaminants (ECs) in drinking water is a converging global environmental and public health concern, resulting from natural geogenic sources, increased industrialization, urbanization and inadequate treatment of drinking water and wastewater. Exposure to toxic metals, such as arsenic (As), cadmium (Cd), lead (Pb), mercury (Hg) and chromium (Cr), as well as to organic ECs including pharmaceuticals, personal care products, artificial sweeteners, endocrine disrupting compounds (EDCs), pesticides and industrial chemicals, have been linked to carcinogenic, neurotoxic, nephrotoxic, reproductive and developmental effects in humans and disruption of aquatic ecosystems.pubmed.ncbi.nlm.nih+8

When considering direct human health risks through the drinking water pathway, the ecological risk assessment results using RQs, SSDs and probabilistic models are consistently medium to high at environmentally relevant concentrations of some of the ECs. This report summarises the existing evidence on the occurrence, sources, fate, ecological risk, human health impacts and regulatory responses to heavy metals and ECs in drinking water, and highlights some areas of research and policy opportunity for future work.

Keywords: emerging contaminants, arsenic (As), cadmium (Cd), lead (Pb), mercury (Hg) and chromium (Cr)

Introduction

Context and Definitions

The safety of drinking water is an essential component of human health; chemical contaminants such as heavy metals and organic micropollutants can add to the chronic disease burden in addition to microbial contaminants. Heavy metals are relatively heavy and dense metallic elements that, in their toxic form, are toxic or carcinogenic at low concentration, particularly if they are mobilized in water as a result of anthropogenic activities. Emerging contaminants are also known as emerging pollutants or contaminants of emerging concern (CEC), and include a wide range of synthetic and naturally occurring chemicals found in water at trace amounts but not well regulated.

Scope of the Problem

Heavy metal contamination of drinking water is well documented globally, both within urban and rural areas, including in some areas where mining, industrial activity or geogenic arsenic in ground water can lead to concentrations well above guideline levels. In contrast, ECs have been reported in surface water sources, reservoirs, drinking and distribution water, and groundwater in a wide variety of ecosystems, indicating a constant source of input and poor removal by conventional water treatment processes. Both contaminant groups can coexist and pose concern due to the possibility of mixture toxicity and cumulative risk, but most regulatory programs focus on single substances (Ali et al., 2019).

Occurrence and Sources in Drinking Water

1. Heavy Metals: Occurrence Patterns

Various studies have found high concentrations of As, Cd, Pb, Hg, Cr, and other metals in drinking water sources such as wells, boreholes, drinking water supplies, bottled or sachet water, and found that the concentrations vary from region to region and are related to land use and industrial emissions. For instance, the mean concentration of Cd, Pb, Cr, As, Al, Fe and Hg was found to be higher in certain zones than the national/ international guidelines in Gujranwala, Pakistan, and the strong correlation between certain metals like As-Hg indicated a co mobilization pathway or a common source for those metals. Hazard quotients (HQs) are used to evaluate human health risk from exposure to various drinking water sources to children and non-cancer risks in Nigeria and India, respectively, when HQs are below acceptable limits for the metal (Pb, Cd, or Hg).Hazard quotients (HQs) have been used to assess human health risk to children from exposure to various drinking water sources in Nigeria and cancer risk in India, respectively, where HQs are above acceptable limits for the metal (Pb, Cd, or Hg) (Briffa et al., 2020).

2. Sources of Heavy Metal Contamination

Some of the main sources of drinking water for contaminants known as heavy metals include:

Natural geogenic sources: Weathering of metal rich rocks and soils, geothermal activity, mobilization of groundwater, especially of As, F and Mn. Industrial and mining activities: untreated or inadequately treated effluents from smelters, tanneries, electroplating, textile, engineering and iron and Steel industries. Apllication of phosphate fertilizers, some pesticides with Cd, As, and other metals that result in leaching in groundwater and surface sources related to agriculture. Urban runoff and waste disposal: leaching from landfills, domestic sewage and corroding plumbing materials (such as Pb pipes) that result in metal load in distribution systems. These sources lead to chronic contamination and metals can remain in sediments and biota due to their non-degradability and bioaccumulation potential.

3. Emerging Contaminants: Occurrence Patterns

ECs have been found in rivers, lakes, reservoirs and groundwater from drinking water sources around the world at concentration levels ranging from $ng \cdot L^{-1}$ to low $\mu g \cdot L^{-1}$ and at high detection frequencies and in complex mixtures. In India, 15 pharmaceuticals and five artificial sweeteners were detected in river and groundwater (untreated as drinking water) in the Ganges River Basin, with total concentrations of PPCPs up to 826 $ng \cdot L^{-1}$ and artificial sweeteners up to 102 $ng \cdot L^{-1}$ in river water. In one study of a drinking water source in China, 346 pollutants were detected using non-targeted screening, with a majority of the most abundant being industrial chemicals, pharmaceuticals, and pesticides, and a few dozen of the more common ECs (such as bisphenol A, nonylphenol, atrazine, and ibuprofen) quantified at median concentrations ranging from approximately 8 $ng \cdot L^{-1}$ to 35 $ng \cdot L^{-1}$ (Diamanti-Kandarakis et al., 2009).

4. Sources of Emerging Contaminants

Unmetabolized pharmaceuticals, personal care products, hormones and artificial sweeteners in municipal wastewater effluents from WWTFs to receiving waters. Veterinary medication, pesticides, herbicides, and plasticizers from film and packaging that are carried by agricultural runoff. Specialty chemicals, surfactants, plasticizers (including bisphenol A) and intermediates from industrial effluents. Diffuse contamination from leaching from solid waste and septic systems into the groundwater. Some endocrine disrupting compounds, such as BPA, nonylphenol, estradiol, and a few pesticides, are poorly removed during wastewater and drinking water treatment processes, and can enter drinking water systems at times of episodic contamination, particularly in low flow conditions in rivers (Jarup, 2003).




Figure 1: Occurrence and Sources in Drinking Water

Regulatory Guideline Values and Standards

1. Guidelines for Heavy Metals in Drinking Water

The World Health Organization (WHO) and national drinking water regulators have established guideline values for metals in drinking water to reduce chronic exposures. The reported values for the guidelines given in recent literature are: 10 $\mu g \cdot L^{-1}$ for As, 3 $\mu g \cdot L^{-1}$ for Cd, 10 $\mu g \cdot L^{-1}$ for Pb and 50 $\mu g \cdot L^{-1}$ for Cr, derived from toxicological data and other health-based and acceptable daily intake (ADI) assumptions. Exceedances of these thresholds are regularly reported in contaminated regions, particularly in areas with ground water of high As concentration and in industrialised regions, suggesting that these thresholds are not being adequately met and that they need to be tightened.

2. Regulatory Approaches to Emerging Contaminants

Most ECs do not have comprehensive drinking water standards; responses are limited to priority lists, watch lists, and environmental quality standards for specific compounds. River catchments can reach or surpass river standards or screening values (often in the low $\mu g \cdot L^{-1}$ to $ng \cdot L^{-1}$ range) for specific EDCs, including BPA, nonylphenol, and estradiol, proposed by the European Union, Canada, and some national agencies. Normative frameworks often use risk-based approaches that take into account the toxicological potency of the substances, environmental occurrence and also feasibility of treating the substances but a large number of ECs and few toxicity data impose significant challenges (Richardson & Kimura, 2020).

Ecological Risks in Aquatic Systems

1. Risk Assessment Frameworks

The risk quotient (RQ) approach is often used for ecological risk assessment, where RQ is the ratio of the measured environmental concentration to the predicted no effect concentration (PNEC), and the probabilistic approach is often used, for example the SSD based on the joint probability curves and the margins of safety. These frameworks include species specific sensitivity distributions and can assess mixture risks, though there are data gaps and variation in exposure concentration that cause uncertainty. PubMed.

2. Ecological Risks of Heavy Metals

In aquatic organisms, heavy metals cause toxicity on plankton, invertebrates, fish and benthic organisms through various mechanisms such as oxidative stress, ion regulation, and enzyme and endocrine pathway interference. The elevated levels of Cd, Pb, Hg and As in drinking water sources downstream of industrial areas are indicated by the reduction in biodiversity and the deterioration of ecosystem functions, but most risk assessments focus on human exposures rather than on ecosystems. Sediments are frequently a reservoir for metals and a source of chronic exposure to benthic communities, and can remobilize under changing redox or hydrological conditions (Schwarzenbach et al., 2010).

3. Ecological Risks of Emerging Contaminants

The medium to high ecological risks has been demonstrated for ECs, especially PPCPs, artificial sweeteners and EDCs at environmental concentrations. In the Ganges River Basin, for example, moderate risk to aquatic organisms was determined using concentrations of caffeine and the concentrations of sulfamethoxazole, triclocarban, triclosan, and sucralose in river water and species-specific toxicity data. Risk quotients were above 1 for BPA, nonylphenol, atrazine and related ECs for certain sampling points in the Pearl River Delta drinking water source area, suggesting a high ecological risk. Canadian lakes have been identified as some of the highest risk compounds (compounds with RQ values > 1), and exposure patterns have been found to be spatially heterogeneous for some compounds, such as drospirenone, which needs to be targeted for mitigation (Zhang et al., 2008).

4. Mixture Toxicity and Food Web Effects

One of the most important ecosystem parameters in regard to ecological risk is mixture toxicity—the simultaneous presence of more than one EC and metal and their additive or synergistic effects on the target organisms. Study results incorporating trophic state indicators with SSD RQ techniques in tropical reservoirs indicate that relationships between eutrophication, contaminant exposure and community level responses may exist, where ECs can influence species composition

and energy flow. The non-targeted screening of hundreds of pollutants in drinking water sources highlights the fact that the risk assessment based on a limited number of pollutants can underestimate the overall ecological risk.




Figure 2: Ecological Risks in Aquatic Systems

Human Health Implications

1. Health Effects of Heavy Metals

There are several organ toxicities and chronic diseases linked to exposure to high levels of certain metals in drinking water. As, Cd, Pb, Hg, and Cr have been epidemiologically and toxicologically associated with an increased risk of cardiovascular disease, diabetes, kidney damage, neurologic disease and several types of cancer, such as skin, lung and kidney cancer. Many metals generate reactive oxygen species, oxidative stress, DNA damage and disruption of cell signaling pathways, which lead to carcinogenesis and organ dysfunction. Children are especially vulnerable to Pb and Hg exposure particularly through contaminated water because they have increased body weight and developing organ systems, and are linked with cognitive deficits, behavioral problems, and permanent neurodevelopmental deficits (WHO, 2022).

2. Quantitative Health Risk Assessments

Hazard quotient (HQ) and hazard index (HI) have been used to assess metals in drinking water in several studies and showed that there were high non-carcinogenic and carcinogenic risks in the impacted areas.In some studies, hazard quotient (HQ) and hazard index (HI) were applied to the drinking water metal concentrations and the results indicated considerable non-carcinogenic and carcinogenic risks in the impacted areas. For instance, the HQ and HI values for children are higher than those for adults in both states in Nigeria and Maharashtra in India, as a result of measured concentrations for Pb, Cd and Hg across various water sources, with the LCR values for As and Pb in some cases being above acceptable risk ranges in both

states. The results would support targeted interventions such as source remediation and advanced treatment, especially for informal water systems and self-supplied wells where there is limited regulatory control (UNEP, 2019).

3. Health Effects of Emerging Contaminants

Determining the human health risk from ECs in drinking water is more challenging given low levels, the fact they are diverse in structure and mode of action, and the lack of sufficient toxicity data, but there is emerging evidence for endocrine, reproductive, developmental, and metabolic effects. Some EDCs like BPA, nonylphenol, estradiol and some pesticides can mimic or interfere with endogenous hormones and can lead to reproductive abnormalities, hormone dependent cancers, metabolic disorders and developmental defects. The drinking water supply systems reviews highlight that the exposure pathway of tap water is relevant and recommend human health risk assessment in drinking water systems based on multi barrier approaches and NABC (Needs, Approaches, Benefits, Challenges) frameworks for guiding management.

4. Perceived vs Actual Risk

Research on perceived vs. actual risks of EDCs in drinking water shows that the public is more concerned with microbial contamination and aesthetic characteristics, and tends to be underestimating the chronic chemical risks, whereas regulatory agencies indicate that existing treatment technologies have the potential to lower the levels of EDCs found in drinking water to below recommended levels in many systems. Trace levels of EDCs are found throughout the environment, but exposure through drinking water may not pose a significant risk to humans compared to other exposure routes; however, the combined effects of multiple exposures and sensitive populations suggest precautionary measures are warranted (Sharma et al., 2014).

Treatment Technologies and Mitigation Strategies

1. Conventional and Advanced Treatment for Heavy Metals

Conventional drinking water treatment technologies (coagulation, sedimentation, filtration, and disinfection) may be able to remove some particulate bound metals but are less effective for dissolved forms of metals; optimized coagulation, lime softening and ion exchange treatment may be necessary to remove the metals. Activated carbon, metal oxide sorbents, and biosorbents have shown high removal efficiencies for As, Pb, Cd, and other metals, but are not yet cost-effective or feasible for concentrate management. Membrane technologies like reverse osmosis and nanofiltration are also very effective for removing As, Pb, Cd, and other metals; however, cost remains a challenge and concentrate management may be difficult. Point of use technology such as reverse osmosis units, activated carbon filters and

certain adsorptive material is recommended at the household level to minimize metal exposure in contaminated areas (Jarup, 2003).

2. Treatment of Emerging Contaminants

Conventional treatment regimens are not effective for many organic micropollutants (OMPs) and other advanced oxidation processes (AOPs), ozonation, UV/ H_2O_2, adsorption on activated carbon and high-pressure membranes are more effective for the removal of such contaminants. Multi barrier systems (biologically active filtration, carbon adsorption, and oxidation) have been found to effectively reduce PPCPs and EDCs in wastewater and drinking water systems, although some PPCPs are transformed and some partial removal occurs which complicates risk assessments. Tailored adsorbents, advanced biofiltration and hybrid processes are being explored to enhance removals and decrease energy consumption as emerging technologies (Schwarzenbach et al., 2010).

Research Gaps and Future Directions

1. Data Gaps and Monitoring Needs

There are significant information gaps about the occurrence and fate of many ECs, toxicity of mixtures of metals and organics, and long-term health effects of chronic low-level exposures. Non targeted screening studies identify hundreds of compounds in drinking water sources that have not yet been monitored, highlighting the need for monitoring strategies that combine targeted and non-targeted monitoring. High-resolution spatial and temporal data are needed in under-studied areas such as rural and informal water systems for conducting comprehensive health risk assessments and designing interventions for heavy metals.

2. Methodological Advances in Risk Assessment

The future risk assessment frameworks should include the consideration of probabilistic approaches, Adverse Outcome Pathways (AOPs) and mixture toxicity modeling to account for the uncertainties in EC and metal exposure. More holistic decision making can be supported by integrating ecological indicators, such as trophic state indexes, species sensitivity distributions and human health indicators and socio-economic vulnerability assessments. Linking environmental monitoring, toxicology, epidemiology and social sciences is important for understanding exposure pathways and risk perception as well as for the design of effective communication and governance strategies.

3. Implications for Low- and Middle-Income Regions

Heavy metals and ECs in drinking water are compounded by limited water infrastructure and informal water supply systems and significant reliance on untreated groundwater in many LMICs, increasing health and ecological risk. However, research in South Asia, Africa, and Latin America shows that

marginalized communities who use commercially unregulated surface water or self-supplied wells are exposed to disproportionate levels of metals and ECs, but there is limited monitoring and enforcement. There is a need to build capacity in laboratories and utilities, invest in treatment and distribution systems, and implement community interventions such as POU technologies and awareness creation campaigns to address these inequities.

Conclusion

The presence of heavy metals and emerging contaminants in drinking water systems is a multifaceted and complex environmental and public health problem with multiple sources, varying toxicity, and high levels of exposure and risk uncertainty. Heavy metals are common and often found at levels above guidelines in industrial and geogenic hotspots around the world, with the potential for increased cancer and non-cancer risks in vulnerable groups and children. ECs are broadly present at trace levels, and many present ecological risk assessments have found moderate to high risks to aquatic organisms, while some are identified as emerging concerns for human endocrine, reproductive and developmental effects.

Integrated strategies that involve upstream source control, advanced multi barrier treatment, comprehensive monitoring, probabilistic risk assessment and equitable governance of vulnerable communities are necessary for mitigation. Ongoing research on mixture toxicity, long term health effect and fate and transport of contaminants, and responsive regulatory measures and public communication will be vital to protect ecological health and human health as contaminant profiles in drinking water systems continue to change.

References

1. Ali, H., Khan, E., & Ilahi, I. (2019). Environmental chemistry and ecotoxicology of hazardous heavy metals: Environmental persistence, toxicity, and bioaccumulation. Journal of Chemistry, 2019, Article 6730305. https://doi.org/10.1155/2019/6730305
2. Briffa, J., Sinagra, E., & Blundell, R. (2020). Heavy metal pollution in the environment and their toxicological effects on humans. Heliyon, 6(9), e04691. https://doi.org/10.1016/j.heliyon.2020.e04691
3. Diamanti-Kandarakis, E., Bourguignon, J. P., Giudice, L. C., et al. (2009). Endocrine-disrupting chemicals: An Endocrine Society scientific statement. Endocrine Reviews, 30(4), 293–342. https://doi.org/10.1210/er.2009-0002
4. Jarup, L. (2003). Hazards of heavy metal contamination. British Medical Bulletin, 68(1), 167–182. https://doi.org/10.1093/bmb/ldg032
5. Richardson, S. D., & Kimura, S. Y. (2020). Water analysis: Emerging contaminants and current issues. Analytical Chemistry, 92(1), 473–505. https://doi.org/10.1021/acs.analchem.9b05269

6. Sharma, B., Singh, S., & Siddiqi, N. J. (2014). Biomedical implications of heavy metals induced imbalances in redox systems. BioMed Research International, 2014, Article 640754. https://doi.org/10.1155/2014/640754
7. United Nations Environment Programme. (2019). Global chemicals outlook II: From legacies to innovative solutions. United Nations Environment Programme. https://www.unep.org/resources/report/global-chemicals-outlook-ii
8. World Health Organization. (2022). Guidelines for drinking-water quality (4th ed., incorporating the 1st and 2nd addenda). World Health Organization. https://www.who.int/publications/i/item/9789240045064
9. Zhang, Y., Geißen, S. U., & Gal, C. (2008). Carbamazepine and diclofenac: Removal in wastewater treatment plants and occurrence in water bodies. Chemosphere, 73(8), 1151–1161. https://doi.org/10.1016/j.chemosphere.2008.07.086
10. Schwarzenbach, R. P., Egli, T., Hofstetter, T. B., von Gunten, U., & Wehrli, B. (2010). Global water pollution and human health. Annual Review of Environment and Resources, 35, 109–136. https://doi.org/10.1146/annurev-environ-100809-125342

ABOUT THE EDITORS



Dr. Esha Yadav

She is presently working as a Professor in the Department of Zoology at Brahmanand P.G. College, The Mall, Kanpur. She has completed her M. Phil. (Gold Medalist) and Ph.D. from the School of Life Sciences, Dr. B. R. A. University, Agra. She has qualified NET-JRF examination conducted by UGC-CSIR. She has 18 years of teaching experience. Her area of research is Environmental Toxicology. She has published more than 30 research papers in reputed National and International journals. She is also the author of 3 books and 6 book chapters. She has attended many National and International Conferences and presented her research work and also organized National Seminars sponsored by various agencies. She is guiding 5 Ph.D. students. She is a life member of 10 professional organisations and learned societies of national and international repute. She has bestowed with fellow awards from different societies and academies viz., Zoological Society of India, The Academy of Environmental Biology, Environment and Social Welfare Society, Society of Life Sciences, Asian Biological Research Foundation, Society of Biological Sciences and Rural Development, Scholars Academic and Scientific Society etc. She is an editor and reviewer for many reputed journals. She has awarded by many scientific organizations for her outstanding academic and research contributions. Congress of Zoology Medal for Outstanding Research & Academic Contribution (2018), Best Faculty Award for Outstanding Contribution in Zoology (2019), ESW Excellency Award for excellence in leadership and co-operation (2021), Paryavaran Shri Samman (2021), Best Academician of the Year Award (2022) are some of them.



Ms. Hemangi Jayram Gavit

She is an academician and researcher currently serving as an Assistant Professor at Rayat Shikshan Sanstha's Annasaheb Awate Arts, Commerce & Hutatma Babu Genu Science College, Manchar, Maharashtra. With six years of dedicated teaching experience, she combines classroom instruction with an active research trajectory. She holds an M.Sc. in Biotechnology from the prestigious Maharaja Sayajirao University of Baroda, Gujarat, and is currently advancing her research pursuits by completing her Ph.D. from Modern College, Ganeshkhind, Pune.A committed contributor to her scientific field, Ms. Gavit has authored seven research articles in reputed, peer-reviewed international journals, contributed four academic book chapters, and holds a registered patent reflecting her innovative approach to the biological sciences. Beyond her own research, she actively mentors undergraduate and postgraduate students in executing their research projects. Her professional commitment is further demonstrated by her participation in International Conferences, her role as an invited guest speaker, and her completion of formal UGC-aligned Orientation and Refresher courses, alongside specialized Faculty Development Programs organized by the Maharashtra State Faculty Development Academy (MSFDA).



Dr. Swapna Sachin Patil

She is serving as Assistant Professor and Head of the Department of Zoology at Rayat Shikshan Sanstha's Annasaheb Awate Arts, Commerce and Hutatma Babu Genu Science College, Manchar, Dist. Pune. She possesses 14 years of teaching and research experience in the field of Zoology, with a specialization in Hydrobiology. Dr. Patil has made notable contributions to scientific research through the publication of 16 research articles in reputed peer-reviewed and UGC CARE-listed journals. She is also actively involved in the academic peer-review process and serves as a reviewer for international journals. Her dedication to teaching, research, and academic excellence has established her as a committed educator and researcher in the field of biological sciences.



Mr. Anup Kumar Verma

Mr. Anup Kumar Verma is a research scholar in Zoology, pursuing his Ph.D. at Dr. Rammanohar Lohia Avadh University, Ayodhya. He holds a Master's degree in Zoology and has strong academic and research expertise in environmental biology, biodiversity and aquatic ecosystems. He has authored several book chapters with national and international publishers and published review articles in reputed journals. He received the International Academic Excellence Award 2026(Young Researcher Award) from the Bhagyashree Foundation for Academic Research and Development, Dhanabad, Jharkhand, in collaboration with Govt. Bilasa Girls Post Graduate Autonomous college, C.G. His research interests include climate change, water pollution, biodiversity conservation and environmental toxicology studies.

Nature Light Publications

309 West 11, Manjari VSI Road, Manjari Bk.,
Haveli, Pune- 412 307.
Website: www.naturelightpublications.com
Email: naturelightpublications@gmail.com
Contact No: +919822489040 / 9922489040

Nature - 119

ISBN: 978-93-49938-98-4

Price- 750/-